

## A Retrospective Study Re-Suicidal Behavior among Adult Psychiatric Inpatients in Iran

Saeed Shoja Shafte\*, Alireza Memarie, Masomeh Rezaie and Masomeh Hamidi

\*Department of Psychiatry, University of Social Welfare and Rehabilitation Sciences (USWR), Razi Psychiatric Hospital, Tehran, Iran.

Received July 17, 2019; Accepted August 20, 2019; Published January 22, 2020

### ABSTRACT

**Introduction:** Suicidal behavior is seen in the context of a variety of mental disorders and while many believe that, in general, first episode psychosis is a particularly high-risk period for suicide, no general agreement regarding higher prevalence of suicide in first episode psychosis is achievable. In the present study, suicides and suicide attempts among psychiatric in-patients has been evaluated to assess the general profile of suicidal behavior among native psychiatric inpatients and probing any relationship between serum cholesterol level and suicidal behavior.

**Methods:** Five acute academic wards, which have been specified for admission of first episode adult psychiatric patients, and five acute non-academic wards, which have been specified for admission of recurrent episode adult psychiatric patients, had been selected for current study. All inpatients with suicidal behavior (successful suicide and attempted suicide, in total), during the last five years (2013-2018), had been included in the present investigation. Also, assessment of serum lipids, including triglyceride, cholesterol, low density lipoprotein and high density lipoprotein, had been accomplished, for comparing the suicidal subjects with non-suicidal ones.

**Results:** Among 19160 psychiatric patients hospitalized in Razi psychiatric hospital during a sixty months period, 63 suicidal behaviors, including one successful suicide and sixty two suicide attempts, had been recorded by the safety board of hospital. The most frequent mental illness was bipolar I disorder, which was significantly more prevalent in comparison with other mental disorders ( $p < 0.04$ ,  $p < 0.02$ ,  $p < 0.007$  and  $p < 0.003$  in comparison with schizophrenia, depression, personality disorders and substance abuse, respectively). Self-mutilation, self-poisoning and hanging were the preferred methods of suicide among 61.11%, 19.44% and 19.44% of cases, respectively. In addition, no significant difference was evident between the first admission and recurrent admission inpatients, totally and separately, particularly with respect to psychotic disorders. Besides, with respect to different components of serum lipids, no specific or significant pattern was evident.

**Conclusion:** While in the present study the suicidal behavior was significantly more evident in bipolar disorder in comparison with other psychotic or no-psychotic disorders, no significant difference was evident between first admission and recurrent admission psychiatric inpatients. Moreover, no significant relationship between suicidal behavior and serum lipids was palpable.

**Keywords:** Psychiatric disorders, Suicide, Suicide attempt, First admission, Recurrent admission, Schizophrenia, Bipolar disorder, Depression, Substance abuse disorder

### INTRODUCTION

Suicide is derived from the Latin word for “self-murder.” It is a fatal act that represents the person's wish to die [1]. Approximately 25%-30% of persons who attempt suicide will go on to make more attempts. Suicidal behavior is seen in the context of a variety of mental disorders, most commonly bipolar disorder, major depressive disorder, schizophrenia, schizoaffective disorder, anxiety disorders, substance use disorders, borderline personality disorder, antisocial personality disorder, eating disorders and adjustment disorders [1]. According to the findings of a

**Corresponding author:** Saeed Shoja Shafte, Professor of Psychiatry, University of Social Welfare and Rehabilitation Sciences (USWR), Razi Psychiatric Hospital, Tehran, Postal Code: 18669-58891, P.O. Box: 18735-569, Iran, Tel: 0098-21-33401220; Fax: 0098-21-33401604; E-mail: ssshafte@gmail.com

**Citation:** Shafte SS, Memarie A, Rezaie M & Hamidi M. (2020) A Retrospective Study Re-Suicidal Behavior among Adult Psychiatric Inpatients in Iran. *J Psychiatry Psychol Res*, 3(1): 123-129.

**Copyright:** ©2020 Shafte SS, Memarie A, Rezaie M & Hamidi M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

study, among male psychiatric population, the absolute risk of suicide was highest for bipolar disorder, followed by unipolar affective disorder and schizophrenia. Among female psychiatric population, as well, the highest risk was found among women with schizophrenia, followed by bipolar disorder [2]. According to data, approximately 5%-6% of individuals with schizophrenia die by suicide, about 20% attempt suicide on one or more occasions, and many more have significant suicidal ideation. Suicidal behavior is sometimes in response to command hallucinations to harm oneself or others. Suicide risk remains high over the whole lifespan for males and females, although it may be especially high for younger males with comorbid substance use. Other risk factors include having depressive symptoms or feelings of hopelessness and being unemployed and the risk is higher, also, in the period after a psychotic episode or hospital discharge [3]. It is interesting that Bleuler [4] had drawn clinicians' attention that the most serious of the schizophrenic symptoms is the suicidal drive. Up to 50 percent of suicides among patients with schizophrenia occur during the first few weeks and months after discharge from a hospital; only a minority commit suicide while inpatients [3]. Having three or four hospitalizations during their 20s probably undermines the social, occupational and sexual adjustment of possibly suicidal patients with schizophrenia. Consequently, potential suicide victims are likely to be male, unmarried, unemployed, socially isolated, and living alone-perhaps in a single room. After discharge from their last hospitalization, they may experience a new adversity or return to ongoing difficulties. As a result, they become dejected, experience feelings of helplessness and hopelessness, reach a depressed state, and have, and eventually act on, suicidal ideas [5]. Abrupt discontinuation of medication, poor treatment compliance, social isolation, and increased expectation of good performance from others and from patients themselves, are risk factors for suicide in schizophrenics [6]. Also, the lifetime risk of suicide in individuals with bipolar disorder is estimated to be at least 15 times that of the general population. In fact, bipolar disorder may account for one-quarter of all completed suicides [7]. Many believe that, in general, first episode psychosis (FEP) is a particularly high-risk period for suicide, in which risk elevates by 60% within a first year of treatment as compared to later stages of illness. In this regard, longer duration of untreated psychosis, greater symptoms of depression and positive symptoms of psychosis were found to increase the odds of experiencing suicidal ideation [8]. While according to some studies depressive symptoms during the index psychotic episode and comorbidity with stimulant abuse at baseline were relevant predictive factors for suicidal behavior during the first years of first affective and non-affective psychotic episodes [9], more depressive symptoms, higher insight, and negative beliefs about psychosis increase the risk for suicidality in FEP [10]. Impulsive behavior such as self-harm, as well as having a family history of severe mental disorder or substance use,

have been stated as important risk factors for suicide in FEP [11,12]. Furthermore, low levels of cholesterol have been described in suicide behavior including among those individuals who have an increased tendency for impulsivity [13,14]. While, as a kind of psychological explanation, some scholars believe that young men in the early stages of their treatment are seeking to find meaning for frightening, intrusive experiences with origins which often precede psychosis, and these experiences invade personal identity, interactions and recovery [15], some suggests that personality characters, specifically, passive-dependent traits can be a predictor of first suicide attempts FEP [16]. On the other hand, no general agreement regarding higher prevalence of suicide in FEP is so far achievable. For example, while researchers like Nordentoft et al. [2], Bornheimer LA [8], Fedyszyn et al. [17] and Cohen et al. [18] have stated that FEP is a particularly high-risk period for suicide, with a risk as high as 10-60% during the first year of treatment, other scholars like Preti et al. [19], Pompili et al. [20], Crumlish et al. [21] and Addington et al. [22] have expected a lower risk or stated that suicide rates are difficult to measure in FEP patients, even in carefully defined samples. In the present study, suicides and suicide attempts among psychiatric in-patients, during the last five years, in Razi psychiatric hospital, as the largest national psychiatric hospital in Iran and region, has been evaluated to assess the general profile of suicidal behavior among native psychiatric inpatients, comparing first admission with recurrent admission patients and probing any relationship between serum cholesterol level and suicidal behavior.

## METHODS

Razi psychiatric hospital in south of capital city of Tehran, as one of the largest and oldest public psychiatric hospitals in the Middle East, which has been established formally in 1917 and with a capacity around 1375 active beds, had been selected as the field of study in the present retrospective assessment. Amongst its separate existent sections, five acute academic wards, which have been specified for admission of first episode adult psychiatric patients, and five acute non-academic wards, which have been specified for admission of recurrent episode adult psychiatric patients, with a collective capacity around two hundred active beds in each cluster (four hundreds beds, totally), had been selected for current study. Among the aforesaid academic divisions, two wards included female inpatients, with around eighty beds and the remaining three wards included male inpatients. All non-academic wards involved male inpatients. For valuation, all inpatients with suicidal behavior (successful suicide and attempted suicide, in total), during the last sixty months, had been included in the present investigation. Besides, clinical diagnosis was based on Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> edition (DSM-5) [23]. Also, assessment of serum lipids, including triglyceride (TG), cholesterol, low density lipoprotein (LDL) and high density lipoprotein (HDL), which was part of routine

laboratory checkups for all patients upon admission, whether for the first time or periodically, had been accomplished, for comparing the suicidal subjects with non-suicidal ones, incidentally.

**STATISTICAL ANALYSES**

While ‘t-test’ has been used for comparison of means as regards mean total level of serum lipids, difference of suicidal behavior between first admission and recurrent admission patients, had been analyzed by ‘comparison of proportions’. Statistical significance as well, had been defined as p value ≤ 0.05. MedCalc Statistical Software version 15.2 was used as statistical software tool for analysis.

**RESULTS**

As said by the results, among 19160 psychiatric patients hospitalized in razi psychiatric hospital, during a sixty months period (2013-2018), sixty three suicidal behaviors, including one successful suicide and sixty two suicide attempts, had been recorded by the security board of hospital (Table 1). Thirty three of patients were male and thirty of them were female, with no significant difference with regard to quantity (Table 2). The most frequent mental illness was bipolar I disorder (34.92%), which was significantly more prevalent in comparison with other mental disorders (p<0.04, p<0.02, p<0.007 and p<0.003 in comparison with

schizophrenia, depression, personality disorders and substance abuse, respectively). The other disorders included schizophrenia (19.04%), major depressive disorder (MDD) (17.46%), personality disorders (borderline & antisocial) (14.28%), substance abuse disorders, especially methamphetamine induced psychosis (MIP) (12.69%), and adjustment disorder (1.58%) (Table 3 and Figure 1). Also, no significant difference was evident between the first admission and recurrent admission inpatients, totally (p<0.31) and separately, particularly with respect to psychotic disorders (Table 3 and Figure 2). The annual incidences of suicidal behavior in both groups were comparable, and they were around 0.035% and 0.030%, in first admission and recurrent admission psychiatric inpatients, respectively (Table 1). While self-mutilation, self-poisoning and hanging were the preferred methods of suicide among 61.11%, 19.44% and 19.44% of cases, respectively, the first style was significantly more prevalent than the other ways (Z=1.96, P<0.059, CI: -0.0088,0.4532). Furthermore, no significant gender-based difference was evident with respect to the style of suicide in the present assessment (Figure 3). Besides, with respect to different components of serum lipids, no specific or significant pattern was evident, except that all hypolipidemic patients (n=7) were diagnosed as major depressive disorder, while 80% of hyperlipidemic patients (n=5) were diagnosed as bipolar I disorder (Table 4).

**Table 1.** Comparing suicidal behavior between first admission and recurrent admission psychiatric patients in Razi psychiatric hospital thru 2013-2018.

Total number of patients	First admissions	%	Annual Incidence	Recurrent admissions	%	Annual Incidence	Z	P	CI
19160	N=34	0.17	0.035	N=29	0.15	0.030	0.890	0.373	-0.09, 0.25

**Table 2.** Gender difference in suicidal behavior.

Suicidal cases	Male	Female	z	P	CI
N=63	33	30	0.53	0.59	-0.12, 0.22

**Table 3.** Frequency of psychiatric disorders among suicidal patients.

Psychiatric disorders	First admission	Recurrent admission	z	P	CI
Major depressive disorder	5	6	-0.42	0.66	-0.50,0.32
Bipolar I disorder	10	12	0.60	0.54	-0.20,0.38
Schizophrenia	8	4	1.63	0.10	-0.06,0.73
Substance abuse	5	3	1.00	0.31	-0.24,0.74
Personality disorders (borderline and antisocial)	5	4	0.47	0.63	-0.35,0.57
Adjustment disorder	1	0	1.41	0.15	-0.38,2.38
Total	34	29	0.89	0.37	-0.09,0.25




Figure 1. Diagrammatic representation of psychiatric disorders among suicidal subjects.





Figure 2. Graphic illustration of first and recurrent admissions' suicidal behavior.

Table 4. Lipid profile of suicidal patients.

Serum Lipids	Mean total levels in normal patients N=51	Mean total levels in Hyperlipidemic patients N=5 (20% MDD; 80% BID)	Mean total levels in Hypolipidemic patients N=7 (100% MDD)
Cholesterol	164.6 ± 18.66	235.33 ± 22.95	135.75 ± 20.75
Triglyceride	129.6 ± 36.67	191.66 ± 91.45	83.75 ± 27.22
LDL	96 ± 9.27	164.66 ± 23.57	86.5 ± 21.05
HDL	46.4 ± 16.71	51.66 ± 13.67	34.25 ± 15.12



**Figure 3.** Prevalence of different styles of suicide among inpatients.

## DISCUSSION

Always in psychiatry, when giving information about the diagnosis, course of illness, and treatment, the therapist should not ignore the risk of suicide [1]. Also, there are high proportion of young people with first-episode psychosis who attempted suicide before their first contact with mental health services [24]. It should constantly be considered that in the psychiatric hospital setting the inpatient at risk for suicide has previously exhibited suicidal behavior, suffers from schizophrenia, was admitted involuntarily, and lives alone [24]. While in terms of post-discharge suicides, the downsizing of psychiatric hospitals has been a success, there is still a substantial need for better recognition of suicidal risk among psychiatric patients [25]. According to a survey, there are 2 sharp peaks of risk for suicide around psychiatric hospitalization, one in the first week after admission and another in the first week after discharge [26]. The accessibility to one or more means of suicide is a recognized factor in psychiatric institutions [27]. But according to another study in FEP, the majority of attempts occurred when patients were treated as outpatients and were in regular contact with the service [17]. As suicide is a relatively rare event in psychotic disorders, general population-based prevention strategies may have more impact in this vulnerable group as well as the wider population [28,29]. While the immediate post-discharge period is a time of marked risk, rates of suicide remain high for many years after discharge and patients admitted because of suicidal ideas or behaviors and those in the first months after discharge should be a particular focus of concern [30]. Back to our discussion and according to the findings of the present study, The most common principal diagnoses among the suicide subjects were bipolar I disorder and schizophrenia, which was similar to the findings of Nordentoft et al. [2],

except that no gender difference regarding prevalence of these disorders among male and female patients was repeatable here. But, our findings were not in complete agreement with the conclusion of Thong et al. [31] who had found only schizophrenia and depression as the most common principal diagnoses among their suicide subjects and Roy et al. [24] who had declared schizophrenia as the major diagnosis among suicide victims. Meanwhile other psychiatric disorders, like personality disorders and substance abuse disorders, in addition to the aforementioned diagnoses, had been designated in the present study, as remarkable diagnoses among suicide subjects; though fewer than the aforementioned diagnoses. The higher prevalence of bipolar disorder in the present assessment is likewise comparable to the outcomes of a further study regarding suicidal behavior among Iranian inpatient youngsters [32]. In the same way, the higher incidence of self-mutilation, as the preferred method of suicide in the present evaluation, was parallel to the said study, except than it's significantly higher prevalence amongst female adolescents [33,34]. Also, in keeping with the results, while the annual incidences of suicidal behavior in both groups were comparable, they were lower than assessments of Björkenstam et al. [11] and Neuner et al. [35] and higher than approximations of Li et al. [36], which could be stemmed from cultural, instrumental, diagnostic and methodical differences. Also, in accordance with the outcomes of the present assessment, no significant difference was evident between the first admission and recurrent admission inpatients, totally and separately, particularly with respect to psychotic disorders. Such an outcome is clearly incongruous with the findings of Nordentoft et al. [2], Bornheimer LA [8], Fedyszyn et al. [17] and Cohen et al. [18] who have stated that first episode psychosis is a particularly high-risk period for suicide and first-episode psychotic disorder, in general, has seemed to be

a high-risk population for suicidal behavior during the first year of treatment. On the other hand, our findings are compatible with the stances of Preti et al. [19], Pompili et al. [20], Crumlish et al. [21] and Addington et al. [22], who have estimated a lower risk or indicated that suicide rates are difficult to measure in FEP patients, and there is relatively little specific information about the risk of suicide at illness onset or retrospectively concerning the untreated psychotic period. Above and beyond, with respect to relationship between serum lipids and suicidal behavior, outcome of the present assessment was not in harmony with the findings of Shrivastava et al. [13] and Ayesa-Arriola et al. [14], because while there was a couple of patients with higher or lower serum level of cholesterol, triglyceride, LDL and HD, no specific or significant pattern was evident in this regard; so, it seems that maybe such a difference was associated more to alteration of appetite, as a secondary phenomenon, rather than core variation of metabolism, as a primary etiologic issue. Anyhow, disregard to outcomes of the present study and its similarities or differences with comparable studies, elements of an inclusive prevention policy can be grouped under five items: securing the hospital environment, optimization of the care of the patients at suicidal risk, training of the medical teams in the detection of the risk and in the care of the suicidal subjects, involvement of the families in the care and implementation of post-event procedures following a completed suicide or an attempt [37]. Along with enhancement of insight, coping strategies should be boosted with a goal of minimizing depression and preventing suicidality [38]. Absence of post-discharge following program, deficiency of documented data regarding the suicidal behavior or its idea before admission, allocating wards to academic and non-academic, which could impact the quality of care, lack of female gender in the non-academic wards for making it more similar to the academic wards, which included an equal mixture of male and female inpatients, were among the weaknesses of the present assessment. In spite of remarkable findings of the current study, more methodical and comprehensive investigations in future, with taking into account the above shortages, can improve the quality and amendment of mental health services for proper response to patients' unavoidable problems.

## CONCLUSION

While in the present study the suicidal behavior was significantly more evident in bipolar disorder in comparison with other psychotic or no-psychotic disorders, no significant difference was evident between first admission and recurrent admission psychiatric inpatients. Moreover, no significant relationship between suicidal behavior and serum lipids was palpable.

## REFERENCES

1. American Psychiatric Association (APA) (2013) Diagnostic and statistical manual of mental disorders.

- 5<sup>th</sup> Edn. Washington (DC): American Psychiatric Association; pp: 801-806.
2. Nordentoft M, Madsen T, Fedyszyn I (2015) Suicidal behavior and mortality in first-episode psychosis. *J Nerv Ment Dis* 203: 387-392.
3. American Psychiatric Association (APA) (2013) Diagnostic and statistical manual of mental disorders. 5<sup>th</sup> Edn. Washington (DC): American Psychiatric Association; p: 104.
4. Kahn MW (2001) Emergency psychiatry: Tools of engagement: Avoiding pitfalls in collaborating with patients. *Psychiatr Serv* 52: 1571-1572.
5. Ballard ED, Stanley III, Horowitz LM, Cannon EA, Pao M, et al. (2013) Asking youth questions about suicide risk in the pediatric emergency department: Results from a qualitative analysis of patient opinions. *Clin Pediatr Emerg Med* 14: 20.
6. Hillard R, Zitek B (2004) Emergency psychiatry. New York, NY: McGraw-Hill Companies.
7. American Psychiatric Association (APA) (2013) Diagnostic and statistical manual of mental disorders. 5<sup>th</sup> Edn. Washington (DC): American Psychiatric Association; p: 131.
8. Bornheimer LA (2018) Suicidal ideation in first-episode psychosis (FEP): Examination of symptoms of depression and psychosis among individuals in an early phase of treatment. *Suicide Life Threat Behav* 23: 45-47.
9. González-Pinto A, Aldama A, González C, Mosquera F, Arrasate M, et al. (2007) Predictors of suicide in first-episode affective and non-affective psychotic inpatients: Five-year follow-up of patients from a catchment area in Vitoria, Spain. *J Clin Psychiatry* 68: 242-247.
10. Barrett EA, Sundet K, Faerden A, Agartz I, Bratlien U, et al. (2010) Suicidality in first episode psychosis is associated with insight and negative beliefs about psychosis. *Schizophr Res* 123: 257-262.
11. Björkenstam C, Björkenstam E, Hjern A, Bodén R, Reutfors J (2014) Suicide in first episode psychosis: A nationwide cohort study. *Schizophr Res* 157: 1-7.
12. Beckman K, Mittendorfer-Rutz E, Lichtenstein P, Larsson H, Almqvist C, et al. (2016) Mental illness and suicide after self-harm among young adults: Long-term follow-up of self-harm patients, admitted to hospital care, in a national cohort. *Psychol Med*; 46: 3397-3405.
13. Shrivastava A, Johnston M, Campbell R, De Sousa A, Shah N (2017) Serum cholesterol and Suicide in first episode psychosis: A preliminary study. *Indian J Psychiatry* 59: 478-482.

14. Ayesa-Arriola R, Canal Rivero M, Delgado-Alvarado M, Setién-Suero E, González-Gómez J, et al. (2018) Low-density lipoprotein cholesterol and suicidal behavior in a large sample of first-episode psychosis patients. *World J Biol Psychiatry* 11: 1-4.
15. Gajwani R, Larkin M, Jackson C (2018) "What is the point of life?": An interpretative phenomenological analysis of suicide in young men with first-episode psychosis. *Early Interv Psychiatry* 12: 1120-1127.
16. Canal-Rivero M, Barrigón ML, Perona-Garcelán S, Rodríguez-Testal JF, et al. (2016) One-year follow-up study of first suicide attempts in first episode psychosis: Personality traits and temporal pattern. *Compr Psychiatry* 71: 121-129.
17. Fedyszyn IE, Harris MG, Robinson J, Edwards J, Paxton SJ (2011) Characteristics of suicide attempts in young people undergoing treatment for first episode psychosis. *Aust N Z J Psychiatry* 45: 838-845.
18. Cohen S, Lavelle J, Rich CL, Bromet E. (1994) Rates and correlates of suicide attempts in first-admission psychotic patients. *Acta Psychiatr Scand* 90: 167-171.
19. Preti A, Meneghelli A, Pisano A, Cocchi A; (2009) Programma 2000 Team. Risk of suicide and suicidal ideation in psychosis: Results from an Italian multi-modal pilot program on early intervention in psychosis. *Schizophr Res*; 113: 145-150.
20. Pompili M, Serafini G, Innamorati M, Lester D, Shrivastava A, et al. (2011) Suicide risk in first episode psychosis: A selective review of the current literature. *Schizophr Res* 129: 1-11.
21. Crumlish N, Whitty P, Kamali M, Clarke M, Browne S, et al. (2005) Early insight predicts depression and attempted suicide after 4 years in first-episode schizophrenia and schizophreniform disorder. *Acta Psychiatr Scand* 112: 449-455.
22. Addington J, Williams J, Young J, Addington D. (2004) Suicidal behavior in early psychosis. *Acta Psychiatr Scand* 109: 116-20.
23. American Psychiatric Association (APA) (2013) *Diagnostic and Statistical Manual of Mental Disorders*. 5<sup>th</sup> Edn. Washington, DC: American Psychiatric Association; pp: 31-715.
24. Roy A, Draper R (1995) Suicide among psychiatric hospital in-patients. *Psychol Med* 25: 199-202.
25. Pirkola S, Sohlman B, Heilä H, Wahlbeck K (2007) Reductions in post-discharge suicide after deinstitutionalization and decentralization: A nationwide register study in Finland. *Psychiatr Serv* 58: 221-226.
26. Qin P, Nordentoft M (2008) Suicide risk in relation to psychiatric hospitalization: Evidence based on longitudinal registers. *Arch Gen Psychiatry* 62: 427-32.
27. Martelli C, Awad H, Hardy P (2010) In-patients suicide: epidemiology and prevention. *Encephale* 36: 83-91.
28. Dutta R, Murray RM, Allardyce J, Jones PB, Boydell J (2011) Early risk factors for suicide in an epidemiological first episode psychosis cohort. *Schizophr Res* 126: 11-19.
29. Dutta R, Murray RM, Hotopf M, Allardyce J, Jones PB, et al. (2010) Reassessing the long-term risk of suicide after a first episode of psychosis. *Arch Gen Psychiatry* 67: 1230-1237.
30. Chung DT, Ryan CJ, Hadzi-Pavlovic D, Singh SP, Stanton C, et al. (2017) Suicide rates after discharge from psychiatric facilities: A systematic review and meta-analysis. *JAMA Psychiatry* 74: 694-702.
31. Thong JY, Su AH, Chan YH, Chia BH (2008) Suicide in psychiatric patients: Case-control study in Singapore. *Aust N Z J Psychiatry* 42: 509-519.
32. Shoja Shafti S, Memarie A, Rezaie M, Hamidi M (2019) Clinical profile of suicide among adolescent psychiatric inpatients in Iran. *Biomed J Sci Tech Res* 18: 1-5.
33. Shoja Shafti S, Memarie A, Rezaie M, Hamidi M (2019) Suicides and suicide attempts among child and adolescent psychiatric inpatients in Iran. *Curr Trends Clin Med Sci* 2: 1-5.
34. Shoja Shafti S, Memarie A, Rezaie M, Hamidi M (2019) Suicidal conduct among youthful psychiatric inpatients: A pilot local survey. *J Forensic Gen Sci* 3: 216-220.
35. Neuner T, Schmid R, Wolfersdorf M, Spiessl H (2008) Predicting inpatient suicides and suicide attempts by using clinical routine data? *Gen Hosp Psychiatry* 30: 324-330.
36. Li J, Ran MS, Hao Y, Zhao Z, Guo Y (2008) Inpatient suicide in a Chinese psychiatric hospital. *Suicide Life Threat Behav* 38: 449-55.
37. Martelli C, Awad H, Hardy P (2010) In-patients suicide: Epidemiology and prevention. *Encephale* 36: 83-91.
38. Flanagan P, Compton MT (2012) A comparison of correlates of suicidal ideation prior to initial hospitalization for first-episode psychosis with prior research on correlates of suicide attempts prior to initial treatment seeking. *Early Interv Psychiatry* 6: 138-144.