

Treatment of Diabetic Ketoacidosis and Coma in Pregnant Women

Viktor Rosival*

*SYNLAB Department of laboratory Medicine, Dérer's Hospital, Limbová 5, SK-833 05 Bratislava, Slovakia.

Received January 07, 2019; Accepted January 30, 2019; Published January 31, 2019

Several papers have discussed the problems in the treatment of diabetic ketoacidosis and coma in pregnant women [1-3]. Now, it is the time to summarize the results of these discussions.

Substantial progress has been made possible with the Nobel-prize 1977 to Yallow for the development of new methods of biochemical analysis that it make possible to measure insulin concentration in human plasma. Since the general acceptance of the hyperosmolar hyperglycemic non-ketonic coma [4] has emerged a complicated situation: how is it possible to explain with insulin deficiency two diametric different phenomena, diabetic ketoacidotic coma and hyperosmolar hyperglycemic non-ketotic coma? Usually, a "very simple" explanation has been proposed; absolute deficiency of plasmatic insulin is the cause of diabetic ketoacidotic coma, relative deficiency of insulin is the cause of hyperosmolar hyperglycemic non-ketotic coma. However, the application of Yallow's methods to measure insulin concentration has shown that this is not correct. 1981 has been published the monograph "Diabetic coma: ketoacidotic and hyperosmolar" [5] and on page 67, Figure 6.3 has the names of 12 authors who have reported sufficient amounts of plasmatic insulin in patients with diabetic ketoacidosis. In contrast, absolute deficiency of plasmatic insulin has been reported in diabetic patients with hyperosmolar hyperglycemic nonketotic coma [6]. On the opinion that absolute deficiency of plasmatic insulin is the cause of diabetic ketoacidosis and coma has been casted doubts already 1973. Munro et al. [7] have reported patients with euglycemic diabetic ketoacidosis and, thus, the question aroused whether insulin deficiency is possible in euglycemia?

The solution of this problem has been made by several publications [8] reporting that the immediate cause of decreased level of consciousness up to coma is decreasing blood-pH (=increasing concentration of hydrogen ions H⁺). The glycolytic enzyme phosphofructokinase is pH-dependent, as its activity is decreasing with decreasing pH and, thus, is decreased also the utilisation of glucose in brain cells (regardless what is the concentration of glucose in the blood). The decrease of blood-pH is caused by increased concentration of up to 36 organic acids [9]; including also

reports on very low blood-pH without increase of acetoacetic and beta-hydroxybutyric acids [10].

In pregnant women, most dangerous is the situation, if the patient is admitted in coma. The treatment with sodium bicarbonate should begin as soon as possible to avoid irreversible damage of the brain cells caused by low blood-pH. If a diabetic pregnant woman is on insulin therapy, this should be carefully administered to avoid the development of hyperosmolar hyperglycemic non-ketotic coma. The blood-pH should be checked in regular intervals and if this is decreased, therapy with sodium bicarbonate infusion is indicated until normal values have been reached and a continuation of the therapy in the hospital should be considered.

SUMMARY

The methods to measure insulin concentration in the blood has made possible better understand the treatment of diabetic ketoacidotic coma and hyperosmolar hyperglycemic non-ketotic coma also in pregnant diabetic women. Successful treatment of ketoacidotic coma is infusions of sodium bicarbonate, insulin administration is important for prevention and treatment of hyperosmolar hyperglycemic non-ketotic coma.

REFERENCES

1. Dalfrà MG, Burlina S, Sartore G, Lapolla A (2016) Ketoacidosis in diabetic pregnancy. *J Matern Fetal Neonatal Med* 29: 2889-2895.
2. Khoo CM, Lee KO (2013) Endocrine emergencies in pregnancy. *Best Pract Res Clin Obstet Gynecol* 27: 885-891.
3. Jonsson M, Nordén-Lindeberg S, Ostlund I, Hanson U (2009) Metabolic acidosis at birth and suboptimal care--

Corresponding author: Dr. Viktor Rosival, SYNLAB Department of laboratory Medicine, Dérer's Hospital, Limbová 5, SK-833 05 Bratislava, Slovakia, Tel: 00421/33/5511831; E-mail: rosivalv@hotmail.com

Citation: Rosival V. (2019) Treatment of Diabetic Ketoacidosis and Coma in Pregnant Women. *Arch Obstet Gynecol Reprod Med*, 2(1): 38-39.

Copyright: ©2019 Rosival V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

illustration of the gap between knowledge and clinical practice. BJOG 116: 1453-1460.

4. Sament S, Schwartz MB (1957) Severe diabetic stupor without ketosis. S Afr Med J 31: 893-894.
5. Schade DS, Eaton RP, Alberti KGMM, Johnston DG (1981) Diabetic coma: Ketoacidotic and hyperosmolar. University of New Mexico Press, Albuquerque, USA.
6. Vinik A, Seftel H, Joffe BI (1970) Metabolic findings in hyperosmolar, non-ketotic diabetic stupor. Lancet 2: 797-799.
7. Munro JF, Campbell IW, McCuish AC, Duncan LJ (1973) Euglycaemic diabetic ketoacidosis. Br Med J 2: 578-580.
8. Nyenwe EA, Razavi LN, Kitabchi AE, Khan AN, Wan JY (2010) Acidosis: the prime determinant of depressed sensorium in diabetic ketoacidosis. Diabetes Care 33: 1837-1839.
9. Niwa T (1995) Mass spectrometry in diabetes mellitus. Clinica Chimica Acta 241: 190-220.
10. Vernon DD, Postellon DC (1986) Non-ketotic hyperosmolal diabetic coma in a child: Management with low-dose insulin infusion and intracranial pressure monitoring. Pediatrics 77: 770-772.