

Enhancing the Structure of Sandy Soil by Biological Molecules: An Innovative Approach to Water Conservation in Newly Reclaimed Desert-Agricultural Land

LM Mohamed, SH Mohamed, FN El Saeid, MZ Sayed, NM Ali, EN Abd El Rahman, M Abdel Wahab,
MM Ibrahim and AE Ghaly*

*Department of Agricultural Engineering, Faculty of Agriculture, Cairo University, Cairo, Egypt.

Received May 15, 2020; Accepted May 20, 2020; Published June 10, 2020

ABSTRACT

The main aim of this study was to enhance the water holding capacity of the reclaimed desert sandy soil using an environmentally friendly biological technique that would minimize the loss of irrigation water from plant root zone, by downward seepage. Two environmentally friendly bacteria (*Azotobacter chroococcum* and *Lactobacillus fermentum*) capable of producing insoluble polysaccharides as a soil pores plugging agent were selected. The polysaccharides production efficiencies of these bacteria were evaluated. The effectiveness of the polysaccharides in enhancing the water holding capacity of the soil was evaluated. The result showed that the sandy soil used in this study had a particle density of 2.58 g/cm³, a bulk density of 1.6 g/cm³ and a porosity of 37.98 %. The particle size varied from 0.150 to 2.000 mm with most of the soil particles having a diameter within the range of 0.425-0.850 mm indicating that the soil was free of silt and clay. This soil had a loose texture, high infiltration rate and low water holding capacity. *Azotobacter chroococcum* and *Lactobacillus fermentum* were capable of producing levan from sucrose. The levan yield was 0.248 g levan/g sucrose (62.78% of theoretical yield) and 0.371 g levan/g sucrose (93.92% of theoretical yield) for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. The cell yield was 0.074 g cell/g sucrose (47.69 % of theoretical yield) and 0.062 g cell/g sucrose (56.92 % of theoretical yield) for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. The polymer was effective as a plugging agent to plug the pores of the high permeability sandy soil. The results showed that increasing the concentration of bacteria had no significant effect on the amount of leachates collect from the soils treated with both bacteria. However, the leachates collected from the soils treated with *Azotobacter chroococcum* were much larger than those collected from the soils treated with *Lactobacillus fermentum*. Also, the leachates collected from the control (soils received no bacterial treatment) were much larger than soils treated with both bacteria. These microorganisms were suitable for production of levan from sucrose, fixing nitrogen in soil and producing growth hormones, thereby improving the water holding capacity of the soil and enhancing its nutrient content and stimulating plant growth.

Keywords: Sandy soil, Particle density, Bulk density, Porosity, Water holding capacity, Infiltration, Seepage, Bio-cementation, Bio-logical sealing

INTRODUCTION

Egypt is a transcontinental country situated mostly in north-eastern Africa, with the Sinai Peninsula in Western Asia. Egypt has a coastline at the Mediterranean Sea in north, and the Gulf of Suez and the Red Sea in east. The country lies in the dry arid region except for the northern part which enjoys a Mediterranean climate during winter (December-March) which is cool, windy and humid, with occasional rains. Summer in Egypt (June-September) is very dry with extremely hot temperatures into the 32-38°C, sometimes breaking into 48°C. Shoulder seasons (April-May and

Corresponding author: Emeritus Professor A. E. Ghaly, Department of Agricultural Engineering, Faculty of Agriculture, Cairo University, Cairo, Egypt, E-mail: abdel.ghaly@dal.ca

Citation: LM Mohamed, SH Mohamed, FN El Saeid, MZ Sayed, NM Ali et al. (2020) Enhancing the Structure of Sandy Soil by Biological Molecules: An Innovative Approach to Water Conservation in Newly Reclaimed Desert-Agricultural Land. Int. J. Biopro. Biotechnol. Advance, 6(1): 188-214.

Copyright: ©2020 LM Mohamed, SH Mohamed, FN El Saeid, MZ Sayed, NM Ali et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

October-November) are particularly pleasant months but with no rain. Egypt receives 20-200 mm of annual precipitation along the narrow Mediterranean Coast and nearly 0 mm in the central and the southern parts of the country [1, 2].

The total land area of Egypt is 1,000,450 km², of which 32,425 km²(3.24%) is the Delta which is made of silt deposits carried by the River Nile and in which is most agricultural land and live most of the population. The Western Desert of Egypt (681,000 km² or 68.07%) is an area of the Sahara which lies west of the River Nile up to the Libyan border, and from the Mediterranean Sea to the border with Sudan. The Eastern Desert (223,000 km² or 22.29%) extends east from the Nile to the Red Sea, and from the Mediterranean Sea to the border with Sudan. The area of Sinai is 61,000 km² (6.10%) and Northern Lakes cover an area of 3025 km² (0.30%) and is main source of aquaculture in Egypt. Generally, deserts are barren areas of landscape (**Figure 1**) where little or no precipitation occurs and, consequently, living conditions are hostile for plant and animal life. With 96% of Egypt's land is uninhabitable desert (never receives any rain) in both sides of the Nile, the population is concentrated around the narrow Nile Valley and Nile Delta, with smaller numbers along the Mediterranean and Red Sea coasts [3].

Egypt's worrying population boom poses very real dangers to the economic development of the country and is considered as a major challenge to the government. In 2000, the United Nations estimated that Egypt's population would hit 96 million in 2026. However, in 2017, there were some 104.5 million Egyptian, of which 9.5 million lived outside the country. With current population growth rate (2.6 million babies born in 2016), Egypt's population is expected to grow to 128 million by 2030 [4]. According to Egypt's Statistical Agency, the population growth rate must be one-third that of economic growth to prevent living standards from deteriorating [5]. Once the breadbasket of the Roman

Empire, Egypt began to import large quantities of wheat in the 1980s and is now importing 50% of its food [6].

The quest to bring desert land under cultivation has been a cornerstone of Egyptian Government Agricultural Policy since the 1952. The total area reclaimed reached 1.92 million *feddans* (*feddan*=0.42 hectare) in 1987. By 2002, the total reclaimable land was estimated at 2.8 million *feddans* [7]. However, the increase in agricultural land has not kept pace with the population increase in Egypt since 1950's. As a result, the country is facing unprecedented challenges as the agricultural lands are increasingly strained due to urban expansion and depletion of scarce water resources as the Nile faces upstream challenges with Ethiopia building Africa's largest dam [8]. Land reclamation in the Egyptian context means converting desert areas into agricultural land by extending water canals into the desert, enhancing soil fertility, and providing infrastructure for new village construction. If the unlimited desert sandy soil can be improved and provided with water, it can grow a lot of food for the growing population.

Therefore, in 2013, the Egyptian Government began an effort to reclaim approximately 1.5 million *feddan* of desert lands for agricultural use as a first stage of a major project aiming at the reclamation of 4 million *feddan*. Due to Nile water shortage, ground water will be used to irrigate 1,322,000 *feddan* (88.5%) and surface water will be used to irrigate 172,000 *feddan* (11.5%). The hope is that, with new wells, desalination plants and better infrastructure (new towns), farmers will be able to grow more wheat [2]. Example of this new development is the land reclamation project that began in 2015 in the Farâfra Depression (980 km²) in Western Desert. The white desert of Farâfra (**Figure 2**) has been converted into agricultural land capable of producing wheat, potato, radish and other produces (**Figure 3**). By adding the new farmland to Egypt's current 8.4 million *Fedden*, it is hoped to free the population from the narrow confines of the Nile Valley and have the capacity to meet food production needs [9].



Figure 1. Nature of Eastern and Western deserts of Egypt.

Egypt desert soils originated by mechanical disintegration and wind deposit. They are mostly loamy sand (of 95-97% sand and 3-5% clay). These soils are coarse, porous and well-drained and have a red to brown color. They contain salts and are high in potassium, phosphorus and nitrates. These soils have very low moisture, very low organic matter and a basic pH (7.5-8.0). Generally, the sandy soils of Sahara are one of the poorest types of soil for growing plants because of their very low nutrients and very poor water holding capacity [10-14]. In addition, surface irrigation in this dry climate can cause the water to evaporate very quickly leaving salts behind on the soil surface causing salinization. Furthermore, water uses (agricultural, industrial, municipal, transportation and electricity generation) and management in Egypt are very complex and there is a great deficit between the demand and supply [15]. Therefore, it is important to consider (a) water conservation through use of new irrigation technology, (b) new water sources such as desalination and municipal wastewater treatment and reuse and (c) improving the quality of the soil and its water holding capacity [4,7,16]. The farmer are already adopting new irrigation technology to conserve water [7] and the government has imparked on major desalination, and wastewater treatment and use projects as well as drilling wells for underground water [17]. However, building an adequate soil structure in the newly claimed land is still a major challenge.

In order to improve the soil properties, farmers are planting crops that fixes nitrogen such as alfalfa, but this process is unduly time-consuming for many farmers [4]. There are, however, several other techniques for land improvement including: (a) addition of biochar which significantly and permanently increase soil cation exchange capacity (the soil's ability to hold nutrients), creates habitats for beneficial microbes and increases water retention [18-22], (b) addition of organic matter such as well-rotted manure or finished compost which decomposes quickly (since microbial activity is so fast in hot climate) and improves the physical properties of the soil [23,27], (c) application of chemical grouting to stabilize soil structure and modify the pore geometry of the soil by chemical reactions or ionic exchange resulting in a reduced fluid movement and improved water holding capacity and reduced water and nutrient seepage [28-32] and (d) application of microorganisms to alter the soil structure in order to reduce porosity and enhance water and nutrient retention [33-48].

In order to improve the soil properties, farmers are planting crops that fixes nitrogen such as alfalfa, but this process is unduly time-consuming for many farmers [4]. There are, however, several other techniques for land improvement including: (a) addition of biochar which significantly and permanently increase soil cation exchange capacity (the soil's ability to hold nutrients), creates habitats for beneficial microbes and increases water retention [18-22], (b) addition

of organic matter such as well-rotted manure or finished compost which decomposes quickly (since microbial activity is so fast in hot climate) and improves the physical properties of the soil [23,27], (c) application of chemical grouting to stabilize soil structure and modify the pore geometry of the soil by chemical reactions or ionic exchange resulting in a reduced fluid movement and improved water holding capacity and reduced water and nutrient seepage [28-32] and (d) application of microorganisms to alter the soil structure in order to reduce porosity and enhance water and nutrient retention [33-48].

There are many microbiological activities that can be used to alter soil structure and improve the properties of soils which include bio-cementation (or bio-mineralization), gleization and bio-sealing. Bio-cementation is the process where microorganisms produce elemental compounds such as calcium carbonate as a basis for bio-grout that can improve the mechanical properties of the soil and decrease its porosity [33-37]. Gleization is a process in which breakdown of soil structure takes place by strong oxidizing or reducing gelatinous agents which are the products of microbial metabolisms [38-41]. Soil bio-sealing is a process in which microbially induced compounds are utilized to plug the soil pores and reduce soil porosity, leading to increased water holding capacity and reduced loss of water and nutrient through seepage [42-48].

Natural bio-seal (biological soil crust) can develop from the intimate association between soil particles and microorganisms that live within soil such as cyanobacteria, green algae, fungi, bacteria, lichens and bryophytes [44]. They are typical of arid and semi-arid regions but can occur in most ecosystems [45-46]. Some strains of bacteria produce water insoluble polysaccharides which appear to be promising selective plugging agents that can be used to create bio-seal in the sandy soil of Egypt [49,50]. Microbial polysaccharides which have potential in the sealing mechanisms include dextran, xanthan, curdlan, indicant, pullulan, heteroglycan and zenflox-polysaccharides. This study proposes to investigate the possibility of applying biological sealing into the sandy soil of reclaimed Egyptian deserts and evaluate its effectiveness in improving water retention.

OBJECTIVES

The main aim of this study was to enhance the water holding capacity of the reclaimed desert sandy soil using environmentally friendly biological technique that will minimize the loss of irrigation water by downward seepage out of the plant root zone. The specific objectives were: (a) to select environmentally friendly bacteria capable of producing insoluble polysaccharides as a plugging agent in order to minimize soil porosity, (b) evaluate the polysaccharides production efficiency of these bacteria and establish the optimum concentrations of the bacterial



Figure 2. White desert of Farâfra before reclamation.



Figure 3. Fields of crops in newly reclaimed agricultural land of Farafra.

cultures and (c) evaluate the effectiveness of the polysaccharides in enhancing the water holding capacity of the sandy soil.

MATERIALS AND METHODS

Selection of polysaccharide

The polysaccharide levan was selected for this study. Levan is a polymer made up of fructose (a monosaccharide sugar) connected in 2, 6 beta glycosidic linkages as shown in **Figure 4** [51, 52]. Levan can be in both branched and linear structures (**Figure 5**) of relatively low molecular weight [52]. In the branched version, levan forms a very small, sphere-like structure. This structure has basal chains of 9 units long which contain 2, 1 branching, allowing for the

methyl ethers to form and create a spherical shape. The ends tend to contain a glucosyl residue. The branched structure of levan tends to be more stable than the linear structure. However, the amount of branching and length of polymerization tends to vary among different species. The shortest levan is 6-kestose, essentially a chain of two fructose molecules and a terminal glucose molecule [52-53]. Levan contains a diverse set of properties (**Table 1**). The beta 2, 6 linkages of levan allow for it to be insoluble in water, oil and many organic solvents (methanol, ethanol, and isopropanol). The branching of levan also allow for it to have a large amount of tensile and cohesive strength, while the hydroxyl groups contribute to adhesion with other molecules [52-55].




Figure 4. Structural Formula of Levan [51,52].




Figure 5. Levan forms [52]. (a) Linear form with beta 2,6 glycosidic linkages. (b) Branched form with beta 2,1 glycosidic linkages.

Levan is diversity distributed in plants and microorganisms. It is usually found in the stems and leaf tissues of Levan is also produced as exopolysaccharides usually from sucrose (a disaccharide sugar containing glucose and fructose) based substrates by a variety of microorganisms

Agropyroncristatum, *Dactylisglomerata*, *Pea secunda*, *Triticumaestivum* and *Pachysadra terminalis* [56, 57], including bacteria, fungi and algae. However, there are some reports indicating that microbial levan can be produced from fructose, glucose and raffinose substrates [56, 58, 59]. The

main reaction in levan biosynthesis is the transfructosylation by the extracellular enzyme levansucrase. The enzyme forms the 2, 1 linkages in the linear basal chains of levan to allow

for branching points to occur. This production of levan is sensitive to temperature, oxygen concentration, pH and other factors [56, 60-62].

Table 1. The main properties of levan [51-70].

Property	Description
Molecular Formula	C ₁₈ H ₃₂ O ₁₆
Molecular Weight	504.4 g/mol
Solubility	Water and oil insoluble due to β-(2→6) linkage
Viscosity	Low viscosity (0.07-0.18 dL/g for molecular weight 16-24 million Da)
Particle Size	Nanoparticle in water = 224.3 nm Nanoparticles in ethanol = 251.8 nm
Stability	High stability to heat, acid and alkali media Melting point = 225 °C Glass transition temperature = 141 °C Boiling point = 900 °C
Enthalpy of Vaporization	150 kJ/mol
Ionic Bonding	Non-ionic
Assembling	Self-assembled in aqueous solution
Tensile Strength	Up to 10.3 MPa (1500 psi)
Polarity	Amphiphilic (poses water and fat loving properties)
Shape or Form	Amorphous (lacking clear structure)
Chemical Compatibility	Compatible with salts and surfactants
Biomedical Benefits	Non-toxic Antioxidant Anti-inflammatory Anticarcinogenic Antihyperlipidemic Antidiabetic Ameliorate stress Hyperglycaemic Prebiotic and immuno-nutrient Not hydrolyzed by human digestive enzymes Nanocarrier system of peptides, proteins and drugs

Selection of microorganism

Selection of microorganisms used in this study was based on the criteria shown in **Table 2**. As the land will be used for agriculture production, contaminated soil with pathogens could spread diseases to crops and vegetables or to healthy animals and human. Thus, the selected microorganisms must be non- pathogenic. Microbial cells smaller than the average pore sizes of the soil are desirable. Insoluble polysaccharide is required to plug the soil pores and form stable sealing. *Arthrobacter* and *Bacillus* are the most common bacterial genera found in soils and any microbes introduced into to the soil for the purpose of clogging the soil pores (bio-sealing) must compete with these indigenous bacteria for substrate [42,49,50].

Table 3 shows some of the levan producing bacteria. The bacterial species *Azotobacter chroococcum* and

Lactobacillus fermentum were selected for this study. *Azotobacter chroococcum* are capable of producing levan and have a full range of enzymes needed to perform nitrogen fixation (ferredoxin, hydrogenase, and an important enzyme nitrogenase). Owing to their ability to fix molecular nitrogen and produce growth hormones, and therefore increase the soil fertility and stimulate plant growth, *Azotobacter* species are widely used in agriculture as a source of nitrogen biofertilizer [71]. *Lactobacillus fermentum* bacteria are a levan producing bacteria. The use of these two microorganisms would be suitable for production of levan from sucrose, fixing nitrogen in soil and producing growth hormones, thereby improving the water holding capacity of the soil and enhancing its nutrient content and stimulating plant growth. The scientific classifications of *Azotobacter chroococcum* and *Lactobacillus fermentum* are shown in **Table 3** and their biological and biochemical characteristics are shown in **Table 4**.

Table 2. Selection criteria of microorganisms used in the study.

Criteria	Descriptions
Pathogenicity	As the treated soil will be used for agricultural production, contaminated crops and vegetables can spread the diseases to healthy animals and human. Thus, the selected microorganism must be non-pathogenic
Size	Cells size must be smaller than the average pore diameter of the soil (the average pore diameter of agricultural soils is 7 - 15 μm and for sandy soil that drains freely by gravity is $>150 \mu\text{m}$)
Type of Polymer Produced	Non-soluble polysaccharide required to form stable sealing
Competition with Soil Microorganisms	<i>Arthrobacter</i> and <i>Bacillus</i> species are the most dominant soil microorganisms and the selected microorganisms must be able to compete with these as well as other microorganisms in the soil such as yeast and fungi.

Soil collection and preparation

The soil was collected from the Teaching and Experimental Farm of the Faculty of Agriculture, Cairo University. About 100 kg of soil were collected in plastic bags and transported to the Bioengineering Laboratory. The visible organic matter was removed from the soil and soil clumps were crushed.

A soil sample of 500g was used to determine the particle size distribution using a mechanical sieving apparatus (Vibratory Sieve Shaker, Series AS200, Retsch GMBH, Haan, Germany). The pan was first placed onto the sieving apparatus. The sieves with the smallest mesh were stacked on the top of the pan and successively larger meshes were placed above. The sample was placed into top sieve and the lid was placed on top of the stack. The shaker was turned on for 30 min. The soil collected from each sieve was weighed and the percentage of each soil fraction from the original soil weight was calculated.

Soil samples of 50 g each were used to measure the soil particle density, bulk density and porosity. A soil sample of 50 g was placed in a 100 ml graduated cylinder and the actual volume of the soil sample was determined. Another soil sample of 50 g was placed into a graduated cylinder containing 100 ml of water. The volume of water that resulted from the addition of soil is considered the volume of the soil particles. The particle density is defined as the weight of the soil particles divided by their volume. The bulk density is defined as actual weight of the soil divided by its apparent volume. The soil porosity is defined as these were calculated as follows:

$$\rho_p = W/V_p \quad (1)$$

$$\rho_b = W/V_b \quad (2)$$

$$P = (\rho_b - \rho_p) / \rho_b \quad (3)$$

where:

P = Porosity (%)

V_b = Volume of the soil (cm^3)

V_p = Volume of the particles (cm^3)

W = Weight of the soil (g)

ρ_b = Soil bulk density (g/cm^3)

V_p = Particles density (g/cm^3)

The rest of the soil was placed in 10 plastic bags, each containing 1 kg of soil. The bags placed in an autoclave (Tabletop Autoclave, Tuttnauer 2340M, Alpha Scientific, Vancouver, British Columbia, Canada) for sterilization at a temperature of 121°C and a pressure of 103 KPa for 20 min. This process was carried out to kill any soil microorganisms. The sterilized soil was used later to test the effectiveness of bio-cementation and bio-sealing (clogging of the soil pores).

The rest of the soil was placed in 10 plastic bags, each containing 1 kg of soil. The bags placed in an autoclave (Tabletop Autoclave, Tuttnauer 2340M, Alpha Scientific, Vancouver, British Columbia, Canada) for sterilization at a temperature of 121°C and a pressure of 103 KPa for 20 min. This process was carried out to kill any soil microorganisms. The sterilized soil was used later to test the effectiveness of bio-cementation and bio-sealing (clogging of the soil pores).

Preparation of the growth medium and microbial cultures

Samples of *Azotobacter chroococcum* and *Lactobacillus fermentum* (Figure 6) were obtained from the Department of Microbiology, Faculty of Agriculture, Ein Shams University and the Department of Microbiology, Faculty of Agriculture, Cairo University, respectively. Liquid growth medium was prepared using Bacto® Nutrient Broth, which was obtained from Difco Laboratories, Detroit, Michigan, USA.

The Nutrient broth is composed of a simple peptone and a beef extract. The peptone contributes organic nitrogen in the form of amino acids and long-chained fatty acids while the

Table 3. Some levan producing bacteria.

Microorganism	Reference
<i>Acetobacter aceti</i> ^{SG-}	Loewenberg and Reese [72], Moonmangmee et al. [73], Tomulescu et al. [74]
<i>Acetobacter diazotrophicus</i> ^{SNG-}	Tomulescu et al. [74], Hernandez et al. [75], Arrieta et al. [76], Tambara et al. [77], Batista et al. [78]
<i>Acetobacter Pasteurianus</i> ^{SNG-}	Loewenberg and Reese [72], Tomulescu et al. [74], Perumpuli et al. [79], Minakami et al. [80], Tayama et al. [81]
<i>Acetobacter xylinum</i> ^{SG-}	Srikanth et al. [56], Tomulescu et al. [74], Tayama et al. [81], Jasson et al. [82], Wong et al. [83]
<i>Acinetobacter nectaris</i> ^{PG-}	Gonzalez-Garcinuno [84], Bansal et al. [85], Taberero et al. [86]
<i>Actinomyces viscosus</i> ^{HAG+}	Pabst [87], Warner and Miller [88], Miller and Somers [89], Igarashi et al. [90]
<i>Achromobacter xylosoxidans</i> ^{HG-}	Han [91], Yamasato et al. [92]
<i>Aerobacteraerogenes</i> ^{HG-}	Han [91], Srinivasan and Quastel [93], Wilkinson et al. [94]
<i>Aerobacter levanicum</i> ^{SG-}	Evans and Hibbert [95], Takeshita et al. [96], Feingold and Gehatia [97]
<i>Arthrobacter Ureafaciens</i> ^{SG+}	Tomulescu et al. [74], Han [91], Song et al. [98], Tanaka et al. [99], Tanaka et al. [100]
<i>Azotobacterchroococum</i> ^{SG}	Tomulescu et al. [74], Hestrin and Goldblum [101], De La Vega et al. [102], Han [103]
<i>Bacillus arrophilus</i> ^{HG+}	Tomulescu et al. [74], Bansal et al. [85], Abou-Taleb et al. [104]
<i>Bacillus atrophaeus</i> ^{SG+}	Tomulescu et al. [74], Bansal et al. [85], Hestrin and Goldblum [101]
<i>Bacillus amyloliquefaciens</i> ^{SG+}	Tomulescu et al. [74], Han [91], Tian et al. [105]
<i>Bacillus lentus</i> ^{SG-}	Tomulescu et al. [74], Bansal et al. [85], Abou-Taleb et al. [104]
<i>Bacillus licheniformis</i> ^{SG+*}	Ghaly [42], Ghaly et al [49], Ramsay et al. [54], Tomulescu et al. [74], Bansal et al. [85], Xavier et al. [106], Kekez et al. [107], Mamay [108], Larpin et al. [109], van Dyke et al. [110]
<i>Bacillus megaterium</i> ^{SG+}	Tomulescu et al. [74], Evans and Hibbert [95], Strube et al. [111]
<i>Bacillus mesentericus</i> ^{SG+}	Tomulescu et al. [74], Han [91], Tanaka et al. [112]
<i>Bacillus methylotrophicus</i> ^{G+}	Tomulescu et al. [74], Zhang et al. [113], Li et al. [114], Jadan et al. [115]
<i>Bacillus polymyxa</i> ^{SG+}	Han and Clark [51], Tomulescu et al. [74], Bansal et al. [85], Han and Watson [116], Han [117], Liu et al. [118]
<i>Bacillus subtilis</i> ^{SG+}	Shih et al. [60], Bansal et al. [85], Jensen et al. [119], Ing-Lung et al. [120], Abdel-Fattah et al. [121], Benigaret al. [122], Ahmed [123], Esawy et al. [124], Vaidya and Prasad [125], Goncalves et al. [126], Abdul Razack [127], Abdel-Fattah et al. [128]
<i>Bradyrhizobiumdenitrificans</i> ^{SNG-}	Bansal et al. [85], Sutherland [129]
<i>Bradyrhizobiumelkanii</i> ^{SNG-}	Bansal et al. [85], Sutherland [129], Sucawara et al. [130]

<i>Bradyrhizobium embiense</i> ^{SNG-}	Bansal et al. [85], Sutherland [129]
<i>Bradyrhizobium japonicum</i> ^{SNG-}	Bansal et al. [85], Sutherland [129], Sudtachat et al. [1317], Dake [132]
<i>Bradyrhizobium oligotrophicum</i> ^{SNG-}	Bansal et al. [85], Sutherland [129], Resenberg et al. [133]
<i>Bradyrhizobium yuanmingense</i> ^{SNG-}	Bansal et al. [85], Sutherland [129]
<i>Brenneriagoodwinii</i> ^{SPG-}	Tomulescu et al. [74], Liu et al. [134], Xu et al. [135]
<i>Clostridium acetobutylicum</i> ^{SG+}	Gao et al. [136], Dahech et al. [137]
<i>Corynebacterium laevaniformans</i> ^{SG+}	Han [91], Dias and Bhat [138]
<i>Corynebacterium beticola</i> ^{SHG+}	Tomulescu et al. [74], Han [87], Chen et al [139]
<i>Erwinia amylovora</i> ^{SPG+}	Tomulescu et al. [74], Wuerges et al. [140], Gross et al. [141]
<i>Erwinia herbicola</i> ^{SPG-}	Tomulescu et al. [74], Benigar et al. [122], Keith et al. [142], Keith et al. [143]
<i>Geobacillus stearothermophilus</i> ^{SG+}	Inthanovong et al. [144], Li et al. [145]
<i>Gluconobacteraquatilis</i> ^{SPG-}	Tomulescu et al. [74], Ua-AraK et al. [146], De Muynck et al. [147]
<i>Gluconobacter cerinus</i> ^{SPG-}	Tomulescu et al. [74], De Muynck et al. [147], Jakob et al. [148]
<i>Gluconobacter oxydans</i> ^{SG-}	Velazquez-Hernandez et al. [149], Park et al. [150]
<i>Gluconoacetobacter diazotrophicus</i> ^{SNG-}	Han [91], Serrato et al. [151], Banguela et al. [152]
<i>Glucoacetobacter xylinus</i> ^{SG-}	Jakob et al. [148], Kommann et al. [153],
<i>Halomonas myrnesensis</i> ^{SG-}	Sarilmiser et al. [59], Kazak et al. [154], Poli et al. [155]
<i>Kozakiabaliensis</i> ^{SWG-}	Ua-AraK et al. [146], Brandt et al. [156]
<i>Lactobacillus fermentum</i> ^{SWG+}	Dutta et al. [157], Badel et al. [158], Heinemann et al. [159], Galle and Avendt [160]
<i>Lactobacillus gasseri</i> ^{SWG+}	Anwar et al. [161], Diez-Municio et al. [162]
<i>Lactobacillus reuteri</i> ^{SWG+}	Tomulescu et al. [74], Sims et al. [163], van Hijum et al. [164], Kaditzky and Vogel [165], Ni et al. [166]
<i>Lactobacillus sanfranciscensis</i> ^{SWG+}	Bansal et al. [85], Tieking et al. [167],
<i>Leuconostoc citreum</i> ^{SWG+}	Tomulescu et al. [74], Han et al. [168], Ortiz-Soto et al. [169], Bounaix et al. [170]
<i>Leuconostoc mesenteroides</i> ^{SWG+}	Han [81], Xu et al. [171]
<i>Mesorhizobium alhagi</i> ^{SNG-}	Bansal et al. [85], Liu et al. [172]
<i>Mesorhizobium amorphae</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Kimbrel [174]
<i>Mesorhizobium australicum</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Reeve et al. [175]
<i>Mesorhizobium ciceri</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Das et al. [176]
<i>Mesorhizobium huakuii</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Chen et al. [177]
<i>Mesorhizobium japonicum</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173]
<i>Mesorhizobium loti</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Kawaharada et al. [178], Kelly et al. [179]

<i>Mesorhizobiummediterraneum</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Ray [180]
<i>Mesorhizobiumplurifarium</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173]
<i>Micbacteriumlaevaniformans</i> ^{SG+}	Bansal et al. [85], Han [91], Bae et al. [181]
<i>Odontomycesviscosus</i> ^{SWHAG+}	Han [91], Krichevsky et al. [182]
<i>Paenibacillusbovis</i> ^{SG+}	Gozalez-Garcinuno [84], Xu et al. [183], Hang et al. [184].
<i>Pediococcusacidilactici</i> ^{SHG+}	Youssef et al [62], Petrov and Petrova. [185]
<i>Phytomonaspruni</i> ^{SPG+}	Han [91], Haworth and Stacey [186], Lyne et al. [187]
<i>Pseudomonas aureofaciens</i> ^{SPG-}	Tomulescu et al. [74], Fuchs [188], Alamäe et al. [189]
<i>Pseudomonas brassicacearum</i> ^{SPG+}	Tomulescu et al. [74], Alamäe et al. [190], Al Qysi [191]
<i>Pseudomonas chlororaphis</i> ^{SG-}	Tomulescu et al. [74], Fuchs [188], Alamäe et al. [189]
<i>Pseudomonas fluorescens</i> ^{SG-}	Jathore et al. [58], Tomulescu et al. [74], Bansal et al [85], Fuchs [188], Alamäe et al. [189]
<i>Pseudomonas syringae</i> ^{SPG-}	Tomulescu et al. [74], Alamäe et al. [189], Kasapis et al. [191], Laue et al. [192]
<i>Rahnellaaquatilis</i> ^{SWHG-}	Yoo et al. [193], Kim et al. [194], Kim et al. [195]
<i>Rhizobium leguminosarum</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173]. Karunaratne [196], Tikhonovich et al. [197]
<i>Rhisobiummeliloti</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Tikhonovich et al. [197]
<i>Rhisobiumraioabacter</i> ^{SNG-}	Bansal et al. [85], Priest and Goodfellow [173], Tikhonovich et al. [197]
<i>Rothiadentocariosa</i> ^{SWHG+}	Tomulescu et al. [74], Han [91], Leshner and Gerencser [198], Willner et al. [199], Hill [200]
<i>Saccharomyces cerevisiae</i> ^{SG-}	Tomulescu et al. [74], Bansal et al. [85], Franken et al. [201], Elorza et al. [202]
<i>Streptococcus mutans</i> ^{SHG+}	Han [91], Yoo et al. [203], Ebisu et al. [203]
<i>Streptococcus salivarius</i> ^{WHG+}	Tomulescu et al. [74], Fuchs [188], Yoo et al. [193], Ebisu et al. [204], Newbrun et al. [204]
<i>Xanthomonas axonopodis</i> ^{SPG-}	Tomulescu et al. [74], Han [91], Yoo et al. [193], Moosavi-Nasab et al. [205]
<i>Zymomonasmobilis</i> ^{SG-}	Silbir et al. [57], Tomulescu et al. [74], Bansal et al. [88], Benigar et al. [122], Abdul Razack [127], Yoo et al. [193], Vigants et al. [206], Calazans et al. [207], Bekers et al. [208], Melo et al. [209], Ananthalakshmy and Gunasekaran [210], Shaheen et al. [211], Santos et al [212], De Oliveira et al. [213]

A = Animal Pathogen

G+ = Gram Positive

G- = Gram Negative

H = Human Pathogen

N = Nitrogen Fixing

P = Plant Pathogen

S = Soil microorganism

W = Water microorganism

beef extract provides vitamins, carbohydrates, salts and other organic nitrogen compounds. An amount of 6.5 g of the nutrient broth was added to two Erlenmeyer flasks, each containing 500 ml distilled deionized water. Each flask was

capped and thoroughly mixed. The flasks were placed in an autoclave (Tabletop Autoclave, Tuttnauer 2340M, Alpha Scientific, Vancouver, British Columbia, Canada) at a temperature of 121°C and a pressure of 103 KPa for 20 min

to sterilize the media. The flasks were left to cool down. One flask was inoculated aseptically with *Azotobacter chroococum* while the other flask was inoculated aseptically with *Lactobacillus fermentum*. The two cultures were grown on a controlled environment laboratory shaker (MaxQ™ 4000 Benchtop Orbital Shaker, Thermo Fisher Scientific, Montreal, Quebec, Canada) at room temperature (21°C) for 24 h.

Table 4. Scientific classification of *Azotobacter chroococum* and *Lactobacillus fermentum* [214-216].

Taxonomy	<i>Azotobacter chroococum</i>	<i>Lactobacillus fermentum</i>
Kingdom	Bacteria	Bacteria
Phylum	Proteobacteria	Firmicutes
Class	Gammaproteobacteria	Baccilli
Order	Pseudomonadales	Lactobacillales
Family	Pseudomonadaceae	Lactobacillaceae
Genus	Azotobacter	Lactobacillus
Species	Azotobacterchroococum	Lactobacillus fermentum

The cell number was determined according to the procedure described by Ghaly and Mahmoud [217].

Culture propagation and polymer production in laboratory

The two cultures were then grown on liquid growth medium containing sucrose. The liquid growth medium consisted of 50.0 g sucrose, 2.5 g tryptone, 2.5 g K₂HPO₄ and 5.9 g yeast extract per liter of distilled deionized water. The media were transferred to several 1 L Erlenmeyer flasks and sterilized in an autoclave (Tabletop Autoclave, Tuttnauer 2340M, Alpha Scientific, Vancouver, British Columbia, Canada) at a temperature of 121°C and a pressure of 103 KPa for 20 min. Each microbe was transferred aseptically from the nutrient broth to ten 750 mL Erlenmeyer flasks, each having 500 ml of sterilized liquid media. Each flask was inoculated with 10% (v/v) of the homogeneous mixture of the nutrient broth culture. The cultures were grown in a controlled environment laboratory shaker ((MaxQ™ 4000 Benchtop Orbital Shaker, Thermo Fisher Scientific, Montreal, Quebec, Canada) at room temperature (21°C) for 5 days. Samples were drawn from the flasks for biomass, sucrose and polysaccharide determination. Sampling was done every 4 h during the first 24 h every 6 h during the period of 24-72 h and then every 12 h until the end of the 5 days. The cell biomass was determined according to the procedure described by Ghaly and Mahmoud [218]. The polysaccharide concentration analysis was determined according to the procedure described by Ramsay [219]. The

sucrose concentration was determined according to the procedure described by Borji et al. [220].

Polymer production in soil (bio-cementing and bio-sealing)

The setup for testing sandy soil bio-cementation and bio-sealing is shown in **Figure 7**. It consisted of 5 infiltration soil columns, each was constructed a PVC cylinder of 7.5 cm diameter and 40 cm height, a plastic filtration funnel of 7.5 cm diameter and a 1 L flask. The funnel was placed on the top of the flask and a filter bad was placed inside the funnel. This filter bad has a pore size smaller the that of the smallest sand particles (does not allow the soil particles to pass through). The cylinder was connected to the funnel and sealed together. One kg of the sterilized soil was placed in the cylinder and packed to achieve field density. This was done to simulate the soil root zone.

The application of microbial culture and water was carried out as shown in **Table 6**. 400 ml of microbial culture of *Azotobacter chroococum* were added on day 1 to each column. On Day 3, 400 ml of the diluted microbial culture (each soil column received different concentration of the microbial) were added to the columns.

On day 5, the moisture content and pH were measured. The moisture content was measured using a portable soil moisture measurement meter (TOR 150 Soil Moisture Meter, Edaphic Scientific, Moorabbin, Victoria, Australia). The pH was measured using a portable pH meter (Hanna H199121 Digital pH Meter, ITM Instruments Inc, Sainte Anne de Bellevue, Quebec, Canada). Then, 400 ml of water were added to each column and the leachate collected in the flasks were measured after 12, 24 and 48 h from addition of water. Finally, the water holding capacity was determined.

After completing the experiment with *Azotobacter chroococum*, the component of each column were dismantled and washed thoroughly with water and disinfected with alcohol. They were the sterilized in an autoclave (Tabletop Autoclave, Tuttnauer 2340M, Alpha Scientific, Vancouver, British Columbia, Canada) at a temperature of 121°C and a pressure of 103 KPa for 20 min. The 5 columns were reassembled again. The same experimental procedure was followed with *Lactobacillus fermentum*.

RESULTS AND DISCUSSION

Soil characteristics

The result showed that the sandy soil used in this study had a particle density of 2.58 g/cm³, a bulk density of 1.6 g/cm³ and a porosity of 37.98 %.

Table 5. The biological and biochemical characteristics of *Azotobacter chroococum* and *Lactobacillus fermentum* [74,101-103,157-160].

Parameter	<i>Azotobacter chroococum</i>	<i>Lactobacillus fermentum</i>
Habitats	Neutral to alkaline soils, aquatic environments and on some plants.	Fermented milk products, sourdough, fermenting plant materials, faeces and ewage
Motility	Free-living microbes	Non motile
Staining	Gram negative	Gram positive
Bacteria Shape	Oval or spherical	Rod-shaped
Bacteria Size	0.6 - 0.9 μm by 1.5 - 3.0 μm	0.5 - 0.8 μm by 2 - 9 μm
Spore Shape	A closed sac containing a cluster of cells	Non spore forming
Oxygen	Obligate aerobes	Facultative anaerobes
Temperature	Optimum growth at 20 - 33oC	Optimum growth at 20 - 30oC
pH	4.8 - 8.5 with optimum at 7.0-7.5	Strong pH tolerance (pH 3)
Tolerance to Salt	Good growth up to 6% NaCl	Ferment sugars up to 6-8% NaCl
Growth on Agar	Large spreading flat, slimy colonies with a diameter of 5-10 mm and a dark brown to green color	Colonies on agar media are usually 2-5 micrometers, convex, entire, opaque, and without pigment
Substrate	Wide variety of carbohydrates and organic metallic salts with mannitol as a source of energy	Ferment ribose, galactose, D-glucose, D-fructose, D-mannose, maltose, lactose, melibiose, saccharose, D-raffinose, D-tagatose and gluconate.
Applications	Fixing nitrogen (bio-fertilizer) Production of growth hormones Production of polysaccharides	Potential probiotic Production of lactic acid Production of polysaccharides

The fraction of the soil collected from each sieve as a percentage of the original soil weight is shown in **Table 7**. The particle size varied from 0.150 to 2.000 mm. Most of the soil particles had a diameter in the range of 0.425-0.850 mm (**Figure 8**). All sand particles have a diameter within the range 0.05 mm and 2.00, all silt particles have a diameter within the range 0.002 mm and 0.05 mm while all clay particles are less than 0.002 mm in diameter as shown in **Figure 9** [221].

The results indicated that the soil used in this study is typical sandy soil with a rough texture and free of silt and clay. This soil has a loose texture resulting in wind erosion, low organic matter, low nutrient content, high infiltration rate, low water holding capacity, high temperature resulting in faster plant growth, high aeration rate resulting in faster decomposition of organic matter [222]. For a sustainable agriculture, it is important to consider applying biotechnological techniques for building an adequate soil structure in these types of soils as well as water conservation by adopting new irrigation technology.

**Figure 6.** Samples of *Azotobacter chroococum* and *Lactobacillus fermentum*.



Figure 7. Bio-cementation and bio-sealing testing

Bacterial growth and levan production in bioreactor

The bacteria *Azotobacter chroococcum* and *Lactobacillus fermentum* were first grown on a sucrose in shake flasks to produce levan. The plate count test performed on the media obtained from the shake flasks containing nutrient broth revealed that there was a count of approximately 7.29×10^8 microbial cells/mL for *Azotobacter chroococcum* and 8.23×10^8 microbial cells/mL for *Lactobacillus fermentum*. The result of batch culture propagation of *Azotobacter chroococcum* and *Lactobacillus fermentum* in the shake flasks are presented in Table 8. The maximum biomass concentration was 3.6 g/L and 3.0 g/L after 22 h and 26 h for *Azotobacter chroococcum* and

Table 6. Microbial culture and water applications.

Time	Application
Day 1	Units 1-5 400 ml microbial culture
Day 3	Unit 1 (control): 400 ml water+0 ml microbial culture
	Unit 2 (treatment): 300 ml water+100 ml microbial culture
	Unit 3 (treatment): 200 ml water+200 ml microbial culture
	Unit 4 (treatment): 100 ml water+300 ml microbial culture
	Unit 5 (treatment): 0 ml water+400 ml microbial culture
Day 5 (Low moisture content)	Units 1-5 400 ml water

Lactobacillus fermentum, respectively. The concentration of sucrose decreased reaching 1.6 g/L and 1.5 g/L after 56 h and 61 for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. With the depletion of sucrose, the bacterial cell mass decreased reaching 0.16 g/L and 0.13 g/L after 68 h and 72 h for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. The bacteria produced the enzyme levansucrase which converts the soluble sucrose into the polysaccharide β -D fructoside (levan) and glucose. The production of levan reached a maximum of 14 g/L and 17 g/L for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively.

During the fermentation process, the bacteria utilize sucrose for production of levan and for cell maintenance and growth. The following equations describe product formation, respiration and energy production and growth and reproduction.

(a) Respiration and energy production




Table 7. Soil Particle size distribution.

Sieve Number	Diameter (mm)	Retained (Kg)	Soil (%)	Passing (%)
10	2	48	9.6	90.4
20	0.85	105	21	69.4
40	0.425	199.5	39.6	29.8
60	0.25	137	27.4	2.4
100	0.15	12	2.4	0
Pan	0.075			

Soil Sample = 500 g






Figure 8. Sand particle sizes and percentage of particles passing through sieves.


(b) Growth and reproduction



(c) Product formation



Equation (4), (5) and (6) can be combined to yield the following equation:






Figure 9. Sand, silt and clay particle sizes [217].

From equation (7), it appears that the theoretical levan yield is 0.395 g levan/g sucrose and the theoretical cell yield is 0.130 g cells/g sucrose. In this study, the levan yield was 0.248 g levan/g sucrose (62.78% of theoretical yield) and 0.371 g levan/g sucrose (93.92% of theoretical yield) for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. The cell yield was 0.074 g cell/g sucrose (47.69 % of theoretical yield) and 0.062 g cell/g sucrose (56.92 % of theoretical yield). The results showed that *Lactobacillus fermentum* was more efficient in converting sucrose to levan than *Azotobacter chroococcum*. However, *Azotobacter chroococcum* produced more bacterial biomass (g cells/g sucrose) than *Lactobacillus fermentum*. This may be due to the fact that *Azotobacter chroococcum* is a nitrogen fixing microorganism, a process that requires organic matter. Thus, some of the sucrose may have been utilized in nitrogen fixation.

The results showed that it is feasible to use growing cultures of *Azotobacter chroococcum* and *Lactobacillus fermentum*. From the biological and biochemical characteristics of the *Azotobacter chroococcum* and *Lactobacillus fermentum*, it appears that the organisms can produce levan from sucrose under most field and soil conditions and they should be able to compete with most common soil microbial species.

The polysaccharide (levan) produced in this study was non-viscous and water insoluble. The viscosity of the culture broth was the same as that of water. The polymer was a non-transparent suspension and was found to deflect visible light. The polymer can be used as a plugging agent to plug the pores of high permeability soils. Microbial levan contains up to 3 million residues compared to plant levan which contains about 100 residues [51]. The polysaccharide levan $(C_6H_{10}O_5)_n$ consists of fructose monomers linked mainly by $\beta(2\rightarrow6)$ linkages [91].

Bio-cementation and bi-sealing

The moisture content and pH measurements taken on day 5 before the application of 400 ml water to each of the soil bio-cementation and bio-clogging columns are presented in Table 9. The results indicated that the moisture content of the soils receiving the bacterial culture of *Azotobacter chroococcum* (22.3%) was lower than that of the soils receiving the bacterial culture of *Lactobacillus fermentum* (25.3%). The soils designated as control (received no bacterial treatment) had lower moisture content than the soils treated with both bacterial cultures. The moisture content of the soils receiving the bacterial culture of *Azotobacter chroococcum* was higher than that of the control by 13.78% while the moisture content of the soil receiving the bacterial culture *Lactobacillus fermentum* was higher than that of the control by 29.59%. However, increasing the concentration of the bacteria cultures that were added on day 3 did not have any significant effect on the moisture content. There was also no change in the soil pH as a result of addition of bacterial cultures or varying the concentration of bacterial culture added on day 3.

The volumes of leachates collected from the soil bio-cementation and bio-clogging experiment after the addition of 400 ml water for *Azotobacter chroococcum* and *Lactobacillus fermentum* are shown in Table 10.

The results showed that increasing the concentration of bacteria from 25 to 100% in the bacterial culture added on day 3 did not have any significant effect on the amount of leachate collected for both bacteria. However, the leachates collected from the soils receiving *Azotobacter chroococcum* (205 ml) were much larger than those collected from the soils receiving *Lactobacillus fermentum* (105 ml). Also, the leachates collected from the control (received no bacterial treatment) were much larger (310 ml) than both soils treated with both bacterial cultures. In other words, 90 ml (22.5%), 190 ml (47.5%) and 295 ml (73.75%) of the added water on day 5 were retained by the control, the soil receiving *Azotobacter chroococcum* and the soil receiving *Lactobacillus fermentum*, respectively. This amount to a water conservation of 100 ml (25%) and 205 ml (51.25%) for the soil receiving *Azotobacter chroococcum* and the soil receiving *Lactobacillus fermentum*, respectively.

The results obtained from the study showed that it is feasible to use growing cultures of *Azotobacter chroococcum* and *Lactobacillus fermentum* to produce a water insoluble levan. The polymer can be used as a plugging agent to plug the pores of the high permeability sandy soils. Upon production of levan, pore spaces would be reduced and, hence, the hydraulic conductivity would be substantially reduced. In addition to producing levan, these bacteria also produce gelatinous agents and elemental compounds that cause soil bio-cementation as shown in Figure 12.

Table 8. Biomass and levan and conversion efficiencies.

Parameter	<i>Azotobacter chroococcum</i>	<i>Lactobacillus fermentum</i>
Cell Count (Cells/mL)	8.23 x 10 ⁸	7.29 x 10 ⁸
Maximum Biomass (g /L)	3.6	3
Time to reach Maximum Biomass (h)	22	26
Minimum Biomass (g /L)	0.16	0.13
Time to reach Minimum Biomass (h)	68	72
Starting Sucrose (g/L)	50	50
Final Sucrose (g/L)	1.6	1.5
Time to Reach Final Sucrose (h)	56	61
Maximum Levan (g/L)	14	17
Time to Reach Maximum Levan	24	28
Theoretical Levan Yield (g levan/g sucrose)	0.395	0.395
Experimental Levan Yield (g levan/g sucrose)	0.248	0.371
Levan Production Efficiency (%)	62.78	93.92
Theoretical Biomass Yield (g cells/g sucrose)	0.13	0.13
Experimental Biomass Yield (g cells/g sucrose)	0.074	0.062
Biomass Production Efficiency (%)	47.69	56.92

Table 9. Soil moisture content and pH before the addition of water on day 5.

Parameter	Bacteria	Value				
		Control	Treatment 2	Treatment 3	Treatment 4	Treatment 5
Moisture Content (%)	<i>Azotobacterchroococcum</i>	19.6	22.4	22.3	22.2	22.3
	<i>Lactobacillus fermentum</i>	19.7	25.4	25.4	25.3	25.4
pH	<i>Azotobacterchroococcum</i>	8.1	8.3	8.2	8.3	8.2
	<i>Lactobacillus fermentum</i>	8.1	8.2	8,3	8.3	8.2

The bacteria could be grown in the laboratory either in the non-polysaccharide producing mode or in the polysaccharide producing mode. The first would permit distribution of the bacteria to the lower soil layers but would delay the production of the polysaccharide due to the extension of the lag period required to produce the enzyme (levansucrase).

Improving soil properties using biological techniques such as gleization, bio-grouting or bio-cementation and bio-plugging or bio-sealing has been reported by many authors. Kumariad and Xiang [33] stated that bio-grout is an excellent technique for reducing the permeability of porous soils and improving their mechanical properties. Mujab et al. [34] reported that bio-cementation binds soil particles

together leading to increased soil strength and stiffness against wind erosion. Ivanove and Chu [36] evaluated the application of bio-cementation and bi-clogging techniques for reducing the porosity and hydraulic conductivity of soils and found facultative and microaerophilic bacteria to be the most suitable organisms for these techniques. McConkey et al. [40] applied an enhanced gleization technique into irrigation canal and reduced water seepage by 30%. Ghaly [42] developed an enhanced bio-sealing mechanism for earthen manure storage using levan producing microorganism and reported that the infiltration rate was affected by the soil type and was correlated to the percentage of sand in the soil. Knapenetal [47] studied the effect of microbiotic crust on soil erodibility by wind and reported a

37% reduction in soil detachment. Ghaly et al. [49] studied the plugging effect of levan produced by *C* in earthen manure storage and found the bacteria converted sucrose into levan under field condition and the exopolysaccharide plugged the pores of a highly porous soil. Stewart and Folger [50] used polymer producing bacteria to modify soil profiles for enhanced oil recovery and reported that the bacteria utilized sucrose to produce exopolymer which created plugged regions of the porous media leading to enhanced oil recovery. Ramsay et al. [54] used *Bacillus licheniformis* to produce water insoluble levan that was used as a selective plugging agent in microbial enhanced oil recovery under a

temperature of 55°C, a pH between 6 and 9, a pressure less than soota and a salt concentration of 4%.

CONCLUSION

The result showed that the sandy soil used in this study had a particle density of 2.58 g/cm³, a bulk density of 1.6 g/cm³ and a porosity of 37.98 %. The particle size varied from 0.150 to 2.000 mm with most of the soil particles having a diameter in the range of 0.425-0.850 mm indicating that the soil was free of silt and clay. This soil has a loose texture, low organic matter, low nutrient content, high infiltration rate, low water holding capacity.

Table 10. Leachates collected after the addition of 400ml water on day 5.

Time (h)	Bacteria	Leachate (ml)				
		Control	Treatment 2	Treatment 3	Treatment 4	Treatment 5
12	<i>Azotobacterchroococcum</i>	309	205	205	205	205
	<i>Lactobacillus fermentum</i>	310	105	104	105	104
24	<i>Azotobacterchroococcum</i>	310	206	205	205	205
	<i>Lactobacillus fermentum</i>	311	105	105	105	104
48	<i>Azotobacterchroococcum</i>	310	206	205	205	204
	<i>Lactobacillus fermentum</i>	311	105	108	105	105




Figure 10. Bio-cementation and clogging of soil.

Azotobacter chroococcum are capable of producing levan from sucrose and have ability to fix molecular nitrogen and produce growth hormones, and therefore increase the soil fertility and stimulate plant growth, *Lactobacillus fermentum* bacteria are a levan producing bacteria. The viscosity of the culture broth was the same as that of water. The polymer can

be used as a plugging agent to plug the pores of high permeability soils. The levan yield was 0.248 glevan/g sucrose (62.78% of theoretical yield) and 0.371 glevan/g sucrose (93.92% of theoretical yield) for *Azotobacter chroococcum* and *Lactobacillus fermentum*, respectively. The cell yield was 0.074 g cell/g sucrose (47.69 % of

theoretical yield) and 0.062 g cell/g sucrose (56.92 % of theoretical yield).

The results showed that increasing the concentration of bacteria had not significant effect on the amount of leachate collect for both bacteria. However, the leachates collected from the soils receiving *Azotobacter chroococcum* were much larger than those collected from the soils receiving *Lactobacillus fermentum*. Also, the leachates collected from the control (received no bacterial treatment) were much larger than soils treated with both bacterial cultures. These microorganisms can be used together for production of levan from sucrose, fixing nitrogen in soil and producing growth hormones, thereby improving the water holding capacity of the soil and enhancing its nutrient content and stimulating plant growth.

ACKNOWLEDGEMENTS

The project was carried out in the Bioengineering Laboratory of the Department of Agricultural Engineering, Faculty of Agriculture, Cairo. The authors appreciate the assistance provided by the Laboratory Manager Ms. D. M. El Nakib and the Workshop Supervisor Mr. A. M. Abdel Haleem.

COMPETING INTRESTS

The authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors. All authors contributed equally in various roles including formulation research goals, development of methodology, performing the experiments and analysing data and writing the initial draft. The corresponding author coordinated the research activity as agreed by all authors. All authors read and approved the final manuscript.

REFERENCES

- Shafy AH, El Shaharty A, Regelsberger M (2010) Rainwater in Egypt: Quality, distribution and harvesting. *Mediterr Mar Sci* 11: 245-257.
- Maksoud A (2018) Estimation of temperature and rainfall in Egypt. *Asian J Adv Res Rep* 1: 1-22.
- Ali WH (2013) Suitability of Egypt deserts for sustainable urban development. *Dev Countries Stud* 3: 164-173.
- Karasapan O, Shah S (2018) Egypt's population: Boom then bust. Brookings Institution, Washington, DC, USA.
- CAPMPS (2019) Current and predicted Egyptian population. Central Agency for Public Mobilization and Statistics, Cairo, Egypt.
- MALR (2019) Food export and import. Ministry of Agriculture and land Reclamation, Cairo, Egypt.
- Habib IM, Morsy AA (2003) Land Reclamation and Improvement. Cairo University Open Education Centre. Giza, Egypt (Arabic with English translation).
- DRC (2018) Land reclamation program in Egypt. Desert Research Centre, Ministry of Agriculture and Land Reclamation, Cairo, Egypt.
- Alary V, Aboul-Naga A, Osman MA, Daoud I, Abdelrahman S, et al. (2012) Desert land reclamation program and family desert land dynamics in western desert of the Nile Delta (Egypt), 1960-2010. *World Development*, 104: 140-153.
- NRCS (1999) Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. Agriculture Handbook Number 436 (2nd Edition). Natural Resource Conservation Services, United States Department of Agriculture, Washington DC, USA, P869.
- Pankova EI, Gerasimova MI (2012) Desert soils: Properties, pedogenic process and classification. *Arid Ecosystems* 2: 69-77.
- Suleiman FH, Sadeq SA (1992) Inventory and classification of desert lands. Cairo University Press, Giza, Egypt (Arabic with English translation).
- El Shwarby MU (1961) Soil Chemistry. Anglo-Egyptian Bookshop, Cairo, Egypt (Arabic with English translation).
- Alaily F (1986) Cracks in sandy soils of the extreme arid part of Sahara. *Soil Science Hamburg* 3: 1023-1024.
- Petit M, Montaigne E, El Hadad-Gauthier F, Alvarez-Coque GMG, Mattas K, et al. (2015) Sustainable agricultural development: Challenges and approaches in Southern and Eastern Mediterranean countries. Cooperative Management, Springer International Publishing, New York, USA.
- Shehata M (2016) Ground water in Egypt: Reality and future prospects. Paper presented at the Physics of the Earth and Treasures Conference, Cairo, Egypt (Arabic with English translation).
- Indraratna B, Chu J (2005) Ground improvement: Case histories. Elsevier Oxford, UK.
- Monnie F (2016) Effect of biochar on soil physical properties, water use efficiency and growth of maize in a sandy loam soil. MPHIL Thesis, University of Ghana Digital collections, Department of Agricultural Engineering, College of Basic and Applied Science, School of Engineering, University of Ghana, Legon Boundary, Accra, Ghana.

19. Ding Y, Liu Y, Liu S, Li Z, Tan X, et al. (2016) Biochar to improve soil fertility: Review. *Agron Sustain Dev* 36: 36-48.
20. Jeffery S, Verheijen FGA, Van Der Velde M, Bastos AC (2011) A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agric Ecosys Environment* 144: 175-187.
21. Major J, Rondon M, Molina D, Riha SJ, Lehmann J (2010) Maize yield and nutrition during 4 years after biochar application to a Colombian savanna oxisol. *Plant Soil* 333: 117-128.
22. Glaser B, Lehmann J, Zech W (2002) Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal: A review. *Biol Fertil Soils* 35: 1719-1730.
23. Abu Dreeh OEA (2017) Effect of organic matter addition on the growth of Abu Sabeen (*Sorghum Bicolor*) in sandy soil. Technical Report, University of Sudan for Science and Technology, Khartoum, Sudan (Arabic with English translation).
24. Gomez-Sagasti MT, Hernandez A, Artetxe U, Garbisu C, Becerril JM (2018) How valuable are organic amendments as tools for the phyto-management of degraded soils? The Knowns, Known Unknowns, and Unknowns. *Front Sustainable Food Sys* 2: 1-16.
25. Bhogal A, Nicklson FA, Rollet A, Taylor M, Litterick A, et al. (2018) Improvements in the quality of agricultural soils following organic material additions depend on both the quantity and quality of the materials. *Appl Front Sustain Food Sys* 2: 1-12.
26. Lal R (2009) Challenges and opportunities in soil organic matter research. *Eur J Soil Sci* 60: 158-169.
27. Smith P, Lutfalla S, Riley WJ, Torn MS, Schmidt MWI (2017) The changing faces of soil organic matter research. *Eur J Soil Sci* 69: 23-30.
28. Spagnoli G (2018) A review of soil improvement with non-conventional grouts. *Int J Geotechnical Eng.*
29. Kazemian S, Huat BBK, Prasad A, Barghochi M (2010) A review of stabilization of soft soils by injection of chemical grouting. *Aus J Basic Appl Sci* 4: 58625868.
30. Kazemian S, Huat BBK (2009) Assessment and comparison of grouting and injection methods in geotechnical engineering. *Eur J Sci Res* 27: 234-247.
31. Karol RH (2003) *Chemical Grouting and Soil Stabilization* (3rd Edition). Marcel Dekker, New York, USA.
32. Krizek RJ (1985) Chemical Grouting in Soils Permeated by Water. *J Geotech Eng* 111: 898-911.
33. Kumari D, Xiang W (2018) Review on biologically based grout material to prevent soil liquefaction for ground improvement. *Int J Geotech Eng* 13: 48-53.
34. Mujah M, Shahin MA, Cheng L (2017) State-of-the-Art Review of Bio-cementation by Microbially Induced Calcite Precipitation (MICP) for Soil Stabilization. *Geomicrobiol J* 34: 524-537.
35. Wang Z, Zhang N, Cai G, Jin Y, Ding N (2017) Review of ground improvement using microbial induced carbonate precipitation (MICP). *Mar Georesour Geotec* 35: 1135-1146.
36. Ivanov V, Chu J (2008) Applications of microorganisms to geotechnical engineering for bio-clogging and bio-cementation of soil in situ. *Rev Environ Sci Biotechnol* 7: 139-153.
37. Whiffin VS, van Paasse LA, Harkes MP (2007) Microbial carbonate precipitation as a soil improvement technique. *Geomicrobiol J* 24: 417-423.
38. Bockheim JG, Hartemink AE (2017) *Soil-forming process. The soils of Wisconsin*, Springer International Publishing, New York, USA.
39. Bockheim JG, Gennadiyev AN (2000) The role of soil-forming process in the definition of taxa in soil taxonomy and the world soil reference base. *Geoderma* 95: 53-72.
40. McConkey BG, Reimer CD, Nicholaichuk W, Jame YW (1990) Sealing of earthen hydraulic structures with enhanced gleization and sodium carbonate: I. Laboratory study on the effect of freeze-thaw cycle and drying interval. *Can Agric Eng* 32: 163-170.
41. McConkey BG, Reimer CD, Nicholaichuk W, Jame YW (1990) Sealing of earthen hydraulic structures with enhanced gleization and sodium carbonate: II. Application for lining an irrigation canal. *Can Agric Eng* 32: 171-176.
42. Ghaly AE (1989) Enhanced biological sealing for earthen manure storages. *Proceedings of the 11th International Congress on Agricultural Engineering, A.A. Balkema Publisher, Amsterdam*, pp: 399-408.
43. Armanise E, Simmons RW, Ahn S, Garbout A, Doerr SH et al. (2018) Soil sealing development under simulated rainfall: Structural, physical and hydrological dynamics. *J Hydrol* 556: 211-219.
44. Belnap J, Gardner JS (1993) Soil microstructure in soils of the Colorado Plateau - the role of the cyanobacterium *Microcoleus vaginatus*. *West N Am Nat* 53: 40-47.
45. Bowker MA, Maestre FT, Escolar C (2010) Biological crusts as a model system for examining the biodiversity-ecosystem function relationship in soils. *Soil Biol Biochem* 42: 405-417.

46. Jeffery S, Harris JA, Rickson RJ, Ritz K (2009) The spectral quality of light influences the temporal development of the microbial phenotype at the arable soil surface. *Soil Biol Biochem* 41: 553-560.
47. Knapen A, Poesen J, Galindo-Morales, De Baets S, Pals A (2007) Effects of microbiotic crusts under cropland in temperate environments on soil erodibility during concentrated flow. *Earth Surf Proc Land* 32: 1884-1901.
48. Assouline S, Mualem Y (2000) Modeling the dynamics of seal formation: Analysis of the effect of soil and rainfall properties. *Water Resour Res* 36: 2341-2349.
49. Ghaly AE, Arab F, Mahmoud NM, Higgins J (2007) Production of Levan by *Bacillus licheniformis* for use as a soil sealant in earthen manure storage. *Am J Biochem Biotechnol* 2: 47-54.
50. Stewart TL, Fogler HS (2001) Biomass plug development and propagation in porous media. *Biotechnol Bioeng* 72: 353-363.
51. Han YW, Clarke MA (1990) Production and characterization of microbial levan. *J Agric Food Chem* 38: 393-396.
52. Snikanth R, Reddy CHSS, Siddartha G, Ramaiah MJ, Uppuluri KB (2014) Review on production, characteristics and application of microbial levan. *Carbohydrate Polym* 120: 102-114.
53. Park JK, Khan T (2009) Other microbial polysaccharides: pullulan, scleroglucan, elsinan, levan, alternant, dextran. *Handbook of Hydrocolloids*, Woodhead Publishing, Elsevier Massachusetts, USA.
54. Ramsay JA, Cooper DG, Newfeld KJ (1989) Effect of oil reservoir conditions and the production of water-insoluble levan by *Bacillus licheniformis*. *Geomicrobiol J* 7: 155-156.
55. Chen X, Gao H, Ploehn HJ (2013) Montmorillonite levan nanocomposites with improved thermal and mechanical properties. *Carbohydrate Polym* 101: 565-573.
56. Srikanth R, Siddartha G, Reddy CHS, Harish BS, Ramalah MJ et al. (2015) Antioxidant and anti-inflammatory levan produced from *Acetobacter xylinum* MCIM2526 and its statistical optimization. *Carbohydrate Polymers* 123: 8-16.
57. Silbir S, Dagbagli S, Yegin S, Baysal T, Goksungur Y (2014) Levan production by *Zymomonas mobilis* in batch and continuous fermentation systems. *Carbohydrate Polym* 99: 454-461.
58. Jathore NR, Bule MV, Tilay AV, Annapure US (2012) Microbial levan from *Pseudomonas fluorescens*: Characterization and medium optimization for enhanced production. *Food Sci Biotechnol* 21: 1045-1053.
59. Sarilmiser HK, Oner ET (2014) Investigation of anti-cancer activity of linear and aldehyde-activated levan from *Halomonas smyrnensis* AAD6T. *Biochem Eng J* 92: 28-34.
60. Shih IL, Yu YT, Shieh CJ, Hsieh CY (2005) Selective production and characterization of levan by *Bacillus subtilis* (Natto) Takahashi. *J Agric Food Chem* 53: 8211-8215.
61. Zhurina D (2009) Identification and potential characterization of transcriptional regulators involved in temperature-dependent expression of levansucrase in *Pseudomonas syringae*. PhD Thesis, Jacob University Bermen, Vegesack, Bermen, Germany.
62. Youssef GA, Youssef YAS, Talha S, El-Aassar SA (2014) Increased fructosyltransferase (levansucrase) production by optimizing culture condition from *Pediococcus acidilactici* strain in shaking batch cultures. *Life Sci J* 11: 33-47.
63. Combie J (2006) Properties of levan and potential medical uses. Chapter: 13, *Polysaccharides for Drug Delivery and Pharmaceutical Applications*, pp: 263-269.
64. Dahech I, Belghith KS, Hamden K, Feki A, Belghith H et al. (2011) Antidiabetic activity of levan polysaccharide alloxan-induced diabetic rats. *Int J Biol Macromol* 49: 742-746.
65. Gupta SK, Das P, Singh SK, Akhtar MS, Meena DK, et al. (2011) Microbial levan, an ideal prebiotic and immuno-nutrient in aquaculture. *World Aquaculture* 61-66.
66. Gupta SK, Pal AK, Sahu NP, Saharan N, Mandal SC, et al. (2014) Dietary microbial levan ameliorates stress and augments immunity in *Ciprinus cario fry* (Linnaeus, 1758) exposed to sublethal toxicity fipronil. *Aquac Res* 45: 893-906.
67. Sezer AD, Kazak H, Öner ET, Akbuğa J (2011) Levan-based nanocarrier system for peptide and protein drug delivery: Optimization and influence of experimental parameters on the nanoparticle characteristics. *Carbohydr Polym* 84: 358-363.
68. Sarilmiser K, Ates O, Ozdemir G, Arga KY, Öner ET (2015) Effective stimulating factors for microbial levan production by *Halomonas smyrnensis* AAD6T. *J Biosci Bioeng* 119: 455-463.
69. Freitas F, Alves VD, Reis MAM (2011) Advances in bacterial exopolysaccharides: From production to biotechnological applications. *Trends Biotechnol* 29: 388-398.
70. Manandhar S, D'Souza N, Vidhate S (2009) Water soluble levan polysaccharide biopolymer electrospun fibers. *Carbohydr Polym* 78: 794-779.

71. Narula N (2000) *Azotobacter* in Sustainable Agriculture. Vedams Books Ltd, New Delhi, India.
72. Loewenberg JR, Reese ET (1957) Observations on microbial fructosans and fructosanases. *Can J Microbiol* 3: 643-650.
73. Moonmangmee S, Kawabata K, Tanaka S, Toyama H, O. Adachi O, et al. (2002) A novel polysaccharide involved in the pellicle formation of *Acetobacteracetii*. *J Biosci Bioeng* 2: 192-200.
74. Tomulescu C, Stoica R, Secenco C, Casarica A, Moscovici M (2016) Levan: A mini review. *Scientific Bulletin Series F: Biotechnology, Vol XX*: 309-320.
75. Hernandez L, Arrieta J, Menedez C, Vasquez R, Coego R, et al. (1995) Isolation and enzymatic properties of levansucrase secreted by *Acetobacter diazotrophicus* SRT4, a bacterium associated with sugar cane. *Biochem J* 309: 113-118.
76. Arrieta J, Hernandez L, Geogo A, Suarez V, Balmori E, et al. (1996) Molecular characterization of the levansucrase gene from the endophytic sugarcane bacterium *Acetobacter diazotrophicus* SRT4. *Microbiol* 142: 1077-1082.
77. Támara Y, Hormaza JV, Pérez C, León A, Arrieta J (1999) Structural analysis and optimized production of fructo-oligosaccharides by levansucrase from *Acetobacter diazotrophicus* SRT4. *Biotechnol Lett* 21: 117-121.
78. Batista F, Hernandez RL, Fernandez JR, Arrieta J, Menedez C et al. (1999) Substitution of Asp-309 by Asn in the Arg-Asp-Pro (RDP) motif of *Acetobacter diazotrophicus* levansucrase affects sucrose hydrolysis, but not enzyme specificity. *Biochem J* 337: 503-506.
79. Perumpuli PABN, Watanabe T, Toyama H (2014) Pellicle of thermotolerant *Acetobacterpasteurianus* strains: characterization of the polysaccharides and of the induction patterns. *J Biosci Bioeng* 2: 134-138.
80. Minakami H, Entani E, Tayama K, Fujiyama S, Masai H (1983) Isolation and characterization of a new polysaccharide-producing *Acetobacter* sp.. *Agric Biol Chem* 84: 2405-2414.
81. Tayama K, Minakami H, Entani E, Fujiyama S, Masai H (2014) Structure of an acidic polysaccharide from *Acetobacter* sp. NBI 1022. *Biol Chem* 48: 9595-966.
82. Jansson PE, Lindberg J, Wimalasiri KS, Dankert MA (1993) Structural studies of acetan, an exopolysaccharide elaborated by *Acetobacter xylinum*. *Carbohydr Res* 245: 303-310.
83. Wong HC, Fear AL, Calhoun RD, Eichinger GH, Mayer R, et al. (1990) Genetic organization of the cellulose synthase operon in *Acetobacter xylinum*. *Proc Nat Acad Sci USA* 87: 8130-8134.
84. Gonzalez-Garcinuno A, Tabemero A, Sanchez-Alvarez JM, Galan MA, del Valle EMM (2017) Effect of bacteria type and sucrose concentration on levan yield and its molecular weight. *Microb Cell Fact* 191: 1-11.
85. Bansal A, Singh K, Karnwal A (2019) Effective abiotic factors on production of levan by microorganisms: A review. *IJBS* 4: 1-6.
86. Taberero A, Gonzalez-Garcinuno A, Sanchez-Alvarez JM, Galan MA, del Valle EMM (2017) Development of a nanoparticle system based on a fructose polymer: Stability and drug release studies. *Carbohydr Polym* 160: 26-33.
87. Pabst MJ (1977) Levan and levansucrase of *Actinomyces viscosus*. *Infect Immun* 15: 518-526.
88. Warner TN, Miller CH (1978) Cell-associated levan of *Actinomyces viscosus*. *Infect Immun* 22: 266-274.
89. Miller CH, Somers PJB (1978) Degradation of Levan by *Actinomyces viscosus*. *Infect Immun* 15: 518-526.
90. Igarashi T, Takahashi M, Yamamoto A, Etoh Y (1987) Purification and characterization of levansucrase from *Actinomyces viscosus* ATCC 19246. *Infect Immun* 55: 3001-3005.
91. Han YW (1990) Production and characterization of microbial Levan. *Agric Food Chem* 38: 393-396.
92. Yamasato K, Akagawa M, Oishi N, Kuraishi H (1982) Carbon substrate assimilation profile and other taxonomic features of *Alcaligenes faecalis*, *Alcaligenes* and *Achromobacter xulosoxydans*. *JGAM* 18: 195-213.
93. Srinivasan S, Quastel JH (1958) Enzymatic syntheses of oligo- and polysaccharides containing D-Glucosamine. *Science* 127: 143-144.
94. Wilkinson JF, Dudman WF, Aspinall CG (1955) The extracellular polysaccharide of *Aerobacter aerogenes* A3 (S1) (*Klebsiella* type 54). *Biochem J* 59: 446-451.
95. Evans TH, Hibbert H (1946) *Advances in Carbohydrate Chemistry*, Academic Press, New York, New York.
96. Takeshita M (1873) Translucent colony form of the gram-negative, levan-producing bacterium, *Aerobacter levanicum*. *J Bacteriol* 16: 503-506.
97. Feingold DS, Gehatia M (1957) The structure and properties of levan, a polymer of D-fructose produced by cultures and cell-free extracts of *Aerobacter levanicum*. *J Polym Sci* 23: 783-790.
98. Song KB, Bae KS, Lee YB, Lee KY, Rhee SK (2000) Characteristics of levan fructotransferase from *Arthrobacter ureafaciens* K2032 and difructose

- anhydride IV formation from levan. *Enzyme Microb Technol* 72: 212-218.
99. Tanaka K, Karigane T, Nishika F, Yoshida N (1983) Action of levan fructotransferase of *Arthrobacter ureafaciens* on levan oligosaccharides. *J Biochem* 94: 1569-1578.
 100. Tanaka K, Shimonishi M, Kitagaki M, Ikanaga M (1990) Action of levan fructotransferase of *Arthrobacter ureafaciens* on three oligosaccharides containing a bifurcateresidue. *Agric Biol Chem* 54: 815-817.
 101. Hestrin S, Goldblum J (1953) Levanpolyase. *Nature* 172: 1064-1047.
 102. De La Vega MG, Celudo FJ, Faneque A (1991) Production of exocellular polysaccharide by *Azotobacterchroococcum*. *Appl Biochem Biotechnol* 30: 273-284.
 103. Han YW (1990) Microbial Levan. *Adv Appl Microbiol* 35: 171-194.
 104. Abou-Taleb K, Abdel-Monem M, Yassin M, Draz A (2015) Production, purification and characterization of levanopolymer from *Bacillus lentus* V8 Strain. *Br Microbiol Res J* 5: 22-32.
 105. Tian FS, Karboune, Hill A (2014) Synthesis of fructo-oligosaccharides and oligolevans by the combined use of levansucrase and endo-inulinase in one-step bi-enzymatic system. *Innov Food Sci Emerg Technol* 22: 230-238.
 106. Xavier JR, Ramana KV (2017) Optimization of levan production by cold-active *Bacillus licheniformis* ANT 179 and Fructo-oligosaccharide synthesis by its levansucrase. *Appl Biochem Biotechnol* 181: 986-1006.
 107. Kekez BD, Gojic-Cvijovic GD, Jakovljevic DM, Kojic JS, Markovic MD, et al. (2015) High levan production by *Bacillus licheniformis* NS032 using ammonium chloride as the sole nitrogen source. *Appl Biochem Biotechnol* 175: 3068-3083.
 108. Mamay D, Wahyuningrum, Hertadi R (2015) Isolation and characterization of levan from moderate halophilic bacteria *Bacillus licheniformis* BK AG21. *Process Chem* 16: 292-298.
 109. Larpin S, Sauvageot N, Pichereau V, Laplace J, Auffray Y (2002) Biosynthesis of exopolysaccharide by a *Bacillus licheniformis* strain isolated from ropy cider. *Int J Food Microbiol* 77: 1-9.
 110. Van Dyk SJ, Ah Kee NL, Frost CL, Pletschke BI (2012) Extracellular polysaccharides production in *Bacillus licheniformis* SVD1 and its immunomodulatory effect. *Bioresourc* 7: 4976-4993.
 111. Strube CP, Homann A, Gammer M, Jahn D (2011) Polysaccharide synthesis of the levansucrase SacB from *Bacillus megaterium* is controlled by distinct surface motifs. *J Biol Chem* 286: 17593-17600.
 112. Tanaka K, Kawaguchi H, Ohno K, Shohji K (1981) Enzymic formation of difructose anhydride IV from bacterial levan. *J Biochem* 90: 1545-1548.
 113. Zhang T, Li R, Qian H, Mu W, Miao M (2014) Biosynthesis of levan by levansucrase from *Bacillus methylotrophicus* SK 21.002. *Carbohydr Polym* 101: 975-981.
 114. Li R, Zhang T, Jiang B (2014) Purification and characterization of an intracellular levansucrase derived from *Bacillus methylotrophicus* SK 21.002. *Biotechnol Appl Biochem* 62.
 115. Jadan J, Narnoliya LK, Agarwal N, Singh SP (2019) Catalytic biosynthesis of levan and short-chain fructo-oligosaccharides from sucrose-containing feedstocks by employing the levansucrase from *Leuconostocmesenteroides* MTCC10508. *IJBM* 127: 486-495.
 116. Han YW, Watson MA (1992) Production of microbial levan from sucrose, sugarcane juice and beet molasses. *J Ind Microbiol* 9: 257-260
 117. Han YW (1989) Levan production by *Bacillus polymyxa*. *J Ind Microbiol* 4: 447-452.
 118. Liu J, Luo J, Ye H, Zeng X (2012) Preparation, antioxidant and antitumor activities in vitro of different derivatives of levan from endophytic bacterium *Paenibacillus polymyxa* EJS-3. *Food Chem Toxicol* 50: 767-772.
 119. Jensen SL, Diemer MB, Lundmark M, Larsen FH, Blennow A, et al. (2016) Levansucrase from *Bacillus subtilis* hydrolyses β -2, 6 fructosyl bonds in bacterial levans and in grassfructans. *Int J Biol Macromol* 85: 514-521.
 120. Ing-Lung S, Yun-Ti Y, Chwen-Jen S, Chien-Yan H (2005) Production and Characterization of Levan by *Bacillus subtilis* (Natto) Takahashi. *J Agric Food Chem* 53: 8211-8215.
 121. Abdel-Fattah AM, Gamal-Eldeen AM, Helmy WA, Esawy MA (2012) Antitumor and antioxidant activities of levan and its derivative from the isolate *Bacillus subtilis* NRC1aza. *Carbohydr Polym* 89: 314-322.
 122. Benigar E, Dogsa I, Stopar D, Jamnik A, Cigiæ IK (2014) Structure and dynamics of a polysaccharide matrix: Aqueous solutions of bacterial levan. *Langmuir* 30: 4172-4182.
 123. Ahmed F (2005) Production of levansucrase from *Bacillus subtilis* NRC 33a and enzymic synthesis of levan and fructo-oligosaccharides. *Microbiology* 51: 402-407.

124. Esawy MA, Ahmed EF, Helmy WA, Mansour NM, El-Senousyand WM (2011) Production of levansucrase from novel honey *Bacillus subtilis* isolates capable of producing antiviral levans. *Carbohydr Polym* 86: 823-830.
125. Vaidya M, Prasad DT (2012) Thermostable levansucrase from *Bacillus subtilis* BB04, an isolate of banana peel. *J Biochem Technol* 3: 322-327.
126. Gonçalves BCM, Mantovan J, Ribeiro MLL, Borsato D, Celligoi MAPC (2013) Optimization production of thermo active levansucrase from *Bacillus subtilis* Natto CCT 7712. *J Appl Biol Biotechnol* 1: 1-8.
127. Abdul Razack S, Velayutham V, Thangavelu V (2013) Medium optimization for the production of exopolysaccharide by *Bacillus subtilis* using synthetic sources and agro wastes. *Turkish J Biol* 37: 280-288.
128. Abdel-Fattah AF, Mahmoud DAR, Esawy MAT (2005) Production of levansucrase from *Bacillus subtilis* NRC 33a and enzymatic synthesis of levan and fructo-oligosaccharides. *Curr Microbiol* 51: 402-407.
129. Southerland LW (2004) *Microbial Exopolysaccharides: Structure diversity and function versatility of polysaccharides*. CRC Press, New York, USA.
130. Sucawara M, Haramaki R, Monaka S, Ezura H, Okazaki S (2007) Rhzobiotoxine production in *Agrobacterium faciens* e58 by *Bradyrhizobium elkiirtx* ACDEFG genes. Annual Reports of Osaka International Center for Biotechnology, Osaka University, Osaka, Japan.
131. Sudtachat, N., N. Ho, S. Eda, H. Mitsui and K.Minamisawa. 2007. Analysis of metabolic features of naturally occurring aromatic compounds in *Bradyrhizobium japonicum* based on array and gene distribution. Annual Reports of Osaka International Center for Biotechnology, Osaka University, Osaka, Japan.
132. Dake, M. 2005. Biodegradable polymers: Renewable nature, life cycle and application. *Microbial Factories* (V. Kalia Ed), New Delhi, India.
133. Rosenberg, e., E. F. DeLong, S. Lory, E. Stackebrandt and F. Thompson. 2014. *The Prokaryotes: Alphaproteobacteria and Betaproteobacteria*. Springer Publishing Company, New York, USA.
134. Liu, Q., S. Yu, T. Zhang, B. Jiang and W. Mu. 2017. Efficient biosynthesis of levan from sucrose by a novel levansucrase from *Brenneria goodwinii*. *Carbohydrate Polymers*, 157: 1732-1740.
135. Xu, W., Q. Liu, Y. Bai, S. Yu, T. Zhang, B. Jiang and W. Mu. 2017. Physicochemical properties of a high molecular weight levan from *Brenneria* sp. EniD312. *International Journal of Biological Macromolecules*, 109: 810-818.
136. Gao, S., X. Qi, D. J. Hart, H. Gao and Y. An. 2017. Expression and characterization of levansucrase from *Clostridium acetobutylicum*. *Journal of Agriculture and Food Chemistry*, 65 (4): 867-871.
137. Dahech, I., H. Ben Ayed, K. S. Belghith, H. Belghith and H. Mejdoub. 2013. Microbial production of levanase for specific hydrolysis of levan. *International Journal of Biological Macromolecules*, 60: 128-133.
138. Dias, F. F. and J. V. Bhat. 1964. Nutritional properties of *Corynebacterium laevaniformans*. *Antone van Leeuwenhoek Journal of Microbiology*, 30(1): 176-184.
139. Chen, Y. F., Y. N.Yin, X. M. Zhang and J. H. Guo. 2007. *Curtobacterium flaccumfaciens* pv. *beticola*, A New Pathovar of Pathogens in Sugar Beet. *Plant Disease*, 91(6): 667-684.
140. Wuerges, J., L. Caputi, M. Cianci, S. Boivin, R. Meijersand S. Benini. 2015. The crystal structure of *Erwinia amylovora* levansucrase provides a snapshot of the products of sucrose hydrolysis trapped into the active site. *Journal of Structural Biology*, 191(3):290-8.
141. Gross, M., G. Geier, K. Rudolph and K Geidar. 1991. Levan and levansucrase synthesized by the fire blight pathogen *Erwinia amylovora*. *Physiological and Molecular Plant Pathology*, 40(6): 371-381.
142. Keith, J. A, B. J. Wiley, D. A. Zorfass, S. Arcidiacono, J. M. Mayer and D. Kaplan. 1989. The Production, purification and properties of the biopolymer levan produced by the bacterium *Erwinia Herbicola*. Technical Report No. 01760-5000, Science and Advanced Technology Directorate, Research, Development and Engineering Center, United States Army, Natick, Massachusetts, USA.
143. Keith, J. A, B. J. Wiley, D. A. Zorfass, S. Arcidiacono, J. M. Mayer and D. Kaplan. 1991. Continuous culture system for production of biopolymer Levan using *Erwinia herbicola*. *Biotechnology and Bioengineering*, 38(5):557-560.
144. Inthanovong L, Tian F, Khodadadi S, Karboune (2013) Properties of *Geobacillus stearothermophilus* levansucrase as potential biocatalyst for the synthesis of levan and fructo-oligosaccharides. *Biotechnol Prog* 29: 1405-1415.
145. Li Y, Triccas JA, Ferenci T (1997) A novel levansucrase gen cluster in *Bacillus stesrothermophilus* ATCC12980. *Biochem Biophys Act* 1353: 203-206.
146. Ua-Arak T, Jakob F, R. F. Vogel RF (2017) Fermentation pH modulates the size distributions and

- functional properties of *Gluconobacter ralbodus* TMW 2.1191 Levan. *Front Microbiol* 8: 1-11.
147. De Muynck C, Pereora CSS, Naessens M, Parmentier S, Soetaert W, et al. (2007) The genus *Gluconobacter* oxydans: Comprehensive overview of biochemistry and biotechnological applications. *Crit Rev Biotechnol* 27: 141-171.
 148. Jakob F, Rudi DM, Vogel F (2012) Comparison of novel GH 68 levansucrases of levan-overproducing *Gluconobacter* species. *AAB* 1(e2): 6-14.
 149. Velazquez-Hernandez ML, Baizabal-Aguirre VM, Cruz-Vazquez F, Trejo-Contreras MJ, Fuentes-Ramirez LE, et al. (2010) *Gluconacetobacter diazotrophicus* levansucrase is involved in tolerance to NaCl, sucrose and desiccation, and in biofilm formation. *Arch Microbiol* 193: 137-149.
 150. Park NH, Choi HJ, Oh DK (2005) Lactosucrose production by various microorganisms harboring levansucrase activity. *Bacteriol Lett* 27: 495-497.
 151. Serrato RV, Meneses CH, Vidal MS, Santana-Filho AP, Iacomini M, et al. (2013) Structural studies of an exopolysaccharide produced by *Gluconacetobacter diazotrophicus* Pal5. *Carbohydr Polym* 98: 1153-1159.
 152. Banguela A, Arrieta JG, Rodríguez R, Trujillo LE, Menéndez C (2011) High levan accumulation in transgenic tobacco plants expressing the *Gluconacetobacter diazotrophicus* levansucrase gene. *J Biotechnol* 154: 93-98.
 153. Kommann H, Duboc P, Marison I, von Stockar U (2003) Influence of Nutritional Factors on the Nature, Yield, and Composition of Exopolysaccharides Produced by *Gluconacetobacter xylinus* I-2281. *Appl Environ Microbiol* 69: 6091-6098.
 154. Sarilmiser HK, Ates O, Ozdemir G, Argaand KY, Oner ET (2015) Effective stimulating factors for microbial levan production by *Halomonas smyrnensis* AAD6T. *J Biosci Bioeng* 119: 455-463.
 155. Poli A, Kazak H, Gürleyendag B, Tommonaro G, Pieretti G, et al. (2012) High level synthesis of levan by a novel *Halomonas* species growing on defined media. *Carbohydr Polym* 78: 651-657.
 156. Brandt JU, Jakob F, Behr J, Geissler AJ, Vogel RF (2016) Dissection of exopolysaccharide biosynthesis in *Kozakiabaliensis*. *Microb Cell Fact* 170: 1-13.
 157. Dutta A, Das D, Goyal A (2001) Purification and characterization of fructan and fructansucrase from *Lactobacillus fermentum* AKJ15 isolated from Kodo ko jaanr, a fermented beverage from north-eastern Himalayas. *Int J Food Sci Nutr* 63: 216-224.
 158. Badel S, Bernardiand T, Michaud P (2011) New perspectives for *Lactobacilli* exopolysaccharides. *Biotechnol Adv* 29: 54-66.
 159. Heinemann C, Johan ET, Veig VH, Janssen DB, Busscher HJ (2000) Purification and characterization of a surface-binding protein from *Lactobacillus fermentum* RC-14 that inhibits adhesion of *Enterococcus faecalis* 1131. *Biotechnol Lett* 90: 117-180.
 160. Galle S, Avendt EK (2014) Exopolysaccharides from sourdough lactic acid bacteria. *Crit Rev Food Sci Nutr* 54: 891-901.
 161. Anwar MA, Kralj S, Pique AV, Leemhuis H, van der Maarel MJEC L (2010) Inulin and levan synthesis by probiotic *Lactobacillus gasserii* strains: Characterization of three novel fructansucrase enzymes and their fructanproducts. *Microbiol* 156: 1264-1274.
 162. Diez-Municio M, de la Rivas B, Jimeno ML, Munoz R, Moreno FJ, et al. (2013) Enzymatic synthesis and characterization of fructo-oligosaccharides and novel maltosylfructosides by inulosucrase from *Lactobacillus gasserii* DSM 20604. *Appl Environ Microbiol* 79: 4120-4131.
 163. Sims IM, Frese SA, Walter J, Loach D, Wilson M, et al. (2011) Structure and functions of exopolysaccharide produced by gut commensal *Lactobacillus reuteri* 100-23. *The ISME Journal* 5: 1115-1124.
 164. Van Hijum SA, Bonting K, van der Maarel MJEC, Dijkhuizen L (2001) Purification of a novel fructosyltransferase from *Lactobacillus reuteri* strain 121 and characterization of the levan produced. *FEMS Microbiol Lett* 205: 323-328.
 165. Kaditzky SV, Vogel RF (2008) Optimization of exopolysaccharide yields in sourdoughs fermented by *Lactobacilli*. *Eur Food Res Technol* 228: 291-299.
 166. Ni D, Xu W, Bai Y, Zhang W, Zhang T, et al. (2018) Biosynthesis of levan from sucrose using a thermostable levansucrase from *Lactobacillus reuteri* LTH5448. *Int J Biol Macromol* 113: 29-37.
 167. Tiekling M, Ehrmann MA, Vogel RF, Gänzle MG (2005). Molecular and functional characterization of a levansucrase from the sourdough isolate *Lactobacillus sanfranciscensis* TMW 1.392. *Appl Microbiol Biotechnol* 66: 655-663.
 168. Han, J., X. Xu, C. Gao, Z. Liu and Z. Wu. 2016. Levan-producing *Leuconostocitreum* strain BD1707 and its growth in tomato juice supplemented with sucrose. *Appl Environ Microbiol* 82: 1383-1390.
 169. Ortiz-Soto ME, Olivares-Illana V, Lopez-Munguia A (2004) Biochemical properties of inulosucrase from

- Leuconostocitreum CW28 used for inulin synthesis. *Biocatal Biotransfor* 22: 275-281.
170. Bounaix MS, Gabriel V, Robert H, Morel S, Simeon MR, et al. (2010) Characterization of glucan producing *Leuconostoc* strains isolated from sourdough. *Int J Food Microbiol* 144: 1-9.
 171. Xu Y, Coda R, Shi Q, Tuomainen P, Katina K, Tenkanen M (2017) Exopolysaccharides production during the fermentation of soybean and fava bean flours by *Leuconostoc mesenteroides* DSM 20343. *J Agric Food Chem* 65: 2805-2815.
 172. Liu X, Luo Y, Mohamed OA, Liu D, Wei G (2014) Global transcriptome analysis of *Mesorhizobium alhagi* CCNWXJ12-2 under salt stress. *BMC Microbiology* 14.
 173. Priest PG, Goodfellow M (2000) *Applied Microbia Systematics*. Springer Science Publisher, New York, USA.
 174. Kimbrel J (2012) Genome-enabled discovery and characterization of type III effect-encoding genes of plant symbiotic bacteria. Ph D Thesis, Oregon State University, Corvallis, Oregon, USA.
 175. Reeve W, Yates RJ, Twari R, Gu W (2013) Complete genome sequence of *Mesorhizobium australicum* strain type WSM2073T. *Stand Genomic Sci* 4: 410-419.
 176. Das K, Rajawat MVS, Saxena AK, Parasanna R (2017) Development of *Mesorhizobium ciceri* based biofilms and analyses of their antifungal and plant growth promoting activity in cichpea challenged *Fusarium* wilt. *Ind J Microbiol* 57: 48-59.
 177. Chen WX, Li GS, Qi YL, Wang ET, Yuan HL, Li JL (1991) *Rhizobium huakuii* sp. nov. isolated from the root nodules of *Astragalus sinicus*. *Int J Syst Bacteriol* 41: 275-280.
 178. Kawaharada YS, Eda K, Miamisawa, Mitsui H (2007) A *Mesorhizobium loti* mutant with reduced glucan content defective invasion of its host plant *Lotus japonicus*. *Annual Reports of Osaka International Center for Biotechnology, Osaka University, Osaka, Japan*.
 179. Kelly SJ, Muszynski A, Kawaharada Y, Hubber AM, Suuivan JT, Sandal N, Carlson RW, Stougaard J, Ronson CW (2013) Controlled requirement for exopolysaccharides in *Mesorhizobium-Lotus* symbiosis. *Mol Plant Microbe Interact* 26: 319-329.
 180. Ray RC (2005) *Microbial Biotechnology in Agriculture and Aquaculture*. SP Science publishing, Enfield, New Hampshire, USA.
 181. Bae IY, Oh IK, Lee S, Yoo SH, Lee GH (2008) Rheological characterization of levan polysaccharides from *Microbacterium laevaniformans*. *Int J Biol Macromol* 42: 10-13.
 182. Krichevsky MI, Howell A, Lim JRS (1969) Levan Formation by *Odontomyces viscosus*. *J Dent Res* 48: 938-942.
 183. Xu X, Gao C, Liu Z (2016) Characterization of the levan produced by *Paenibacillus bovis* sp. nov. BD3526 and its immunological activity. *Carbohydr Polym* 144: 178-186.
 184. Hang F, Wang Q, Chen W (2017) Effect of oxygen supply on milk-clotting activity expressed by *Paenibacillus* spp. strain BD3526. *Food Sci Technol* 82: 437-445.
 185. Petrov KK, Petrova P (2017) Sugar transport systems involved in fructooligosaccharides utilization by the probiotic bacterium *Pediococcus acidilactici*. *Compte Rendus de Academic Bulgare des Sciences* 70: 1263-1270.
 186. Haworth N, Stacey M (1940) The chemistry of immunology saccharides. *Ann Rev Biochem* 17: 97-114.
 187. Lyne RR, Peat S, Stacey M (1940) Polysaccharides: The constitution of certain levans formed by bacterial action. *J Chem Soc* 47: 237-244.
 188. Fuchs A (1956) Synthesis of levan by pseudomonads. *Nature* 178: 921.
 189. Alamäe T, Visnapuu T, Mardo K, Mäe A, Zamfir AD (2012) Levan sucrases from *Pseudomonas* bacteria: Novel approaches for protein expression, assay of enzymes, fructooligosaccharides and hetero-oligofructans. *Carbohydr Chem* 38: 176-191.
 190. Al Qaysi SAS (2016) Levan production using *Pseudomonas brassicacearum* isolated from rhizosphere soil of cowpea farm in Iraq. *Iraqi J Biotechnol* 15: 83-89.
 191. Kasapis S, Morris ER, Gross M, Rudolph K (1994) Solution properties of levan polysaccharide from *Pseudomonas syringae* pv. *phaseolicola*, and its possible primary role as a blocker of recognition during pathogenesis. *Carbohydr Polym* 23: 55-64.
 192. Laue H, Schenk A, Li H, Lambertsen L, Neu TR, Molin S, Ullrich MS (2006) Contribution of alginate and levan production to biofilm formation by *Pseudomonas syringae*. *Microbiol* 152: 2909-2918.
 193. Yoo SE, Yoon E, Cha, Lee H (2004) Antitumor activity of levan polysaccharides from selected microorganisms. *Int J Biol Macromol* 34: 37-41.
 194. Kim H, Yang JY, Lee HG, Cha J (2001) Cloning and sequence analysis of a levansucrase gene from *Rahnella aquatilis* ATCC15552. *J Microbiol Biotechnol* 11: 693-699.
 195. Kim HJ, Park HE, Kim MJ, Lee KG, Yang JY, Cha L (2003) Enzymatic characterization of a recombinant

- levansucrase from *Rahnella aquatilis* ATCC 15552. *J Microbiol Biotechnol* 13: 230-235.
196. Karunaratne DN (2012) The complex world of polysaccharides. In Tech Open Publishing Company, London, England.
197. Tikhonovich IN, Provorov NA, Romanov VI, Newton WE (1995) Nitrogen fixation fundamental and application. Proceedings of the 10th International congress on Nitrogen Fixation, Kluwer Academic Publisher, Norwell, Massachusetts, USA.
198. Leshner R, Gerencser VF (1977) Levan production by a strain of *Rothia*: Activation of complement resulting in cytotoxicity for human gingival cells. *J Dent Res* 56: 1097-1105.
199. Willner SZ, Imam, Hader I (1977) Case Report in Cardiology. 1-3.
200. Hill MJ (2018) Microbial metabolism in digestive tract. CRC Press, New York, USA.
201. Franken J, Brandt BA, Tai SI, Bauer FF (2013) Biosynthesis of levan: A bacterial polysaccharide, in yeast *Saccharomyces cerevisiae*. *PLoS One* 8: 1-14.
202. Elorza MV, Villanueva JR, Sentandre R (1977) The mechanism of catabolite inhibition of invertase by glucose in *Saccharomyces cerevisiae*. *Biochem Biophys Acta* 475: 103-112.
203. Ebisu S, Kato K, Kotani S, Misaki A (1975) Structural differences in fructans elaborated by *Streptococcus mutans* and *Streptococcus salivarius*. *J Biotechnol* 78: 879-887.
204. Newbrun E, Baker S (1968) Physico-chemical characteristics of the levan produced by *Streptococcus salivarius*. *Carbohydr Res* 6: 165-170.
205. Moosavi-Nasab MB, Layegh L, Aminlari MB (2010) Microbial production of levan using date syrup and investigation of its properties. *Int J Biol Biomol Agric Food Biotechnol Eng* 4: 603-607.
206. Vigants A, Hicke HG, Marx SP (2001) A simple and efficient method for the purification of membrane-bound levansucrase from *Zymomonas mobilis*. *Curr Microbiol* 42: 415-418.
207. Calazans GMT, Lima RC, de Franca FP, Lopes CE (2000) Molecular weight and antitumor activity of *Zymomonas mobilis* levans. *Int J Biol Macromol* 27: 245-247.
208. Bekers MD, Upite E, Kaminska J, Laukevics M, Grube A et al. (2005) Stability of levan produced by *Zymomonas mobilis*. *Process Biochem* 40: 1535-1539.
209. Melo IR, Pimentel MF, Lopes CE, Calazans GMT (2007) Application of fractional factorial design to levan production by *Zymomonas mobilis*. *Braz J Microbiol* 38: 45-51.
210. Ananthalakshmy VK, Gunasekaran P (1999) Optimization of levan production by *Zymomonas mobilis*. *Braz Arch Biol Technol* 42: 291-298.
211. Shaheen S, Aman A, Siddiqui NN (2017) Influence of metal ions, surfactants and organic solvents on the catalytic performance of Levansucrase from *Zymomonas mobilis* KIBGE-IB14. *J Basic Appl Sci* 13: 41-46.
212. Santos VAQ, Garcia-Cruz VL, Del Bianchi VL (2014) Effect of initial pH in levan production by *Zymomonas mobilis* immobilized in sodium alginate. *Acta Sci Technol* 36: 349-354.
213. De Oliveira MR, de Silva RSSF, Buzato JB, Celligoi MAPC (2007) Study of levan production by *Zymomonas mobilis* using regional low-cost carbohydrate sources. *Biochem Eng J* 37: 177-183.
214. Garrity GM, Brenner DJ, Krieg NR, Staley JT (2005) *Bergey's Manual of Systematic Bacteriology*. Springer Publishing, New York, USA.
215. Sneath PHA, Mair NS, Sharpe ME, Holt JG (1986) *Bergey's Manual of Systematic Bacteriology*. Lippincott Williams and Wilkins, Baltimore, Maryland, USA.
216. Breed RS, Murray EGD, Smith NR (1957) *Bergey's Manual of Determinative Bacteriology*. The Williams and Wilkins Company: Baltimore, Maryland, USA.
217. Ghaly, A. E. and R. M. Ben-Hassan. 1994. Kinetics of batch production of single cell protein from cheese whey. *Appl Biochem Biotechnol* 50: 79-92.
218. Ghaly AE, Mahmoud NS (2006) Optimum condition for measuring dehydrogenase activity of *Aspergillus niger* using TTC. *Am J Biochem Biotechnol* 2(4): 186-194.
219. Ramsay (1987) Microbial products in enhanced oil recovery. PhD thesis. Department of Chemical Engineering, Faculty of Engineering, McGill University, Montreal, Quebec, Canada.
220. Borji AF, Borji, Jourani A (2017) A new method for the determination of sucrose concentration in a pure and impure systems: Spectrophotometric method. *Int J Anal Chem* 217/8214120: 1-6.
221. Schipper L (2013) Soil properties: Relative sizes of sand, silt and clay particles. Science Learning Hub, University of Waikato, Hamilton, New Zealand. Accessed on December 3rd 2019. Available online at: <https://www.sciencelearn.org.nz/resources/957-soil-properties>

222. Braud AC, Hartman, Lesturgez G (2005) Management of tropical soils for sustainable agriculture: A holistic approach for sustainable development of problem soils in the tropics. KhonKaen University Press, KhonKaen, Thailand.