

Identification and Evaluation of Antibiotic Sensitivity Pattern of Bacteria Associated with Gastro-Intestinal Disorder in Damaturu Yobe State, Nigeria

Muhammad S Abadallah¹, Ibrahim U Imam² and Muhammad Ali^{3*}

¹Desert Research Monitoring and Control Centre, Yobe State University, Damaturu, Nigeria

²Department of Vector and Parasitology, Nigerian Institute for Trypanosomiasis and Onchocerciasis Research, Kaduna, Nigeria

³Department of Microbiology, Federal University, Gusau, Nigeria.

Received February 25, 2019; Accepted February 27, 2019; Published September 05, 2019

ABSTRACT

Enteric bacterial pathogens are one of the major causes of food borne gastroenteritis in humans and remain an important health problem worldwide. The study was aimed to identify and determine antibiotic sensitivity pattern of some bacteria associated with gastro-intestinal disorder among patients attending General Sani Abacha Specialist Hospital in Damaturu Yobe State, Nigeria. A total of thirty (30) sterile swab sticks containing diarrheic stool samples were collected from patients complaining of gastro-intestinal infection attending the hospital were streaked onto the surface of plate containing Nutrient agar using standard method. The bacterial isolates were identified based on Gram staining, cultural characteristics and biochemical test. The bacteria isolates were subjected to antibiotic susceptibility testing using the agar disc diffusion method. The results showed that six different bacteria were isolated and identified namely; *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, Enterobacter, Shigella and Salmonella. The findings of the study on sensitivity pattern of the isolates against antibiotics indicated Sensitivity testing of the isolate against antibiotics indicated that most of the isolates were resistant to rifampicin, streptomycin, gentamicin and chloramphenicol, but sensitive to ciprofloxacin, amoxicillin, norfloxacin and levifloxacin. Findings of this study justify the presence of enteric bacteria as one of the cause of gastric disorder.

Keywords: Bacteria, Identification, Sensitivity pattern, Gastro-intestinal disorder

INTRODUCTION

Food borne diseases are an important cause of morbidity and mortality worldwide. There are over 200 different types of illness that may be transmitted by food [1]. The causes of food borne illness are bacteria, viruses, parasites and chemicals and bacterial contamination is the most common cause of illness [2]. The human gastrointestinal tract is home to a complex microbial community with more than 100 trillion members of prokaryotic cells. Most food borne bacterial infections cause self-limiting diarrhea, however, systemic infection and death can occur, particularly in vulnerable groups such as the elderly, people with diminished immunity or infants and young children [3]. Bacteria have accounted for more than 70% of deaths associated with food borne transmission [2,4].

Gastrointestinal infections are responsible for a high rate of morbidity in the world, although its highest incidence occurs in the developing countries, according to the World Health Organization (WHO) global health occur [1]. 700 million diarrheal diseases each year and kills 760,000 children under 5 each year for severe dehydration for this reason it is

considered the second leading cause of death in the population at this age, even in developed countries, children under 3 years of age have on average 3 diarrhea episodes per year [5]. A broad spectrum of microorganisms may be responsible for the infection and present in common the same clinical manifestations being the most common symptoms diarrhea, vomiting, fever, abdominal pain, malaise, anorexia and dehydration at different levels of severity [6]. The medical diagnosis is usually relatively simple once a correct anamnesis and epidemiological history

Corresponding author: Muhammad Ali, Department of Microbiology, Federal University, Gusau, Nigeria, E-mail: alimuhd4real@gmail.com

Citation: Abadallah MS, Imam IU & Ali M. (2019) Identification and Evaluation of Antibiotic Sensitivity Pattern of Bacteria Associated with Gastro-Intestinal Disorder in Damaturu Yobe State, Nigeria. J Allerg Res, 1(2): 19-24.

Copyright: ©2019 Abadallah MS, Imam IU & Ali M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

has been made, it is important to perform microbiological laboratory studies to determine the etiological agent responsible for the disease and initiate appropriate treatment [7]. Diarrhea is generally defined as three or more unformed stools evacuations per day and it is often associated with other enteric symptoms [8]. The etiology of diarrhea, abdominal pain, and digestive disorders may be related to more than forty infectious pathogens which could be grouped in viruses, bacteria and parasites [8]. Viral gastroenteritis is a common cause of morbidity and mortality in humans around the world and is considered the most common infectious gastroenteritis. The most common causative agents are Rotavirus, Norovirus, Sapovirus, Adenovirus and Astrovirus [9]. Some bacterial pathogens are able to develop persistent gastrointestinal symptoms and other complications; among the most frequently been identified: *E. coli*, *Campylobacter*, *Salmonella* and *Shigella* [7]. In a study conducted by Nas et al. [10] on identification of some enteric bacterial isolates from diarrheic stool sample of adult male patients attending Murtala Muhammad Specialist Hospital Kano, Nigeria showed the presence of *Escherichia coli*, *Salmonella* spp., *Klebsiella* spp., *Pseudomonas* spp., *Citrobacter* spp. and *Shigella* spp. with the following prevalence 22%, 19%, 14%, 12%, 14%, 19%, respectively. Considering the public importance of gastrointestinal infection, laboratory surveillance of such infection is utilized in many countries for safety and prevention efforts [11]. In this, we isolate and identify bacterial associated with gastro-intestinal disorder among patients attending General Sani Abacha Specialist Hospital Damaturu Yobe state Nigeria. The antibiotic sensitivity pattern of the isolates against antibiotics was also evaluated.

MATERIALS AND METHODS

Ethical approval

Ethical approval for the study was obtained from Ministry of Health Damaturu, Yobe State based on the consent of General Sani Abacha Specialist Hospital Damaturu Yobe state, Nigeria Ethical Committee.

Samples collection

A total of thirty (30) sterile swab sticks containing diarrheic stool samples were collected from patients complaining of gastro-intestinal infection attending Microbiology Department of General Sani Abacha Specialist Hospital Damaturu Yobe state, Nigeria. The samples were immediately transported to Biology Laboratory in the Department of Biological Science, Yobe State University Damaturu for bacteria isolation and identification.

Isolation of bacteria

The swab sticks containing stool samples were streaked onto the surface of plate containing Nutrient agar (Life save Biotech, USA) using standard method of Harley and Prescott [12]. The procedure was applied for each sample and the

plates were incubated at 37°C for 24 h. The presumptive colonies of each isolate on agar plates were further sub cultured to get pure culture. There covered pure isolates were preserved for further bacterial identification.

Identification of bacterial isolates

The bacterial isolates were identified based on Gram staining, cultural characteristics and biochemical test. Gram staining was done for each individual isolates according to method described by Holt et al. [13] and Sherman [14]. Colonial morphology and cultural characteristics of plates were made and recorded for the different growth on MacConkey agar (Life save Biotech, USA). The isolates were also characterized by biochemical tests (indole test, Methyl Red test, Vogues Proskauer test and Citrate utilization test) as well as Lactose fermentation Reaction, Nitrate reduction and motility tests by standard method given by Holt et al. [13]; Sherman [14] and Cheesbrough [15].

Gram staining

A drop of normal saline was placed on a well labeled clean grease-free glass slide using a sterile inoculating loop; a colony of an overnight culture of the bacterial isolate was emulsified with the normal saline to make a thin smear. The smear was air dried and then heat fixed. The slide was flooded with primary stain (crystal violet) for 30 s after which the stain was rinsed from the slide with water. The smear was flooded with Lugolr iodine (mordant) to fix the primary stain. The iodine was rinsed with water after 60 s. The slide was then flooded with a decolorizer (acetone) and rinsed off almost immediately. The counter stain; safranin was added and left for 30 s before being rinsed off. The stained smear was air dried and then observed under the microscope using 100x oil immersion objective lens of the microscope [14].

Indole test

Tryptophan broth was inoculated with an isolate of the test organism and incubated at 37°C for 24 h. About 0.5 ml of Kovack's reagents was added to the broth culture.

Methyl red test

MR-VP broth was inoculated with an isolate of the test organism using sterile inoculating loop and incubated at 37°C for 24 h. About 5 drops of Methyl-red reagent was added to the broth culture.

Voges Proskauer

MR-VP broth was inoculated with an isolate of the test organism using sterile inoculating loop and incubated at 37°C for 24 h. Six milliliters (6 ml) of 5% alpha naphthol was added followed by 0.2 ml of KOH. The tube was shaken gently and remained undisturbed for 5 min.

Citrate utilization test

Simmons’s citrate agar was streaked back and forth with inoculums of the test organism and incubated aerobically at 37°C for 24 h.

Nitrate reduction test

Nitrate broth was inoculated with an isolate of the test organism using sterile inoculating loop and incubated at 37°C for 24 h. A dropper full of sulfanilic acid and that of α-naphthalamine were added to the broth.

Motility test

A semi solid medium in a test tube was inoculated with an isolate of the test organism using straight sterile wire and making a single stab at the centre of the test tube. The test tube was incubated at 37°C and examined the stab line at various intervals to determine motility.

Antibiotic sensitivity test

The bacteria isolates were subjected to antibiotic susceptibility testing using the agar disc diffusion method as described by Bauer et al. [16]. Mueller Hilton agar (MHA) (Life save Biotech, USA) plates were inoculated with overnight culture of each isolate by streak plating method. The standard antibiotic sensitivity discs were then

aseptically placed at equidistance on the plates and allowed to stand for 1 h. The plates were then incubated at 37°C for 24 h. Sensitivity pattern of the isolates to Augmentin (30 µg/disc), Taravid (10 µg/disc), Streptomycin (30 µg/disc), Ampicillin (30 µg/disc), Gentamicin (20 µg/disc), Reflacin (10 µg/disc), Ceporex (30 µg/disc), Nalixidic acid (20 µg/disc), Ciprofloxacin (10 µg/disc) and Septrin (30 µg/disc), produced by Abtek pharmaceutical limited, were determined. Isolates were divided into three groups based on the zone of inhibition produced by the antibiotic disc; susceptible, intermediately susceptible and resistant according to the Clinical and Laboratory Standards Institute (CLSI) guideline; Performance Standards for Antimicrobial Susceptibility Testing [17]. The experiment was conducted in triplicate and the average value was calculated.

RESULTS

Cultural characteristics of the isolates

The Cultural characteristics of the recovered isolates on MacConkey agar is presented in **Table 1**. The result showed how the recovered isolate were characterized on the basis of colony morphology and staining characteristics. Colony morphology ranges from mucoid pink, non-mucoid darker pink, colorless colony with jagged edge, transparent colorless colony.

Table 1. Biochemical characterization of bacteria isolated from stool samples.

Isolate code	Gram staining	Cultural characteristics on MacConkey agar
IS ₁	Negative/rod	Non-mucoid, dry, flat and darker pink colony
IS ₂	Negative/rod	Mucoid pinkish colony
IS ₃	Negative/rod	Colorless colony with characteristic foul smell
IS ₄	Negative/rod	Produce pinkish colony
IS ₅	Negative/rod	Colorless colony with jagged edge
IS ₆	Negative/rod	Transparent colorless colony

Biochemical characteristics of the isolates

The biochemical characteristic of the isolates is presented in **Table 2**. The isolates were characterized based on indole, methyl-red, Voges Proskauer, citrate utilization and motility

tests. Nitrate reduction test and lactose and fermentation tests were also conducted. The results showed that six different bacteria species were isolated and identified namely; *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Enterobacter*, *Shigella* and *Salmonella*.

Table 2. Biochemical characterization of bacteria isolated from stool samples.

Code	IN	MR	VP	CI	LF	NR	MO	CA	Isolates
IS ₁	+	+	-	-	+	+	+	+	<i>Escherichia coli</i>
IS ₂	-	-	+	+	+	+	-	+	<i>Klebsiella pneumoniae</i>
IS ₃	-	+	-	+	-	+	+	+	<i>Proteus mirabilis</i>
IS ₄	-	-	+	+	+	+	+	+	<i>Enterobacter</i> spp.
IS ₅	+	+	-	-	-	+	-	-	<i>Shigella dysenteriae</i>
IS ₆	-	+	-	-	-	+	+	+	<i>Salmonella typhi</i>

Key: IN: Indole; MR: Methyl Red; VP: Voges Proskauer; CI: Citrate; LF: Lactose Fermentation; NR: Nitrate Reduction; MO: Motility Test; CA: Catalase Test

Antibiotic sensitivity test

The zone of inhibition of antibiotic sensitivity disc against the bacterial isolates is presented in **Table 3**. Most of the antibiotics are active against the isolates. *E. coli* is sensitive to norfloxacin, gentamicin, erythromycin, levifloxacin and chloramphenicol. *Klebsiella* is resistant gentamicin,

amoxicillin, rifampicin, ampicillin and chloramphenicol. *Proteus mirabilis* is resistant to ciproloxacin rifampicin, streptomycin, ampicillin and erythromycin. *Enterobacter* is sensitive to all antibiotics but resistant streptomycin and rifampicin. *Salmonella* and *Shigella* were resistant to most antibiotic tested but sensitive to ciprofloxacin, Amoxicillin, ampicillin and levofloxacin.

Table 3. Mean zone of inhibition of antibiotics against bacterial isolates.

Isolates	Antibiotics (µg/disc)/Average mean zone of inhibition (mm)									
	CIP (10)	NOR (10)	GEN (10)	AMO (20)	STR (30)	RIF (20)	ERY (30)	AMP (20)	LEV (20)	CHL (20)
<i>Escherichia coli</i>	16	22	21	10	17	10	21	10	20	19
<i>Klebsiella</i>	20	21	10	10	18	10	20	10	19	10
<i>Proteus mirabilis</i>	10	19	10	21	15	10	13	10	18	22
<i>Enterobacter</i> spp.	21	19	22	22	15	15	22	21	20	21
<i>Shigella</i> spp.	20	21	10	20	10	10	10	22	21	10
<i>Salmonella typhi</i>	21	10	10	21	10	10	10	20	19	10

Key: CIP: Ciprofloxacin; NOR: Norfloxacin; GEN: Gentamicin; AMO: Amoxil; STR: Streptomycin; RIF: Rifampicin; ERY: Erythromycin; AMP: Ampicillin; LEV: Levifloxacin; CHL: Chloramphenicol

DISCUSSION

The human gut is natural habitat for various bacteria species and majority of them participate in metabolic activities. Enteric bacteria are microbes that reside in the guts of animals and humans. However there are some among them that reside in intestinal tract of animals that can cause diseases and harsh reactions when human become infected with them [18]. They can cause a mild infection, such as food poisoning or severe community-infections like diarrhea. Such examples of enteric bacteria include *Salmonella*, *Escherichia coli*, *Shigella*, *Klebsiella*, *Campylobacter*, *Enterobacter*, *Yersinia*, *Vibrio* and *Citrobacter* [19].

In the presence study 6 different bacteria were isolated from stool samples of patients complaining of gastro-intestinal disorder. The bacterial isolate include *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Enterobacter*, *Shigella* and *Salmonella*. Bacterial were tested for gram staining and biochemical test (**Table 2**). Bacteriological analysis of the isolates was done using MacConkey agar. MacConkey agar is both selective and differential. The medium contain lactose as the sole fermentable source of carbohydrate. Some enteric bacteria ferment lactose while others do not. Colonies of bacteria that ferment lactose produce acid end product and thus the colony will turn pink while non-lactose fermenters produce translucent or colorless colony. *E. coli*, *Klebsiella* and *Enterobacter* were found to be lactose fermenter. *Salmonella*, *Shigella* and *Proteus mirabilis* do not ferment lactose. Several studies

were conducted to characterize bacteria associated with gastro-intestinal infection.

The finding of this study was in conformity with that of Nas et al. [10] on identification of some enteric bacterial isolates from diarrheic stool sample of adult male patients attending Murtala Muhammad Specialist Hospital Kano, Nigeria which found the presence of *Escherichia coli*, *Salmonella* spp., *Klebsiella* spp., *Pseudomonas* spp., *Citrobacter* spp. and *Shigella* spp. with the following prevalence 22%, 19%, 14%, 12%, 14%, 19%, respectively. Similar study was conducted by Obi et al. [19] on enteric bacterial pathogen in stools of residents of urban and rural regions of Nigeria, the results shows the most frequently encountered pathogens in rural area are *E. coli*, followed by *Salmonella*, *Shigella*, *Enterobacter* and *Campylobacter*. Similarly, the result of this study correspond to the one conducted others on bacteria isolated from diarrhea patients in Korea which reveals that *Escherichia coli* as one of the most prevalent bacteria, then followed by *Clostridium* and *Salmonella*. A study conducted by Lopez et al. [20] on Entero-pathogenic agents isolated in persistent diarrhea, the result shows that *Salmonella*, *Escherichia coli* and *Shigella* as the major enteric bacteria responsible for gastric disorder. This finding supported the present study.

On the sensitivity pattern of the isolates against the antibiotics used, most of the antibiotics are active against the isolates. Most of the antibiotics are active against the isolates. *E. coli* is sensitive to norfloxacin, gentamicin, erythromycin, levifloxacin and chloramphenicol. *Klebsiella*

is resistant gentamicin, amoxicillin, rifampicin, ampicillin and chloramphenicol. *Proteus mirabilis* is resistant to ciprofloxacin rifampicin, streptomycin and erythromycin. Enterobacter is sensitive to all antibiotics but resistant streptomycin and rifampicin. Salmonella and Shigella were resistant to most antibiotic tested but sensitive to ciprofloxacin, Amoxicillin, ampicillin and levifloxacin. The result of this study supported the result of a study conducted by Dessalegn et al. [21]. They found that *E. coli* were resistant to ampicillin (100%), Proteus was resistant to ampicillin (100%) and Salmonella isolates were susceptible to ciprofloxacin.

LIMITATIONS OF THE STUDY

This study was limited to certain bacterial pathogens and other medically important enteric organisms such as Campylobacter, Citrobacter, Vibrio, Aeromonas and Yersinia were left unidentified due to some technical problems.

CONCLUSION

The finding of the present study showed that *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, Enterobacter, Shigella and Salmonella represent some bacteria associated with gastro-intestinal disorder because they reside in intestinal tract of animals that can cause diseases and harsh reactions when human become infected with them. Sensitivity testing of the isolate against antibiotics indicated that most of the isolates were resistant to rifampicin, streptomycin, gentamicin and chloramphenicol, but sensitive to ciprofloxacin, amoxicillin, norfloxacin and levifloxacin. Findings of this study justify the presence of enteric bacteria as one of the cause of gastric disorder.

ACKNOWLEDGEMENT

The authors wish to acknowledge the staff of Microbiology department, General Sani Abacha Specialist Hospital Damaturu for sample provision. Thanks to technical staff of Biological Science Department Yobe State University for use of the department's laboratory facilities. Sincere thanks to Yobe State Government through Ministry of health for granting us with ethical clearance for the conduct of the research.

REFERENCES

- World Health Organization (2013) Enfermedades diarreicas [Online]. Available from: <http://www.who.int/mediacentre/factsheets/fs330/es/>
- Lynch MJ, Painter R, Woodruff and Braden C (2006) Surveillance for foodborne disease outbreaks -- United States, 1998-2002. CDC 55: 1-34.
- Kennedy MR, Villar DJ, Vugia T, Rabarsky-Ehr MM, Farley M, et al. (2014) Enteric bacteria isolated from diarrheal patients in Korea in 2004. Osong Public Health Res Perspect 6: 233-240.
- Hughes C, Gillespie IA, O'Brien SJ; The Breakdowns in Food Safety Group (2007) Food borne transmission of infectious intestinal disease in England and Wales, 1992-2003. Food Control 18: 766-772.
- Yurist-Doutsch S, Arrieta MC, Vogt SL, Finlay BB (2014) Gastrointestinal microbiota-mediated control of enteric pathogens. Annu Rev Genet 48: 361-382.
- Mengelle C, Mansuy JM, Prere MF, Grouteau E, Claudet I, et al. (2013) Simultaneous detection of gastrointestinal pathogens with a multiplex Luminex-based molecular assay in stool samples from diarrheic patients. Clin Microbiol Infect 19: E458.
- Dupont HL (2014) Acute infectious diarrhea in immunocompetent adults. N Engl J Med 370: 1532-1540.
- Zboromyrska Y, Vila J (2016) Advanced PCR-based molecular diagnosis of gastrointestinal infections: Challenges and opportunities. Expert Rev Mol Diagn 16: 631-640.
- Becker SL, Chatigre JK, Gohou JP, Coulibaly JT, Leuppi R, et al. (2015) Combined stool-based multiplex PCR and microscopy for enhanced pathogen detection in patients with persistent diarrhea and asymptomatic controls from Côte d'Ivoire. Clin Microbiol Infect 21: 591.e1-10.
- Nas F, Ali M, Yahaya A (2017) Identification of some enteric bacteria from diarrheic stool samples of patients attending Murtala Muhammad Specialist Hospital Kano, Nigeria. Int J Adv Acad Res Sci Technol Eng 3: 28-36.
- Kendall ME, Crim S, Fullerton K, Han PV, Cronquist AB, et al. (2012) Travel associated enteric infection diagnosed after return to the United State, Food Borne Diseases Active Surveillance Network (FoodNet), 2004-2009. Clin Infect Dis 54: S480-487.
- Presscot LM, Harley JP, Klein DA (2005) Microbiology. 6th Edn. McGraw Hill International edition, New York.
- Holt JG, Krieg NR, Senath PHA, Staley JT, Williams ST (1994) Bergey's Manual of Determinative Bacteriology. 9th Edn. Baltimore Md., Williams and Wilkins.
- Sherman N (2005) Microbiology: A laboratory manual. 6th Edn. ISBN 81: 265-267.
- Chessbrough M (2006) District laboratory practice in tropical countries. 2nd Edn, Part two. Cambridge University Press, pp: 80-85.

16. Bauer AW, Kirby WM, Sherris JC, Turck M (1996). Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 45: 493-496.
17. CLSI (2010) Performance standards for antimicrobial susceptibility testing. 20th Informational Supplement, Clinical and Laboratory Standards Institute Doc. M100eS20.
18. Singh V, Pandey PC, Jain DK (2013) A text book of Botany. 4th Edn. Rastogi Publication, India.
19. Obi CL, Coker AO, Epoke J, Ndip RN (1997) Enteric bacterial pathogen in stools of resident of urban and rural region in Nigeria. *J Diarrhrel Dis Res* 15: 241-247.
20. Lopez MD, Gonzales ES, Valdes DM, Fragoso AT, Albizu CJ (1996) Wales, 1992-2003. *Food Control* 18: 766-772.
21. Dessalegn L, Techalew S, Endale T (2014) Aerobic bacterial isolates from post-surgical wound and their antimicrobial susceptibility pattern: A hospital based cross-sectional study. *E3 J Med Res* 3: 18-23.