

Risk of Dementia in Head Injuries and the Role of Altered Cerebral Blood Flow

Satish G*

*Baylor Scott & White Medical Center-Centennial Hospital, USA.

Received March 15, 2021; Revised April 12, 2021; Accepted April 15, 2021

ABSTRACT

Traumatic brain injury and its consequences on cognitive function has been a long-debated topic. In this article we try to review available literature that tries to explain the consequences of head injury, the change in cerebral flow dynamics and other neurochemical changes that play a role in the development of neurocognitive deficits and other symptoms as a consequence of TBI.

Keywords: Dementia, Brain Injury, Cognitive deficits, Head Injury, TBI

CEREBRAL TRAUMATIC INJURY (CTI)

Traumatic brain injury is disruption of normal brain function caused by an external mechanical force. Force can be transmitted to the brain from impact to the skull, jaw or body. The constraints of shear force on the brain and blood vessels, which results from acceleration or a sudden and often rotational deceleration of the head, are responsible for most of the damage [1]. TBI can heal completely on its own or produce serious disability or even lead to death. Overall, falls are the main causes of TBI especially in young (under 14) and elderly people (65 years old and over). Motor vehicle accidents are the second most common leading causes of TBI and death in young adults. Two-thirds of TBIs occur in men, but lesions of the same severity may develop more unfavorably in women than in men. Participation in sports and being victims of assault are other common causes of head trauma [1]. In the workplace, the causes of TBI are falls, falling of objects, motor vehicle or machine accidents. certain professions with the most risk are construction, transport, agriculture, forestry, fishing and emergency medical services. Twenty percent of workplace injuries are believed to be attributable to falls on surfaces, uneven or wet or caused by objects out of place, and they, therefore, appear to be largely preventable. Brain trauma can be either "closed", that is, without lesions of the scalp, or "open", accompanied by lacerations of the scalp. A fractured skull or lacerations of the dura (the outermost and the most resistant of the three membranes separating the skull from the brain) can be associated. There may also be contusions or lacerations of the brain [2]. A closed linear skull fracture is insignificant in itself, except that it is a testament to the

intensity of the force exerted on the head. Fractures of closed skullcaps, however, may cause embarrassment (localized depression of the cranial vault) which can damage the brain (Levin, Shum and Chan, 2014). Open trauma is a cause of intracranial infections, such as meningitis, subdural empyema and brain abscess. Open trauma can be external or internal. In case of internal open trauma, the skull fracture can affect the paranasal sinuses or mastoid pneumatic cells at the base of the skull and, in combination with damage to the dura mater and arachnoid membrane, serve as bacterial contamination and lead to the aforementioned intracranial infections [3].

WHAT IS DEMENTIA?

Dementia can be explained as the loss of mental capacity which is considered to be sufficient for undertaking normal routine activities for more than six months, not present at the time of birth and not being associated with the alteration and loss of consciousness. Dementia is an umbrella term which covers several types of medical conditions which have common symptoms resulted due to the gradual death of brain cells. Loss within cognitive abilities due to dementia

Corresponding author: Satish G, Baylor Scott & White Medical Center-Centennial Hospital, USA, Tel: 7087147661; E-mail: drsatishgaddam@gmail.com

Citation: Satish G. (2022) Risk of Dementia in Head Injuries and the Role of Altered Cerebral Blood Flow. J Neurosurg Imaging Techniques, 7(1): 441-446.

Copyright: ©2022 Satish G. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

can lead to memory impairment along with drastic changes in behavior (Peterson et al., 2019). Chronic traumatic encephalopathy (CTE) is referred to be a progressive degenerative disease having a history of receptive brain trauma inclusive of symptomatic concussions and asymptomatic head injuries. Boxers were the first to be the sufferers of this type of dementia. Now CTE is considered to be widespread amongst soccer, football, ice hockey as well as other sports athletes who hold inherent risk of concussive or sub concussive injuries (Dewan et al., 2018). The word "dementia" is a generic term that groups together more than a hundred diseases that affect brain function. It is mainly the cognitive abilities, such as thinking, memory, orientation in time and space, as well as language, which are affected. As the disease progresses, they are more and more limited in their daily and / or professional activities and can become dependent on the help of others [4].

DEMENTIA OR SIMPLE OCCASIONAL FORGETFULNESS!

We are all subject to occasional forgetfulness, the frequency of which varies according to each one. There is no reason to worry if it happens from time to time that you no longer know whether you have locked the door or lost your glasses. But if you notice that your memory problems are more common and affect more than one area of your cognition, you may want to get it evaluated. That said, memory loss is just one symptom of dementia. Depending on its form, dementia can manifest in other ways, such as changes in social behavior [5].

IS THERE A CURE FOR DEMENTIA?

Dementia can have many causes. Some forms of dementia are reversible and can therefore be partially or even completely cured with appropriate treatment. This is the case, for example, with dementia due to a metabolic disease such as hypothyroidism or disorders linked to vitamin B12 deficiency. Most of the time, however, the brain is directly affected in causes of irreversible dementia. The best-known forms of these degenerative brain diseases are Alzheimer's disease and vascular dementia. Lewy body dementia, frontotemporal dementia and Parkinson's disease dementia. These degenerative forms of dementia cannot be cured or slowed down, with any known therapies. But with the right treatment, those affected can maintain their independence for longer and see their symptoms improve. Provided, however, that the diagnosis has been made as soon as possible by the specialists [6].

RELATIONSHIP BETWEEN DEMENTIA AND HEAD INJURIES

Head injuries, depending on their severity and the area of the brain that is affected, can be the cause of cognitive impairment and memory problems similar to those that occur in cases of dementia.

The sequel of injuries of this type can be, among others:

- Vertigo
- Headache
- Mood alteration
- Apathy
- Depressive or anxiety states
- Sleep disturbance
- Memory problems

There can be a significant loss of mental abilities in head injury patients (concentration, organizational skills, reasoning skills, decision-making for problem solving, etc.), of paying attention, and of doing things called "executive functions" (planning, organizing, using abstract reasoning, problem solving, and judgment).

In relation to the loss of memory of past and recent events and the acquisition of new memories, amnesia can be reached. Severe head injuries are becoming more common in developed countries. Traffic accidents, the practice of risky sports or work accidents are some of the most frequent causes. However, it is also common for this type of trauma to occur in habitual users of alcohol and other addictive substances [7]. Likewise, it is important to note that some patients with previous head injuries, whether multiple or isolated, develop dementia over time, characterized by the existence of problems when expressing themselves and understanding what is said to them (aphasia), the inability to make movements that you are asked to perform (apraxia) or the inability to describe an object (agnosia).

In fact, in studies carried out after the autopsy on people, and even in children, who have suffered serious accidents of this type, it has been found that in 30% of cases there are amyloid plaques that form shortly after the trauma and that are characteristic of some dementias. One of the best-known types of dementia due to head trauma is the so-called boxer dementia, which is the result of repeated trauma in the exercise of this sport and whose symptoms progress over time. When dementia is the consequence of a single brain injury, it usually does not progress and if it does, one must consider the existence of an additional pathology [8]. More and more data points to the relationship between head injuries and senile dementia. The term "senile dementia" is too broad and vague a concept, which is under review. This contains a significant number of symptoms, with varied origins. Likewise, the word senile, etymologically speaking, refers to old age, or to this period in people's lives. This is related to progressive degeneration and the loss of physical, motor and mental capacities due to advanced age. Likewise, it applies to degenerative damage produced in the brain, but which can occur before reaching the "third age". In any case, what the specialists are looking for is to give a name and

surname to the different symptoms related to dementia, according to their causes and other specificities [9].

ETIOLOGY OF TBI AND DEMENTIA

The most leading cause of TBI amongst the population includes motor vehicle accidents, falls, and sports-related injuries as well as assaults [10]. From amongst these falls, are rendered to be the leading cause TBI in every age. However, elderly persons aging 75 years and above reported highest rate of hospitalization due to TBI and hence long-term cognitive impairments or death because of fall. Moreover, other cause identified for TBI includes repetitive head trauma from contact sports as well as combat-related injuries in the head amongst soldiers that return from wars [11]. Additionally, brain injuries can also be the result of bullet wounds or any other injury which tends to penetrate within the brain or the skull [11]. Basically, there are two assumptions related to the etiology of TBI where the first is the anterior and frontal cortices are susceptible to neural contusion whereas, the second is the rotational and linear forces act over axon bundles resulting in axonal injury [12,13]. TBI is rendered to be the most established environmental risk for developing dementia. The recently available data over dementia indicates that (a) adequate amount of data for the connection between moderate to severe TBI and dementia having a higher risk level of developing dementia between 2 - 4 fold for such patients and (b) scarce evidence is available for the connection between mild TBI associated with the loss of consciousness and developing of dementia; (c) insufficient or inadequate evidence is available for understanding whether there exists an association between mild level of TBI without losing consciousness and development of dementia [14]. However, no evidence can be traced which indicates that a single episode of mild TBI can increase the risk of dementia. However, in boxers, the risk of developing dementia is deeply associated with the rounds boxed instead of the number of times the boxer was knocked off. This eventually suggests that repetitive mild TBIs which do not result in losing conscious might increase this risk level [15].

HEAD INJURIES: COMMON SUSPECTS IN SENILE DEMENTIA

Any type of head injury can have serious consequences for the person involved, as insignificant as it may seem. The episodes of brain trauma are responsible for a significant number of cases that lead to pathologies. Among these we find Alzheimer's and Parkinson's disease are more significant. The problem is that these head injuries seriously affect the lymphatic system, which is responsible for removing toxins from the brain. This exclusive brain cleansing program works during sleep hours. With it, all mental operations are left in optimal conditions to face a new day. A slight impact within this complex network of pipes creates difficulties in the expulsion of waste [16].

The most delicate part of this negative progression is that, unlike most cells in the human body, brain cells do not have regenerative capacities. For this reason, when dying there is no possibility of recovery or replacement. In the absence of adequate cleaning, toxic proteins can accumulate setting the stage for the onset of neurodegeneration.

HEAD TRAUMA: THE BEGINNING OF A DEGENERATIVE PROCESS THAT TAKES TIME

Experts assure that fatal consequences on brain cells do not usually occur simultaneously with traumatic episodes. It is a process that can take more time, depending on the severity of the injury itself. However, brain contusions cause immediate damage. They are the beginning of what has been called a "biochemical cascade". This refers to a path that can eventually lead to irreversible states of mental decline [17].

TRAUMATIC BRAIN INJURY AND DEVELOPMENT OF DEMENTIA AND THE ROLE OF CHANGES IN CEREBRAL FLOW DYNAMICS

Traumatic brain injury (TBI) has emerged as a very common clinical issue which is also a major source of mortality and morbidity in the current times. Common reasons associated with head injury include falls or accidents. Traumatic brain injuries can be classified as severe, moderate or mild depending upon whether the brain injury has resulted in unconsciousness and its duration as well as severity levels of symptoms [18]. Though, TBIs are usually classified to be mild since they do not pose a life-threatening situation, even this type of TBI can have long-lasting and severe effects. A much-feared consequence related to TBI is dementia and several studies indicate that experiencing TBI in early or in mid-life can lead towards the risk of developing dementia [18]. Some common areas related to cognitive impairment that are a result of TBI involves executive and attention functioning, memory information processing speed and others amongst those experiencing that of mild brain injury. Additionally, another observation indicates that repetitive mild brain trauma that are, most of the times experienced by professional sportsperson like footballers, hockey players and boxers can be associated with a high degree of chronic traumatic encephalopathy (CTE). Basically, CTE is a form of dementia having distinct pathologic and clinical features.

Cognitive deficiency in the longer run might incur despite of having a normal looking brain within traditional neuroimaging studies. The risk of dementia increases with the number and severity of head injuries, according to a large prospective study conducted on a Danish cohort, the results of which are published in the "Lancet Psychiatry". This study was indeed conducted on nearly 2.8 million people, for a total of 27,632,020 person-years. It shows that people with repeated head trauma have an increase in their risk of dementia by 24%, and the specific risk of Alzheimer's disease increase by 16%. The risk is also increased even

with a head injury of moderate severity (concussion), with an increase of 17% [19].

A DANISH STUDY

The link between TBI and dementia had already been suggested, mainly for veterans of the wars in Iraq and Afghanistan, and for certain sportsmen (football, American football, hockey, boxing). But the studies on the subject lacked robustness (too small samples and duration of follow-up, lack of information on the number, severity and date of head injuries, lack of a comparative control population) and the results were contradictory. This study is the first to follow a sample of sufficient size for a long time. It thus considered in the Danish registers the cases of head trauma that had led to hospitalization or an emergency room visit between 1977 and 2013. The association of the severity, number and date of head injury with the onset of dementia was adjusted for other risk factors such as diabetes, heart disease, depression and drug use [20]. However, other risk factors such as level of education were not taken into account. "It's important to understand how robust this study is," insists a commentary in the same issue of the "Lancet Psychiatry," Carol Brayne of Cambridge University School of Medicine.

A RISK THAT INCREASES WITH THE NUMBER OF HEAD INJURIES

Between 1977 and 2013, 4.7% of the 2.8 million people considered underwent at least one head trauma. And between 1999 and 2013, 4.5% of people over 50 were diagnosed with dementia (with an average age at diagnosis of 80.7 years). The risk of dementia increases with the number of traumatic injuries, from 22% with a single head trauma to 183% (almost three times as much) with 5 or more injuries. The study also found that a single severe head injury increased the risk of dementia by 35% while a moderate head trauma increased it by 17%. In addition, the earlier an individual suffers a head injury, the greater the risk of dementia. Gender also has an influence since men are slightly more at risk of developing dementia than women in the same situation. Finally, the study shows that people who have had traumatic fractures outside the sites of the head and neck are not at increased risk of dementia.

Every year, 50 million people around the world experience head trauma, and the main causes are falls, car accidents and assaults.

CLASSIFICATION OF TBI AND DEMENTIA

A traumatic brain injury incurs when an outside force hits hard over the head which is strong enough to cause human brain to violently move within the skull hence disrupting normal functioning of the brain. Depending upon whether the injury resulted in unconsciousness and how long it was, brain injury can be classified into mild injury (mTBI) when there is no loss of consciousness or in case of

unconsciousness, it doesn't last for more than 30 minutes. Moderate TBI is when the unconsciousness lasts for 30 minutes where severe brain injury means where unconsciousness exceeds 24 hours [21-23].

EPIDEMIOLOGY

According to Hyder [23], almost 10 million people across the globe are being affected by TBI on an annual basis. As a matter of fact, this figure is also underestimated as it excludes patients suffering from mTBI who tend to frequently dismiss their symptoms of TBI and fail to seek any type of medical treatment (almost 25%) or those who visit private clinics or treat themselves through primary care providers (almost 14%) and does not report their clinical findings [24,25]. Estimation carried out by The Centers for Disease Control and Prevention (CDC), almost 1.7 million Americans tend to suffer from severe TBI which can result in hospitalization, treating in emergency care units or even in deaths on an annual basis [14]. World Health Organization (2015) states that TBI has the potential to surpass various diseases and can become one of the leading causes of disability or death by the year 2020. A large number of unreported as well as reported cases, TBI are rendered to be a 'silent epidemic' of the world. In addition to this, Holm [26] estimates that almost 85% - 90% of all treated brain injuries can be categorized as mild which corresponds to an expected incidence 100-300/100,000 cases as per the task force of WHO [26]. The same study proposes that one from every three mTBI victims suffer from long term cognitive deficiencies [12], which can incur irrespective to normal results obtained from conventional neuroimaging studies. Based upon the data available, a calculation related to the percentage of people having this disease attributed to TBI is between 10% - 15% [14].

PATHOPHYSIOLOGY

Head injury tends to initiate an inflammatory force which releases amino acids like aspartate and glutamate along with free radicals which results in damaging of the tissues [27]. Other potential elements include endogenous opioid peptides like catecholamines, naloxone, thyrotropin-releasing hormone (TRH), acetylcholine, adenosine and lactate along with nitrous oxide [28]. Diffuse axonal injury (DAI) usually incurs after moderate to severe TBI and is an important neuropathological consequence of brain traumas. This aspect involves various abnormalities including direct damaging of the axonal cytoskeleton to that of incurring damage to the disruption of axoplasmic membrane transport, proteolysis, and swelling [29]. Many studies have been undertaken to determine the connection between inheritance of the apolipoprotein $\epsilon 4$ (APOE $\epsilon 4$) allele and the risk of developing dementia (particularly Alzheimer's disease post a head injury). A study carried out in Manhattan [30] people having a history of traumatic brain injury and the development of APOE $\epsilon 4$ allele were associated with a 10-fold increased risk of dementia whereas, APOE $\epsilon 4$ in the

absence of TBI resulted in only a 2-fold increased risk of dementia. The same study has found no evidence related to developing dementia because of TBI when APOE ϵ 4 is not present [30].

PATHOLOGY

Within rodent models, brain injury has resulted in neurodegeneration as well as progressive brain atrophy which can carry on for at least a year after the occurrence of the [30,31]. The various types of proteins that are associated with neurodegenerative diseases within humans have also been demonstrated for the accumulation of experimental brain injury in rodents. Amyloid precursor protein is up regulated on an immediate basis after the brain injury and β -amyloid peptide which accumulates over weeks as well as over months after a trauma [32]. This developed hypothesis holds therapeutic implications in the recent developments within dementia and related diseases and might have a very important role in the management of traumatic brain injuries [33]. More evidences which indicate that neurodegeneration after a head injury may share some similar features with Alzheimer's disease (AD) in imaging studies. Cerebral atrophy after an injury is not diffused instead it is regionally selective and those regions where this atrophy is the most prominent after an injury like amygdala, hippocampus, parietal and frontal cortices and precuneus, tends to overlap with the regions of predominant β -amyloid deposition, reduced uptake of glucose as well as progressive atrophy in AD [34]. The findings of these studies can be associated with common molecular mechanisms that are shared between TBI-related neurodegeneration and AD [14]. Though CTE and AD are the distinct elements. Several studies that were made over retired professional athletes who have suffered multiple concussions and have also developed dementia have reported very prominent tau-immunoreactive NFTs and astrocytic tangles; however, β -amyloid pathology was found in less in than half of the cases [14,22].

MEDICAL TREATMENT FOR DEMENTIA IN CASE OF A BRAIN INJURY

A brain injury which resulted in dementia can benefit from education and emotional support. This can include the following: A detailed discussion of all the different treatment modalities is beyond the scope of this topic.

Cognitive rehabilitation, behavior modification, family intervention, social services Medications including dopaminergic agents, Antidepressants, Antipsychotic agents, Mood stabilizers, Antiepileptic drugs. Other therapeutically strategies for dementia in head injury patients include activity changes, diet etc.

CONCLUSION

The risk of dementia resulted because of a brain injury is difficult to predict in certainty. It depends upon multiple factors including the seriousness of the injury. Some may

completely recover from severe injuries whereas others might suffer from disabilities for a long time even after milder injuries. Additionally, dementia resulting from a brain injury is different from other forms of dementia. Dementia resulting from a brain injury usually does not worsen with time. Multiple factors including changes in CSF flow dynamics, alterations in neuro chemicals, sex, age and other comorbidities all play a role in development and recovery of dementia in head trauma patients. Long term population-based studies are required to further the knowledge base in this specific set of patients.

REFERENCES

1. Centers for Disease Control and Prevention (2016) Injury prevention & control: Traumatic brain injury & concussion Available online at: <https://www.cdc.gov/traumaticbraininjury/data/rates.html>
2. Levin HS, Shum D, Chan RC (2014) Understanding traumatic brain injury: Current research and future directions. New York (NY): Oxford University Press.
3. Smith M (2008) Monitoring intracranial pressure in traumatic brain injury. *Anesth Analg* 106(1): 240-248.
4. Fratiglioni L, Paillard-Borg S, Winblad B (2004) An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol* 3: 343-353.
5. Launer LJ (2005) The epidemiologic study of dementia: A life-long quest? *Neurobiol Aging* 26: 335-340.
6. American Psychological Association (2012) Guidelines for the evaluation of dementia and age-related cognitive change. *American Psychologist* 67(1): 1-9.
7. Beason-Held LL, Goh JO, Kraut MA, O'Brien RJ, Ferrucci L et al. (2013) Changes in brain function occur years before the onset of cognitive impairment. *J Neurosci* 33: 18008-18014.
8. Tran HT, LaFerla FM, Holtzman DM, Brody DL (2011) Controlled cortical impact traumatic brain injury in 3xTg-AD mice causes acute intra-axonal amyloid- β accumulation and independently accelerates the development of tau abnormalities. *J Neurosci* 31: 9513-9525.
9. Plassman BL, Havlik RJ, Steffens DC, Burke JR, Newman TN et, al. (2000) Documented head injury in early adulthood and risk of Alzheimer's disease and other dementias. *Neurology* 55(8): 1158-1166.
10. Wintermark M, Sanelli PC, Anzai Y, Tsiouris AJ, Whitlow CT (2015) Imaging Evidence and Recommendations for Traumatic Brain Injury: Conventional Neuroimaging Techniques. *J Am Coll Radiol* 12: e1-e14.

11. Okie S (2005) Traumatic brain injury in the war zone. *N Engl J Med* 352: 2043-2047.
12. Büki A, Povlishock JT (2006) All roads lead to disconnection? Traumatic axonal injury revisited. *Acta Neurochir (Wien)* 148: 181-184.
13. Meythaler JM, Peduzzi JD, Eleftheriou E, Novack TA (2001) Current concepts: Diffuse axonal injury-associated traumatic brain injury. *Arch Phys Med Rehabil* 82: 1461-1471.
14. Shively S, Scher AI, Perl DP, Diaz-Arrastia R (2012) Dementia Resulting from Traumatic Brain Injury. *Arch Neurol* 69: 1245-1251.
15. Lee B, Newberg A (2005) Neuroimaging in traumatic brain imaging. *NeuroRx* 2: 372-383.
16. Crane PK, Gibbons LE, Dams-O'Connor K, Trittschuh E, Sonnen J, et al. (2016) Association of traumatic brain injury with late-life neurodegenerative conditions and neuropathologic findings. *JAMA Neurol* 73(9): 1062-1069.
17. Dams-O'Connor K, Gibbons LE, Bowen JD, McCurry SM, Larson EB et al. (2013) Crane PK. Risk for late-life re-injury, dementia and death among individuals with traumatic brain injury: A population-based study. *J Neurol Neurosurg Psychiatry* 84(2): 177-182.
18. Faul M, Xu L, Wald MW, Coronado VG (2010) Traumatic Brain Injury in the United States. Available online at: https://www.cdc.gov/traumaticbraininjury/pdf/blue_book.pdf
19. Savonenko AV, Melnikova T, Hiatt A, Tong L, Worley PF, et al. (2012) Alzheimer's therapeutics: Translation of preclinical science to clinical drug development. *Neuropsychopharmacology* 37(1): 261-277.
20. Caye A, Axelrud L, Salum GA (2018) Traumatic brain injury and dementia. *Lancet Psychiatry* 5(10): 782-783.
21. Barnes DE, Byers AL, Gardner RC, Seal KH, Boscardin WJ, et al. (2018) Association of mild traumatic brain injury with and without loss of consciousness with dementia in US military veterans. *JAMA Neurol* 75(9): 1055-1061.
22. McKee AC, Cantu RC, Nowinski CJ, Robert AS, Andrew EB, et al. (2009) Chronic traumatic encephalopathy in athletes: Progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol* 68: 709-735.
23. Hyder AA, Wunderlich CA, Puvanachandra P, Gururaj G, Kobusingye OC (2007) The impact of traumatic brain injuries: A global perspective. *Neuro-Rehabilitation* 22: 341-353.
24. Langlois JA, Rutland-Brown W, Wald MM (2006) The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil* 21: 375-378.
25. Papa L, Edwards D, Ramia M (2015) Exploring Serum Biomarkers for Mild Traumatic Brain Injury. CRC Press.
26. Holm L, Cassidy JD, Carroll LJ, Borg J (2005) Summary of the WHO Collaborating Centre for Neurotrauma Task Force on Mild Traumatic Brain Injury. *J Rehabil Med* 37: 137-141.
27. Johnson VE, Stewart W, Smith DH (2013) Axonal pathology in traumatic brain injury. *Exp. Neurol* 246: 35-43.
28. Leker RR, Shohami E (2002) Cerebral ischemia and trauma-different etiologies yet similar mechanisms: Neuroprotective opportunities. *Brain Res Brain Res Rev* 39: 55-73.
29. Morganti-Kossmann MC, Rancan M, Stahel PF, Kossmann T (2002) Inflammatory response in acute traumatic brain injury: A double-edged sword. *Curr Opin Crit Care* 8: 101-105.
30. McKee AC, Robert CC, Nowinski CJ, Budson AE, Veronica ES, et al. (2010) Chronic traumatic encephalopathy in athletes: Progressive tauopathy following repetitive head injury. *J Neuropathol Exp Neurol* 68(7): 709-735.
31. Bramlett HM, Dietrich WD (2002) Quantitative structural changes in white and gray matter 1 year following traumatic brain injury in rats. *Acta Neuropathol* 103: 607-614.
32. Iwata A, Chen XH, McIntosh TK, Browne KD, Smith DH (2002) Long-term accumulation of amyloid-beta in axons following brain trauma without persistent upregulation of amyloid precursor protein genes. *J Neuropathol Exp Neurol* 61: 1056-1068.
33. Uryu K, Laurer H, McIntosh T, Domemico P, Susan L, et al. (2002) Repetitive mild brain trauma accelerates Abeta deposition, lipid peroxidation, and cognitive impairment in a transgenic mouse model of Alzheimer amyloidosis. *J Neurosci* 22: 446-454.
34. Buckner RL, Sepulcre J, Talukdar T, Hesheng L, Hedden T, et al. (2009) Cortical hubs revealed by intrinsic functional connectivity: Mapping, assessment of stability, and relation to Alzheimer's disease. *J Neurosci* 29: 1860-1873.