BioMed Research Journal

BMRJ, 3(S1): 09 www.scitcentral.com



Abstract: Open Access

Exploration of Cervical-Squamous-Epithelial-Cell (CSEC) Biometrics from Thin-Prep Pap Smear on GA Genotype of *Fas-Promoter-670* Gene in Indonesian Women Living with High-Risk HPV and *Candida* sp. Infection: A Case Report

Mauritius Lambertus Edy Parwanto^{1*}, Raditya Wratsangka¹, Assangga Guyansyah¹, Kirana Anggareni¹, Reza Aditya Digambiro¹, David David¹, Hanslavina Arkeman¹, Hosea Jaya Edy², Yosua Jaya Edy^{3,4}, Sisca Sisca¹ and Haryo Ganeca Widyatama¹

*1 Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Study Program of Pharmacy, Faculty of Math and Natural Sciences, Universitas Sam Ratulangi, Manado, Indonesia

³Center for Development of Civil Servants, State Personnel Board, The Republic of Indonesia, Bogor, Indonesia.

⁴Department of Sociology, Faculty of Social Politic, Universitas Kristen Indonesia, Jakarta, Indonesia.

Published November 01, 2019

ABSTRACT

Background: Human Papilloma Virus (HPV) infection can change Cervical-Squamous-Epithelial Cells (CSEC) to be abnormal. Epigenetically, this change in the direction of abnormality in CSEC is partly due to the polymorphism of the *Fas-promoter-670* gene. High-risk HPV patients can be infected by other microbes, for example *Candida* sp. The purpose of this study was to show the appearance of *Candida* sp. and CSEC based on epigenetics on the *Fas-promoter-670* gene polymorphism in Indonesian women with high-risk HPV infection. Biometric quantification is carried out with the following CSEC performance characteristics.

Case presentation: Indonesian women at the age of 28 years, living with high-risk HPV and Candida sp. infection, on April 21, 2016 had a Pap smear examination using a thin-prep method and also blood drawn from the cubital vein (vena cubiti). Blood and thin-prep samples were detected of Fas-promoter-670 gene polymorphisms by Restriction Fragment Length Polymorphism (RFLP) with BstN1 enzyme. Epigenetically, the subjects in this case had the GA genotype of Fas-promoter-670 gene in both blood and thin-prep samples. Candida sp. infection in patients is divided into early and advanced stages. Candida sp. infection in the early stages with characteristic by appearance of polymorphonuclear leukocytes on the hyfa, whereas at an advanced stage without polymorphonuclear leukocytes on the hyfa. CSEC performance (n cells=27) as follow: Cell Length (CL)=5.38 ± 0.77 μm (CI 95%=5.10-5.67 μm), Cell Width (CW)=4.37 ± 0.57 μm (95% CI=4, 16-4.58 µm); Cell Area (CA)= $20.29 \pm 4.46 \ \mu m^2$ (95% CI=18.62-22.01 μm^2); Cell Perimeter (CP)= $16.52 \pm 1.83 \ \mu m$ (95% CI=15.80-17.21 (μ m); Nucleus Length (NL)=0.81 ± 0.11 μ m (95% CI=0.77-0.86 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 ± 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 \pm 0.09 μ m (95% CI=95% CI 0.58-0.92 μ m); Nucleus Width (NW)=0.62 \pm 0.09 μ m); Nucleus Width (NW)=0.62 \pm 0.09 \mum); Nucleus Width (NW 0.65 μ m); Nucleus Area (NA)=0.41 \pm 0.10 μ m² (95% CI=0.38-0.45 μ m²); Nucleus Perimeter (NP)=2.38 \pm 0.27 μ m (95% CI=2.29-2.49) μ m); Nucleus Length Index (NLI)=15.43 ± 3.03 (95% CI=14.31-16.62); Nucleus Width Index (NWI)=14.46 ± 3.38 (95% CI=13.32-15.78); Nucleus Area Index (NAI)=2.16 ± 0.75 (95% CI=1.90-2.45) and the Nucleus Perimeter Index (NPI)=14.61 ± 2.29 (95% CI=13.77-15.49). **Conclusion:** Epigenetically, the subject in this case had the GA genotype of the *Fas-promoter-670* gene in blood and thin-prep samples. We found the early stages of *Candida* sp. infection which is characterized by appearance of polymorphonuclear leukocytes on the hyfa, whereas at an advanced stage without polymorphonuclear leukocytes on the hyfa. Base on the result of biometrics characteristic such as CL, CW, CA, CP, NL, NW, NA, NP, NLI, NWI, NAI and the NPI, CSEC in this case is a normal. Our team recommends a re-examination after 6 months.

Keywords: Epigenetically, Fas promoter-670 gene, Candida sp., Cervical-squamous-epithelial-cells, Cell biometrics

Corresponding author: Mauritius Lambertus Edy Parwanto, Department Food Science, College of Agriculture, Tikrit University, Iraq, E-mail: edyparwanto@trisakti.ac.id

Citation: Parwanto MLE, Wratsangka R, Guyansyah A, Anggareni K, Digambiro RA, et al. (2019) Exploration of Cervical-Squamous-Epithelial-Cell (CSEC) Biometrics from Thin-Prep Pap Smear on GA Genotype of *Fas-Promoter-670* Gene in Indonesian Women Living with High-Risk HPV and *Candida* sp. Infection: A Case Report. BioMed Res J, 3(S1): 09.

Copyright: ©2019 Parwanto MLE, Wratsangka R, Guyansyah A, Anggareni K, Digambiro RA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.