

Sudden Blindness in Children Passing Round Worm per Oral

Avinash Shankar^{1*}, Amresh Shankar², Anuradha Shankar³ and Shubham⁴

¹Institute of Applied Medicine, National Institute of Health and Research, Warisaliganj (Nawada), Bihar, India

²Aarogyam Punarjeevan, Ram Bhawan, Patna, India

³Centre for Indigenous Medicine and Research, Regional Institute of Ayurveda Research, Itanagar, Arunachal, India

⁴Department of Pediatrics, GMC, Thiruvananthapuram, Kerala, India.

Received October 28, 2019; Accepted November 05, 2019; Published August 07, 2020

ABSTRACT

Ascaris lumbricoides infestation is the most prevalent parasitic infection among the children in tropical and developing country but incidence of sudden blindness on passing the worm per oral is undocumented, the lag period depend on the prodromes and varies from 5 days, investigation reveal mere raised eosinophilic count and decreased hemoglobin with normal CT scan and CSF examination.

Materials: 10 cases of sudden blindness investigated and treated at various centre without any positive response attended our centre after 30-45 days of incidence, during January 2018 to March 2019 were selected.

Methods: Selected patient's parent were interrogated for the course of disease, treatment taken and their response, patients were clinically examined, investigated for basic bio parameters, vision and were treated with the prescribed regime containing pyridoxin, methyl cobalamin, nicotinamide, pantothenic acid and herbal neurovitalizer composite NEUROVIT.

Results: All patients had progressive vision gain and attended complete vision on 6 months therapy without any adversity and residual effect or any alteration in hepato-renal profile.

Conclusion: Sudden blindness in children after passing round worm or with history of round worm must be suspected for photoreceptor blockade by roundworm toxin and be treated with pyridoxin and herbal neurovitaliser to assure complete recovery.

Keywords: *Ascaris lumbricoides*, CT scan, CSF, Photo receptor, Neurovitaliser, Recovery

INTRODUCTION

Prevalence of intestinal worm infection is 49.35% and *Ascaris lumbricoides* is most common parasitic infection 46.85% soil transmitted Helminth infection form the most important group of intestinal worm affecting 2 billion people world-wide causing considerable morbidity. *Ascaris lumbricoides* remain the most prevalent parasitic infection, i.e., 75% despite of therapeutic response of Albendazole and Mebendazole, but eradication is difficult due to recurrent infection. Considering the changing effect of worm infestation GOI has launched a program to combat the worm infestation, i.e., National deworming day for children of age group 1-19 years biannually. As per WHO >836 million children are at risk of parasitic manifestation worldwide and 214 million children are of age group 1-14 years [1-14]. In addition evidence of disproportionate worm infestation [15] and self-drug use resulting resistance to available deworming agent and presently a combination of parasiticide, i.e., Albendazole and Ivermectine [16] is in quite consideration. As these agents only act on adult worm

not on cyst or ova its recurrent dose must be prescribed as on 45th day every ova is developed to active adult round worm.

MATERIALS AND METHODS

Materials

10 children attending the centre for critical care with complaints of sudden blindness after passing round worm per oral having treated at various hospitals without any positive response and were suggested brain surgery,

Corresponding author: Dr. Avinash Shankar, Chairman, Institute of Applied Medicine, National Institute of Health and Research, Warisaliganj (Nawada), Bihar, 805130, India, E-mail: dravinashshankar@gmail.com

Citation: Shankar A, Shankar A, Shankar A & Shubham. (2020) Sudden Blindness in Children Passing Round Worm per Oral. J Neurosurg Imaging Techniques, 5(2): 262-270.

Copyright: ©2020 Shankar A, Shankar A, Shankar A & Shubham. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ophthalmological examination, CT brain shows no evident pathology except blood showing high eosinophilic count.

Patients	Age/Sex	Clinical presentation	Lag period
A	10/F	History of passing round worm per oral, lose motion, vomiting fever, loss of vision both side	3 days
B	9/M	Lose motion, vomiting, pain in abdomen, itching in the body, loss of vision both side passage of round worm per oral	2 days
C	12/M	Vomiting, pain in abdomen, fever urticarial rash over the extremity, passage of round worm per oral, loss of vision both side	4 days
D	6/M	Vomiting, lose motion, involuntary body movement, urticarial rash, passing round worm per oral sudden blindness	2 days
E	13/F	Lose motion, urticarial rash, fever, shivering, nausea, pain in abdomen, passage of round worm per oral, sudden blindness	3 days
F	12/F	Lose motion of white color, dark urine, intense itching with rash, fever, vomit of round worm, abdominal pain, loss of vision	5 days
G	8/F	Agonising pain in abdomen, lose motion, vomiting, urticarial rash, passage of round worm per oral, loss of vision	4 days
H	8/M	Nausea, vomiting, urticarial rash, passage of round worm per oral, loss of vision	5 days
I	6/F	Vomiting, loss of appetite, urticarial rash, passage of round worm, loss of vision	4 days
J	14/F	Fever, pain in abdomen, vomiting, headache, urticarial rash, passage of round worm both from mouth and stool	1 day

Method

All the patients presenting with sudden blindness and associated history of passing round worm per oral and treated at various hospitals without any vision improvement in spite of medication and no pathology were detected on various investigation like CT brain, retinal examination and

various hematological examination, were interrogated examined thoroughly, investigated for basic hematological, hepatic and renal profile.

All the selected patients were administered the following irrespective of age and presentation:

I.V Mannitol 10% with glycerine. 10% in therapeutic dose.

Inj Methyl cobalamin, Nicotinamide, Pyridoxin and Pantothenic acid with betamethasone. 1 ml intravenous every 4th day very slow.

Syr Herbal neuro energiser (NEUROVIT) 2.5 ml-5 ml every 8 h.

Susp Albendazol 400 mg plus Ivermectin 3 mg at bed time for 5 days.

Bland and simple high carbohydrate diet.

Herbal neurovitaliser NEUROVIT constitutes:

Each 5 ml	
<i>Acorus calamus</i>	100 mg
<i>Herpestis monnieri</i>	100 mg
<i>Convolvulus pluricaulis</i>	100 mg
<i>Nardostachys jatamansi</i>	100 mg
<i>Cassia angustifolia</i>	100 mg

Patients parent were instructed to practice daily to ascertain visual response by finger counting or light reflex, in addition also suggested to mark any adversity or new emerging manifestation if any and report immediately.

Patients were routinely examined on every week to ascertain response of the therapy and safety profile. At the end of therapy when patient ensured their complete vision patients were examined by ophthalmologist for vision and visual acuity.

OBSERVATIONS

Selected patients were of age group 6-14 years (**Table 1**) and among them 04 was male and 06 female (**Figure 1**),

though they approached for Medicare within the lag period of 3-5 days at appropriate centre, investigated for CT scan, ophthalmological examination to asses vision and retina status which remain within normal limit in all cases, except raised eosinophilic count (**Table 2**). Patients were treated with many neurotropics and topical eye drops without any positive response. Majority patients attended our centre after 30-45 days of the onset of blindness and lag period of onset of blindness and passing the round worm per oral was 1-5 days while patients presenting with associated CNS manifestation like involuntary movement and headache has very short lag period, i.e., 1 or 2 days. At our centre hematological examination show raised eosinophil count with other normal other parameters, i.e., hepatic and renal.

Table 1. Showing distribution of patients.

Age group (in years)	Number of patients		
	Male	Female	Total
6-8	02	02	04
8-10	01	01	02
10-12	-	-	-
12-14	01	03	04




Figure 1. Pie diagram showing sex composition of the patients.

Table 2. Presentation of the patients.

Sudden blindness
Passing worm per oral
Nausea and vomiting
Fever
Lag period of onset of blindness and passing the worm per oral: 24-72 h
Blurring of vision
Sign of avitaminosis /xerosis /bitots spot

RESULTS

All patients started visual improvement by 8th day of therapy and complete visual recovery by 6th month of therapy without any visual debility; optometry confirmed the vision

in all patients as 6/6 in both eyes (**Tables 3 and 4 and Figure 2**). No adversity or sequel is noted in any case or any evident of post therapy withdrawal affect, i.e., decline in vision or visual acuity or any CNS manifestation.

Table 3. Showing bio parameter status at various states.


Basic bio parameters	Number of Patients		
	(A)	(B)	(C)
Hematological			
Absolute eosinophil count			
<200/cc	-	-	8
200-300/cc	3	4	2
300-400/cc	6	6	0
400-500/cc	1	0	0
TLC			
<6000/cc	1	0	0
6000-7000/cc	8	10	10
>7000/cc	1	0	0
Hemoglobin parent			
<10 g%	5	3	0
>10 g%	5	7	10
Diabetic profile			
Blood sugar			
Fasting:			
<100 mg%	9	10	10
>100 mg%	1	0	0
Post prandial:			
<150 mg%	10	10	10
>150 mg%	0	0	0

Hepatic profile			
SGOT			
<30 IU	7	8	10
>30 IU	3	2	0
SGPT			
<30 IU	7	8	10
>30 IU	3	2	0
Alkaline phosphatase			
<140 mg%	10	10	10
>140 mg%	0	0	0
Renal profile			
Blood urea			
<26 mg%	10	10	10
>26 mg%	0	0	0
Serum creatin			
<1.5 mg%	10	10	10
>1.5 mg%	0	0	0
CT scan			
Altered	None	None	None
Unaltered	10	10	10
Vision			
Status of eye	Normal	Normal	Normal
PL			
Absent	10	10	0
Present	0	0	10
PR			
Absent	2	0	0
Present	8	10	10
Vision	Absent	Absent	Normal

A – At first centre of treatment; B – At our centre on admission; C – On completion of treatment

Table 4. Outcome of the study.

Particulars	Number of patients
Perception of light	10
Finger counting	10
Blurred vision	None
Clear distant vision	10
Clear distant vision	10
Completely normal vision	10
Safety profile	
Hematological	Improved in all
Hepatic profile	Normal in all
Renal profile	Normal in all

**Figure 2.** Round worm kinetics in intestine.

DISCUSSION

Round worm infestation is very common but manifestation like blindness after passing the worm per oral is very uncommon or remain unmarked, in addition variable lag

period of onset of blindness and worm passage, i.e., 1-5 days suggest its dependence on prodromes, those who had CNS prodromes like headache and involuntary movement had earlier onset [17-22]. Patients presentation on passing worm per oral suggest worm irritation leading to release of a

polypeptide ASCARON which stimulate the intestinal mucosal nerve endings resulting in nausea, vomiting and loose motion, absorption of toxin in blood causes anaphylactic reaction resulting in fever and urticarial rash while access to CSF results in neurosuppression due to inhibition of neurotransmitter GABA as a result of inhibition

of coenzyme pyridoxal phosphatase enzyme by the toxin [23,24]. Sudden blindness is due to effect of neuroconduction suppression results in blockade of neurotransmission from photo receptor of retinal fovea (**Figures 3 and 4**) [25,26]. No change in bio parameters are observed in any case and eosinophil count came to normal.




Figure 3. Schematic presentation of pathogenesis of sudden blindness.




Figure 4. Showing pattern of vision improvement.

A: Perception of light; B: Finger counting; C: Blurred vision; D: Clear vision

All patients recovered of blindness having progressive vision gain from perception of light to normal vision in 6 months duration with the treatment is attributed to:

- Intravenous mannitol 10% with glycerine 10% relieved neural edema.
- Supplementation of pyridoxin as injection of methyl cobalamin, pyridoxin, nicotinamide and pantothenic acid competitively inhibit polypeptide and activate pyridoxal phosphatase and ensure increased neurotransmitter GABA, methyl cobalamin and pantothenic acid promote neuro conduction.
- Herbal composite NEUROVIT constituents ensure neurovitalization and photoreceptor activation [27].
- Administration of albendazole plus ivermectin ensures worm eradication.
- Nutritious diet support recovery.

CONCLUSION

Sudden blindness after passing round worm or without patient must be duly taken care suspecting *Ascaris* toxin as a factor and duly treatment will ensure cure and safety from undue expenses especially in tropical countries where round worm infestation is very common. Herbal composite and pyridoxine supplementation proves boon for cure.

REFERENCES

1. Seltzer E (1999) Ascariasis: Tropical infectious disease - Principles, Pathogen and Practice. 1st Edn. Guerrant RL, Weller PF (Eds), Philadelphia: Churchill Livingstone, p: 553.
2. Salam N, Azam S (2017) Prevalence and distribution of soil-transmitted helminth infections in India. BMC Public Health 17: 201.
3. WHO (2019) Prevalence of soil transmitted helminths.
4. Lobo DA, Velayudhan R, Chatterjee P, Kohli H, Hotez PJ (2011) The neglected tropical diseases of India and South Asia: Review of their prevalence, distribution and control or elimination. PLoS Negl Trop Dis 5: e1222.
5. Padmaja N, Swaroop PS, Nageswararao P (2014) Prevalence of intestinal parasitic infections among school children in and around Amalapuram. J Public Health Med Res 2: 36-38.
6. Panda S, Rao UD, Sankaram KR (2012) Prevalence of intestinal parasitic infections among school children in rural area of Vizianagaram. IOSR J Pharm Biol Sci 3: 42-45.
7. Ragunathan L, Kalivaradhan SK, Ramadass S, Nagaraj M, Ramesh K (2010) Helminthic infections in school children in Puducherry, South India. J Microbiol Immunol Infect 43: 228-232.
8. Golia S, Sangeetha K, Vasudha C (2012) Prevalence of parasitic infections among primary school children in Bangalore. Int J Basic Appl Med Sci 4: 12-18.
9. Krishnan A, Sekar U, Sathanantham DK (2013) Prevalence and pattern of helminthic infection among children in a primary school of rural Tamil Nadu. Acad Med J India 1: 40-42.
10. Fernandez MC, Verghese S, Bhuvaneswari R, Elizabeth SJ, Mathew T, et al. (2002) A comparative study of the intestinal parasites prevalent among children living in rural and urban settings in and around Chennai. J Commun Dis 34: 35-39.
11. Dhanabal J, Selvadoss PP, Muthuswamy K (2014) Comparative study of the prevalence of intestinal parasites in low socio-economic areas from south Chennai, India. J Parasitol Res 630968.
12. Sunish I, Rajendran R, Munirathinam A, Kalimuthu M, Kumar VA, et al. (2015) Impact on prevalence of intestinal helminth infection in school children administered with seven annual rounds of diethyl carbamazine (DEC) with albendazole. Indian J Med Res 141: 330-339.
13. Clark A, Turner T, Dorothy KP, Goutham J, Kalavati C, et al. (2003) Health hazards due to pollution of waters along the coast of Visakhapatnam, east coast of India. Ecotoxicol Environ Saf 56: 390-397.
14. Nikolay B, Brooker SJ, Pullan RL (2014) Sensitivity of diagnostic tests for human soil-transmitted helminth infections: A meta-analysis in the absence of a true gold standard. Int J Parasitol 44: 765-774.
15. Shankar A (2008) Disproportionate worm infestation, a cause of anthelmintic default. IJM Today 03: 31-32.
16. Shankar A (2009) Albendazole and ivermectin in management of helminthiasis. IJM Today 04: 50-52.
17. Wani ML, Ashraf HZ, Ahangar AG, Wani SN, Nayeem UH, et al. (2012) Unusual presentation of *Ascaris lumbricoides*. J Clin Case Rep 2: 174.
18. Coursin DB (1969) Vitamin B6 and brain function in animals and man. Ann N Y Acad Sci 166: 7-15.
19. Gale K (1985) Mechanisms of seizure control mediated by γ -aminobutyric acid: Role of the substantia nigra. Fed Proc 44: 2414-2424.
20. Kaufman DL, Houser CR, Tobin AJ (1991) Two forms of the gamma-aminobutyric acid synthetic enzyme glutamate decarboxylase have distinct intraneuronal distributions and cofactor interactions. J Neurochem 56: 720-723.

21. Martin DL, Martin SB, Wu SJ, Espina N (1991) Cofactor interactions and the regulation of glutamate decarboxylase activity. *Neurochem Res* 16: 243-249.
22. Martin DL, Rimvall K (1993) Regulation of γ -aminobutyric acid synthesis in the brain. *J Neurochem* 60: 395-407.
23. Miller LP, Walters JR, Martin DL (1977) Post-mortem changes implicate adenine nucleotides and pyridoxal-5'-phosphate in regulation of brain glutamate decarboxylase. *Nature* 266: 847-848.
24. Tews JK (1969) Pyridoxine deficiency and brain amino acids. *Ann N Y Acad Sci* 166: 74-82.
25. (2016) How the human eye sees? WebMD. Ed. Alan Kozarsky. WebMD.
26. Than K (2016) How the human eye works? LiveScience. TechMedia Network.
27. Shankar A (2019) Herbal composite constituents. *Pharmacological Basis of Indigenous Therapeutics*, Bhalani Publication: Mumbai.