

Prevalence of Anal Cancer in Patients Presenting with Different Proctological Issues: A 9 years' Experience (2010-2019)

Anastasios G Xiarchos¹, Fernand Tshijanu^{1*}, Evanthia D Chatzigianni², Michael N Vorias³, George A Xiarchos⁴ and Luc Mubaminyi⁵

¹Department of Surgery, Athens Medical Center-Peristeri Clinic, Athens, Greece

²Department of Anaesthesiology, Athens Medical Center-Peristeri Clinic, Athens, Greece

³Department of Surgery, Mediteraneo Hospital-Glyfada, Athens, Greece

⁴Medical School of European University of Cyprus, Cyprus

⁵Medical School of National Kapodistrian University of Athens-Greece, Athens, Greece.

Received November 07, 2020; Revised December 04, 2020; Accepted February 18, 2021

ABSTRACT

From epidemiological perspective, anal cancer is a relatively uncommon condition, comprising respectively 1,5 % and 1-8% of all digestive System and anorectal malignancies. Traditionally this situation had been regarded as a nosologically entity of older women in 7th decade of life, with a sex ratio of 5 females vs 1 male [1,9 and 17]. These articles peruse is to determine the rate of anal malignancies occurrence among different proctological issues treated in our Surgical Department, in the times interval of 2010 to 2019. Thus, we enrolled in toto 29569 patients, 15115 (51 %) females and 14454 (48,8%) males, the median age was 18-70 years, 35 patients (0,11 %) were diagnosed with anal cancer, furthermore carcinoma in situ (Bowen's disease) was the most prevalent (54 %) followed respectively by squamous cell carcinoma (31 %) and adenocarcinoma (9%). In hemorrhoids, group anal cancer was seen in 11 patients (0,09%) with an occurrence frequency of 1 in 1000 patients, in the group of anal fissures 2 patients (0,03%) had anal cancer with a frequency of 1 in 3000 patients. Anorectal fistula and perianal abscess were not associated with anal cancer. Anal HPV infection was more associated with anal cancer in the rate of 3,07% or 21 patients with an occurrence frequency of 1 in 30 patients. Pilonidal sinus was associated with anal cancer in 0,02% or 1 patient. In toto, the occurrence frequency of anal cancer in proctological issues is 1 in 1000 patients.

Keywords: Proctological issues, Anal cancer, Prevalence

INTRODUCTION

Despite its increased incidence in both genders over the last decades, anal cancer remains relatively a rare event presenting 1,5 % of all digestive system malignancies and 1 to 8 % of all anorectal malignancies. Its sex ratio remains standard with 5 females vs 1 male. Traditionally, anal cancer used to involve older women especially from the 7th decade of age. Unfortunately, in this new generation, the increased prevalence of this neoplastic disease is directly correlated to different factors such as, receptive anal intercourse, immunodepression of different etiologies (HIV, corticosteroid use...), HPV infection especially serotypes 16,18, and smoking [1,2,9]. Most of anal cancer mimic benign proctological conditions regarding symptomatology features such as haemorrhoids, anal fissures, anal fistula, pruritus anis [10,11]. In this article, we aim to determine the occurrence of anal cancer in the group of patients presenting with any proctologic issue in our surgical Department during a times interval of 9 years as mentioned in the title.

RESULTS

Statistical analysis

29569 patients with different proctologic issues were enrolled, 15115 (51%) females and 14454 (48,8%) males, anal cancer was diagnosed in 35 patients (0,11 %), carcinoma in situ (Bowen's disease) was more prevalent (54 %) followed respectively by squamous cell carcinoma (31%)

Corresponding author: Fernand Tshijanu, Department of Surgery, Athens Medical Center-Peristeri Clinic, Athens, Greece, Tel: (30)6996154491; E-mail: tshijanufernand@yahoo.fr

Citation: Xiarchos AG, Tshijanu F, Chatzigianni ED, Vorias MN, Xiarchos GA, et al. (2021) Prevalence of Anal Cancer in Patients Presenting with Different Proctological Issues: A 9 years' experience (2010-2019). Int J Surg Invasive Procedures, 4(2): 184-188.

Copyright: ©2021 Xiarchos AG, Tshijanu F, Chatzigianni ED, Vorias MN, Xiarchos GA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and adenocarcinoma (9%) (Tables 1 and 2 and Figure 1). In the group of haemorrhoids, anal cancer was seen in 11 patients (0,09%) with an occurrence frequency of 1 in 1000 patients, anal fissure was associated with anal cancer in the range of 2 patients (0,03%) with a frequency of 1 in 3000 patients. Anorectal fistula and perianal abscess were not associated with anal cancer. Occurrence anal HPV infection

(Condyloma) was more associated with anal cancer in the rate of 3,07 % or 21 patients with an occurrence frequency of 1 in 30 patients. Pilonidal sinus was associated with anal cancer in 0,02% or 1 patient. In toto, the occurrence frequency of anal cancer in proctologic issues is 1 in 1000 patients (Table 3 and Figure 1).

Table 1. Patients Characteristics.

Proctology issue	Patient	Median age	Gender		Pathology	
			Female	Male	Benign	Anal cancer
Hemorrhoids	11780	50-60	9650	2130	11769	11
Anal fissure	6320	60-70	2200	4120	6318	2
Anorectal Fistula	4102	60-70	1271	2831	4102	0
Perianal abscess	1870	70-80	758	1112	1870	0
Anal condyloma	682	40-50	253	429	661	21
Pilonidal sinus	4815	18-40	983	3832	4814	1
Total	29569		15115	14454	29534	35

Table 2. Pathologic types of anal cancer.

ANAL CARCINOMA	
Carcinoma in situ (BOWNS disease): 19	Adenocarcinoma: 3
Squamous cells: 11	
Skins cancer: Basal cells: 2	
Melanoma: 0	

Table 3. Rate of anal cancer occurrence out of proctologic issues.

Proctology issue	Percentage of cancer	Frequency
Hemorrhoids	0.09%	1 in 1000 patients
Anal fissure	0.03%	1 in 3000 patients
Anorectal Fistula	0%	0
Perianal abscess	0%	0
Anal condyloma	3.07%	1 in 30 patients
Pilonidal sinus	0.02%	1 in 5000 patients
Total	0.12%	1 in 1000 patients




Figure 1. Pathologic types of anal cancer.




Figure 2. Rate of anal cancer in patients with proctological issues.

DISCUSSION

From anatomic perspective we mean by anal canal, the segment between the dentate line and the anal verge. Moreover, the surgical view considers the anal canal, as the area from the palpable anorectal ring to the peri-anal skin. This latter definition is the most widely accepted in practice and the most preferred by the American Joint Committee on Cancer (AJCC). Therefore, any tumour involving the

anorectal junction is stated as a rectal cancer in this case, the topography epicentre is more than 2 cm proximal to dentate line. In addition to, anal cancer is defined if the epicentre is 2 cm or less from the dentate line. Taxonomically, tumours of the peri-anal skin are designated as perianal cancers and resemble biologically to other skin tumours [72,13,30]. This contrast is crucial since anal cancers are more aggressive than cancers of the peri-anal skin, furthermore its incidence

is up to 5 times more common than that of anal margin lesions [12,13,16,30].

Consistently, anal cancer was regarded as a disease of older women, with onset in the 7th decade of life. This neoplastic entity used to be of a very rare occurrence. However, over the past decades its incidence has raised especially in women and in homosexual men. Additionally, for the past decade, its escalated incidence has redheaded the range of 2,2 % annually. Furthermore, this cancer represented 0,5 % of all cancers diagnosed in the USA [1,9,17-19]. Etiologically, the nowadays occurrence of anal cancer is more related to Human Papilloma Virus infection in 91 % so that the frequency of this malignancy is merged in younger people of both genders affected by this virus. Therefore, since the probability of developing this cancer is 120-fold when there is a pre-existing HPV ano-rectal infection, anal cancer in young people can be regarded as a sexually transmitted neoplastic disease which needs a prophylactic measure involving a sexual education in puberty and adolescent age groups as well as, a vaccination against this infectious agent. And the most oncogenic serotypes are HPV 16 and 18. thus, the biopsy is recommended once there is ano-rectal HPV infection especially when there is a coexistent HIV infection [2-8,14,26-28]. Results of peer-reviewed studies of anal cancer, including detection of HPV DNA, demonstrated that the prevalence of HPV16 and 18 in 72 % of patients, and includes screening for vaginal and cervical cancer and male external genitalia. Anal cancer has numerous risk factors apart from HPV such as a non-protected sexual life in term of multiple partners, anal receptive intercourse in both genders, history of vulvar or vaginal and cervical dysplasia or cancer, advanced age, smoking and immunosuppression related to different organs such as HIV infection or solid [15,16,20,22-25].

A study using data from the Danish Cancer Registry demonstrated that the probability of developing anal cancer after a diagnosis of cervical cancer or cervical intraepithelial neoplasia was three to five times as high as the probability of developing stomach or colon cancer with a strong association between cervical cancer and anal cancer. Frisch et al. have found an association between anal cancer and lymphoma or leukaemia and raises a possible role of immunodeficiency in the development of anal cancer [29].

Regarding pathologic perspective, 85 % of anal cancers are of squamous type [21,29] but in us study the most prevalent histologic type was carcinoma *in situ* (Bowen disease) with 54% followed squamous cell carcinoma with 31 % and adenocarcinoma with 9 %, skin cancer (basal cell) 6%. There was no case of anal melanoma in our study.

CONCLUSION

Anal cancer remains a nosologic entity of rare occurrence even among different proctologic issues. Its incidence is nowadays increasing because of the epidemiologic profile of

HIV infection in both genders, especially in young population. We suggest a good sexual education in adolescent population as well as vaccination against this etiological as well as a screening ano-rectal biopsy in women with history of vulvo-vaginal HPV infection or cancer and in general in high risk patients such as immunocompromised patients from different origins.

REFERENCES

1. Nelson VM, Benson AB (2017) "Epidemiology of Anal Canal Cancer". Surg Oncol Clin North Am 26 (1): 9-15.
2. Anal Cancer Prevention (2014) National Cancer Institute. Accessed on: 30 May 2019.
3. National Cancer Institute (2009) Anal Cancer Treatment (PDQ) Patient Version. Archived July 14, 2009, at the Way back Machine 13 June 2008. Accessed on: 26 June 2009.
4. Margaret AS, David MW, Jane SC, Goon, Peter KC (2012) HPV infection, anal intra-epithelial neoplasia (AIN) and anal cancer: Current issues. BMC Cancer 12 (1): 398.
5. Frisch M (2002). "On the etiology of anal squamous carcinoma". Danish Med Bullet 49 (3): 194-209.
6. Frisch M, Glimelius B, van den Brule AJ, et al. (1997) Sexually transmitted infection as a cause of anal cancer. N Engl J Med 337 (19): 1350-1358.
7. Parkin DM (2006) The global health burden of infection-associated cancers in the year 2002. Int J Cancer 118 (12): 3030-3044.
8. Fred Hutchinson Cancer Research Centre (2007) Changing trends in sexual behaviour may explain rising incidence of anal cancer among American men and women. Fred Hutchinson Cancer Research Centre (fhcrc.org).
9. "Anal Cancer" American Cancer Society (2014).
10. Esiashvili N, Landry J, Matthews RH (2007) "Carcinoma of the Anus Management". Armenian Health Network, Health.
11. Saleem AM, Paulus JK, Shapter AP, Baxter NN, Roberts PL, et al. (2011) "Risk of anal cancer in a cohort with human papillomavirus-related gynaecologic neoplasm". Obstet Gynaecol 117 (3): 643-649.
12. American Joint Committee on Cancer (2018) AJCC Cancer Staging Form Supplement, AJCC Cancer Staging Manual, Eighth Edition, Last updated 05 June 2018.

13. MB Amin, SB Edge, FL Greene, et al. (2017) AJCC Cancer Staging Manual. 8th edition, New York: Springer.
14. Tuller, David (2007-01-31) "HPV vaccine may help to prevent anal cancer". International Herald Tribune.
15. Agency for Healthcare Research and Quality (2008) "Hospital HIV Clinic Offers Convenient, Proactive Screening for Anal Cancer, Enabling Identification and Treatment of Precancerous Lesions".
16. National Comprehensive Cancer Network (2013) "NCCN Clinical Practice Guidelines in Oncology: Anal Carcinoma. V 1".
17. Cancer Research (2008) "Detailed Guide: Anal Cancer What Are the Key Statistics About Anal Cancer?"
18. Cancer.net (2019) "Anal Cancer: Statistics" Cancer.net.
19. Cancer.net (2014) "Anal Cancer Statistics" Cancer Research UK.
20. Tong WW, Hillman RJ, Kelleher AD, Grulich AE, Carr A (2014) Anal intraepithelial neoplasia and squamous cell carcinoma in HIV-infected adults. HIV Med 15: 65-76.
21. Flejou JF (2015) An update on anal neoplasia. Histopathology 66:147-160.
22. Berry JM, Jay N, Cranston RD, Darragh TM, Holly EA, et al. (2014) Progression of anal high-grade squamous intraepithelial lesions to invasive anal cancer among HIV-infected men who have sex with men. Int J Cancer 134: 1147-1155.
23. Nagle D (2009) Anal squamous cell carcinoma in the HIV-positive patient. Clin Colon Rectal Surg 22: 102-106.
24. Ouhoumane N, Steben M, Coutlee F, Vuong T, Forest P, Rodier C, et al. (2013) Squamous anal cancer: Patient characteristics and HPV type distribution. Cancer Epidemiol 37: 807-812.
25. Mendez-Martinez R, Rivera-Martinez NE, Crabtree-Ramirez B, Sierra-Madero JG, Caro Vega Y, et al. (2014) Multiple human papillomavirus infections are highly prevalent in the anal canal of human immunodeficiency virus-positive men who have sex with men. BMC Infect Dis 14: 671.
26. Garcia-Espinosa B, Moro-Rodriguez E, Alvarez-Fernandez E (2013) Human papillomavirus genotypes in human immunodeficiency virus-positive patients with anal pathology in Madrid, Spain. Diagn Pathol 8: 204.
27. Cachay E, Agmas W, Mathews C (2015) Five-year cumulative incidence of invasive anal cancer among HIV-infected patients according to baseline anal cytology results: an inception cohort analysis. HIV Med 16: 191-195.
28. Wexler A, Berson AM, Goldstone SE, Waltzman R, Penzer J, Maisonet OG, et al. (2008) Invasive anal squamous-cell carcinoma in the HIV-positive patient: Outcome in the era of highly active antiretroviral therapy. Dis Colon Rectum 51: 73-81.
29. Abunassar M, Reinders J, Jonker DJ, Asmis T (2015) Review of anal cancer patients at the Ottawa hospital. Eur J Surg Oncol 41: 653-658.
30. Sinnatamby CS (1999) Last's anatomy: Regional and applied. 10th edition, Edinburgh: Churchill Livingstone.