

No Mother Should Die of Peripartum Cardiomyopathy

James D Fett^{1,2*}

¹Department of Internal Medicine, Hospital Albert Schweitzer, Deschapelles, Haiti.

²Department of Cardiology, University of Pittsburgh Medical Center, Pittsburgh, PA, USA.

Received December 06, 2018; Accepted December 17, 2018; Published February 08, 2019

ABSTRACT

Peripartum cardiomyopathy (PPCM) is a unique form of heart failure characterized by systolic dysfunction developing during or shortly after pregnancy in a subject with a previously healthy heart; and is one of the leading causes of maternal mortality world-wide. Ventricular tachyarrhythmias with sudden cardiac arrest (SCA) and sudden cardiac death (SCD) do occur in peripartum cardiomyopathy subjects; but the occurrence is infrequent and most often only where there is severe systolic dysfunction with LVEF less than 0.35. Earlier diagnosis, with better-preserved LVEF, is the best way to prevent either SCA or SCD. Those with LVEF less than 0.35 can be protected by continuous cardiac rhythm monitoring with a wearable cardiac defibrillator (WCD) in the early stages; or with implantable cardiac defibrillator (ICD) if the LVEF does not rise above the level of 0.35 after 5 to 6 months. "Evidence-based" treatment utilizing beta-blockers (BB) and angiotensin converting enzyme inhibitors (ACEI) or angiotensin releasing hormone blockers (ARB) also increases the threshold, helping to prevent these malignant arrhythmias from occurring. Controlled studies are still needed to compare use vs. non-use of prolactin inhibition treatment. At this time, we are on the cusp of recognizing the mothers who are at greatest risk for the development of PPCM as well as those who, although apparently fully recovered, are still at greatest risk for the development of heart failure in a post-PPCM pregnancy. "State-of-the-art" knowledge about PPCM and its diagnosis/management obligates us all as medical caregivers, as well as all subjects in pregnancy, to make sure that any loss of life due to PPCM becomes rare or non-existent. No mother should die of PPCM. I foresee the day when that will become fact, as well as full recovery for almost all.

BRIEF CASE SUMMARY

24 year old gravida 2, para 2. First pregnancy uncomplicated, vaginal birth, healthy male newborn. Current pregnancy received regular prenatal care and experienced increasing discomfort during the last month of pregnancy. Attempted pitocin drip induction of labor at 40 weeks gestation intermittently over a 24 h period. Persistent tachycardia observed during pitocin infusion. Cardiology consultation requested, subsequent echocardiography revealed severe dilated cardiomyopathy, left ventricular ejection fraction (LVEF) 0.35, left ventricular internal diameter diastole (LViDd) 7.17 cm. Treatment initiated for heart failure with immediate Cesarean section under epidural anesthesia on 8 December, healthy female newborn. Experienced some improvement, with brief discharge from hospital, but required readmission in NYHA Class III heart failure. Repeat echocardiography on 9 December, LVEF 0.30; and again on 9 January, LVEF 0.15. Despite standard heart failure therapy, deterioration continued with development of cardiogenic shock and acute renal failure, died on 13 January.

COMMENTARY

Peripartum cardiomyopathy (PPCM) is a unique form of heart failure characterized by systolic dysfunction developing during or shortly after pregnancy in a subject with a previously healthy heart; and is one of the leading causes of maternal mortality world-wide [1,2]. Over the past two decades, maternal mortality rates have diminished from over 50% to less than 5%, with variation world-wide. With appropriate timely diagnosis and treatment, no mother should die of PPCM.

In the USA, the incidence of PPCM in mothers of African heritage is double that of those who do not have African heritage [3]. The reasons for this are not yet clear, but genetic factors, higher incidence of hypertension in

Corresponding author: James D Fett, MD, Department of Cardiology, University of Pittsburgh Medical Center, Pittsburgh, PA, USA, Tel: 360-438-5270, E-mail: fett.sprunger@comcast.net

Citation: Fett JD. (2019) No Mother Should Die of Peripartum Cardiomyopathy. J Cardiol Diagn Res, 2(1): 37-39.

Copyright: ©2019 Fett JD. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

pregnancy and later post-partum recognition of PPCM appear to play a role (**Figure 1**).




Figure 1. Comparative incidence of PPCM.

Ventricular tachyarrhythmias with sudden cardiac arrest (SCA) and sudden cardiac death (SCD) do occur in peripartum cardiomyopathy subjects; but the occurrence is infrequent and most often only where there is severe systolic dysfunction with LVEF less than 0.35 [4]. Earlier diagnosis, with better-preserved LVEF, is the best way to prevent either SCA or SCD. Those with LVEF less than 0.35 can be protected by continuous cardiac rhythm monitoring with a wearable cardiac defibrillator (WCD) in the early stages; or with implantable cardiac defibrillator (ICD) if the LVEF does not rise above the level of 0.35 after 5 to 6 months [1,5].

“Evidence-based” treatment utilizing beta-blockers (BB) and angiotensin converting enzyme inhibitors (ACEI) or angiotensin releasing hormone blockers (ARB) also increases the threshold, helping to prevent these malignant arrhythmias from occurring. Guidelines are provided in the American Heart Association/American College of Cardiology publication recommendations for the treatment of heart failure with reduced LVEF [6,7]. If the diagnosis of PPCM is made antenatal, it is safe to use hydralazine in place of ACEI/ARB. The use of hydralazine can be accompanied by the use of nitrates, depending upon possible heart rate increase from the hydralazine. BB treatment is safe to use during breastfeeding; but of the ACEI group, only enalapril is safe to use while breastfeeding, in which case ARBs should also be avoided. Likewise, the newer Entresto (valsartan/sacubitril) is not deemed safe to use while breastfeeding. “Controlled studies are still needed to compare use vs non-use of prolactin inhibition treatment.

Recent reports from the Peripartum Cardiomyopathy Network of North America, through the Investigations of Pregnancy-Associated Cardiomyopathy (IPAC) study, have clearly confirmed that adverse events, including chronic heart failure, following a diagnosis of PPCM, correlate with

diagnosis not made until there is already severe systolic dysfunction, $LVEF < 0.30-0.35$ [1].

Increasing awareness of PPCM enables earlier recognition, with a diagnosis made closer to the time of delivery, even antenatal. It appears that this earlier diagnosis is associated with better-preserved cardiac function, in which case there is better response to the effective treatment available, with fewer adverse events and fuller recoveries.

What other adverse events threaten the life of the new mother with PPCM? The most serious are major thromboembolic phenomena and severe heart failure transitioning to chronic cardiomyopathy requiring eventual cardiac transplantation; an option not available to many PPCM subjects in the “developing world” [8]. These adverse events can also be avoided if diagnosis can be made with relatively intact systolic function, i.e., $LVEF > 0.35$ at diagnosis [8,9].

At this time, we are on the cusp of recognizing the mothers who are at greatest risk for the development of PPCM as well as those who, although apparently fully recovered, are still at greatest risk for the development of heart failure in a post-PPCM pregnancy [10]. This recognition will come through continuing development of a “genetic risk profile” [11,12], a “hypertensive risk profile” [1] an “angiogenic risk profile” and a “contractile reserve risk profile” (**Figure 2**) [13].




Figure 2. Risk for relapse of heart failure in a post-PPCM pregnancy [14,15].

CR: Contractile Reserve on Exercise Stress Echo

“State-of-the-art” knowledge about PPCM and its diagnosis/management obligates us all as medical caregivers, as well as all subjects in pregnancy, to make sure that any loss of life due to PPCM becomes rare or non-existent. No mother should die of PPCM. I foresee the day when that will become fact, as well as full recovery for almost all.

REFERENCES

1. McNarmara DM, Elkayam U, Alharethi R, Damp J, Hsich E, et.al. (2015) Clinical outcomes for peripartum cardiomyopathy in North America: Results of the Investigations of Pregnancy Associated Cardiomyopathy (IPAC) study. *J Am Coll Cardiol* 66: 905-914.
2. Fett JD (2016) Peripartum cardiomyopathy: Challenges in diagnosis and management. *Expert Rev Cardiovasc Ther* 14: 1035-1041.
3. Brar SS, Khan SS, Sandhu GK, Jorgensen MB, Parikh N, et al. (2007) Incidence, mortality and racial differences in peripartum cardiomyopathy. *AM J Cardiol* 100: 302-304.
4. Harper MA, Meyer RE, Berg CJ (2012) Peripartum cardiomyopathy: Population-based birth prevalence and 7 year mortality. *Obstet Gynecol* 120: 1013-1019.
5. Saltzberg MI, Szymkiewicz S, Bianco NR (2012) Characteristics and outcomes of peripartum versus non-peripartum cardiomyopathy in women using wearable cardiac defibrillator. *J Cardiol Fail* 18: 21-27.
6. American Heart Association; Filure H, Fuster V (2009) The AHA guidelines and scientific statements. Oxford: Wiley-Blackwell.
7. Yancy CW, Jessup M, Bozkurt B (2013) A report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol* 62: e147-e239.
8. Fett JD (2013) Earlier detection can help avoid many serious complications of peripartum cardiomyopathy. *Future Cardiol* 9: 809-816.
9. Fett JD (2014) Peripartum cardiomyopathy: A puzzle closer to solution. *World J Cardiol* 6: 87-99.
10. Fett JD, Shah T, McNamara DM (2015) Why do some recovered peripartum cardiomyopathy mothers experience heart failure with a subsequent pregnancy? *Curr Treat Options Cardiovasc Med* 17: 354-359.
11. Ware JS, Li J, Mazaika E, Yasso CM, DeSouze T, et al. (2016) Shared genetic predisposition in peripartum and dilated cardiomyopathies. *NEJM* 374: 233-241.
12. Sheppard R, Hsich E, Damp J, Elkayam U, Kealey A, et al. (2016) GNB3 C825T polymorphism and myocardial recovery in peripartum cardiomyopathy: Results of the multicenter investigations of pregnancy-associated cardiomyopathy study. *Circ Heart Fail* 9: e002683.
13. Fett JD (2017) Reducing the risks for relapse of heart failure in a subsequent pregnancy after peripartum cardiomyopathy? *Future Cardiol* 13: 305-310.
14. Fett JD, Fristoe KL, Welsh SN (2010) Risk of heart failure relapse in subsequent pregnancy among peripartum cardiomyopathy mothers. *Int J Gynecol Obstet* 109: 34-36.
15. Fett JD (2010) Monitoring subsequent pregnancy in recovered peripartum cardiomyopathy mothers. *Crit Pathw Cardiol* 9: 1-3.