

Protective Effect of 20-Hydroxyecdysterone against Lipopolysaccharides-Induced Acute Lung Injury in Mice

Guoying Song¹, Xi-Chao Xia^{1,2*}, Ke Zhang², Yuhong Ma³, Ruixue Yu², Bingjie Li², Mengbo Li², Xiaozhu Yu², Junfeng Zhang² and Shipeng Xue³

¹Library of Pingdingshan University, Pingdingshan, 467000, Henan Province, China

²Basic Medical College, Pingdingshan University, Pingdingshan, 467000, Henan Province, China

³Basic Medicine College, Nanyang Medical University, Nanyang, 473041, Henan Province, China.

Received January 09, 2019; Accepted January 09, 2019; Published May 07, 2019

ABSTRACT

Aim: To investigate the protective effect of 20-hydroxyecdysterone (20E) against lipopolysaccharide-induced acute lung injury (ALI) in mice.

Materials and methods: Mice were respectively administrated a single intratracheal instillation with normal saline (normal group) and 4 mg/kg lipopolysaccharide (LPS) (LPS cohort). Next, animals of LPS cohort were subsequently divided into the model group, the control group, the low-dose group of 20E, middle-group of 20E and high-dose group of 20E. Histological changes of lung were examined by hematoxylin and eosin. Expression levels of tumor necrosis factor- α (TNF- α), interleukin-2 (IL-2), IL-6, IL-8, IL-4 and IL-10 were determined by real-time PCR and enzyme-linked immuno-sorbent assay, respectively.

Results: 20E treatment could result in a decrease of lung damage in the ALI mice. In the 20E treated groups, expression of TNF- α , IL-2, IL-6 and IL-8 was significantly inhibited compared with that model group, respectively. In addition, expression of IL-4 and IL-10 was induced in the 20E treated groups, respectively.

Conclusion: The results suggest that 20E play a protective-role in ALI of mice probably by inhibiting the pro-inflammatory cytokine expression and enhancing the anti-inflammatory cytokine expression.

Keywords: 20-hydroxyecdysterone, Acute lung injury, Protective effects, Pro-inflammation cytokines, Anti-inflammatory cytokines

INTRODUCTION

Acute lung injury (ALI) is considered as a major health problem for elderly population and characterized by activation of the pulmonary endothelium, disruption of the endothelial and alveolar epithelial barriers, and increase of the microvascular permeability [1]. Lung, a crucial airway for pathogens into the body, is widely contacted with a large number of micro-organisms those can cause acute inflammation and ALI [2]. Evidences have demonstrated that that inflammation is closely associated with the occurrence of ALI. In the inflammation, over-production of inflammatory cytokines are involved into injury of lung tissues [3,4]. Lipopolysaccharides (LPS) presented in cell wall of gram-negative bacterium can cause ALI [5]. Lung injury induced LPS is close link with release of macrophage-derived pro-inflammatory cytokines as well as anti-inflammatory cytokines, such as tumor necrosis factor- α

(TNF- α), interleukin (IL)-1, IL-2, IL-6, IL-8, IL-4, IL-10 and interferon- γ [6,7].

Ecdysterone was first isolated from insect and plays a key role in the molting, development and reproduction of animals. Afterward, it was confirmed that the active component of 20-hydroxyecdysterone (20E) is widespread

Corresponding author: Xi-Chao Xia, Associate Chief, Basic Medical College, Pingdingshan University, Pingdingshan 473061, China Henan, P. R. China; Tel: +86 0377 63526123, E-mail: xiachao8336@163.com

Citation: Song G, Xia XC, Zhang K, Ma Y, Yu R, et al. (2019) Protective Effect of 20-Hydroxyecdysterone against Lipopolysaccharides-Induced Acute Lung Injury in Mice. J Pharm Drug Res, 2(3): 109-114.

Copyright: ©2019 Song G, Xia XC, Zhang K, Ma Y, Yu R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

across the plant kingdom and sustain higher concentration in them [8,9]. It has demonstrated that 20E have remarkable pharmacological properties in mammals, including lowering cholesterol levels and blood glucose, stimulating protein synthesis, promoting carbohydrate and lipid metabolism, and inducing stem cell differentiation [10]. Recently, it has been suggested that 20E has obvious antioxidant activity by radical scavenging tests *in vitro* and *in vivo* [11]. Meanwhile, our study indicates that 20E can function as an important player in counteracting memory deficits in rats with diabetes, possibly through enhancing the anti-oxidative ability in the brain [12]. But, little is known about the effect of 20E on the ALI. Thus, the current study was designed to shed light of the effect of 20E on the ALI.

MATERIALS AND METHODS

Reagents

The reagents used in the experiment and their sources were as follows: lipopolysaccharide (LPS) and 20E were obtained from Sigma-Aldrich Co (St. Louis, MO, USA), dexamethasone-21-acetate (Dex) and sodium pentobarbital were purchased from Aladdin (Shanghai, China). PCR primers were synthesized by Sangon Biological Engineering Technology Company (Shanghai China). Trizol reagent and RT-PCR kit including RTase M-MLV, dNTP mixture, Taq DNA polymerase, RNase inhibitor and SYBR Premix Ex Taq™ were obtained from Takara (Dalian, China). Enzyme-linked immuno-sorbent assay (ELISA) kit of mice was purchased from BOSTER (Wuhan, China). All other chemicals were of reagent grade.

Animals

The experimental study was performed following approval from the Pingdingshan University Animal Care and Ethics Committee. All experimental and surgical procedures were conducted by the Pingdingshan University Surgical Application and Research Center. Animal cultures were performed according to the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Male ICR mice (20-24 g) were obtained from Henan Laboratory Animals Center for Medical Science and Research. The animals were fed standard laboratory chow (Wuhan University Center for Animal experiment, Wuhan, China) with free access to water and housed individually at a controlled temperature of 19-25°C with a 12:12 h light/dark cycle.

Animals were randomly divided in six groups (n=15) including normal group, model group, control group (Dex group), low-dose group of 20E, middle-dose group of 20E and high-dose group of 20E. Except normal group, mice in other groups were anaesthetized with an intraperitoneal injection of 40 mg/kg of sodium pentobarbital and treated with a single intratracheal instillation with 4 mg/kg LPS. Based on previous study [12], animals in the low-dose group of 20E, middle-dose group of 20E and high-dose group of

20E received 20E solution at a concentration of 0.1 mg/kg, 1 mg/kg and 10 mg/kg, respectively, and Dex group received Dex at 10 mg/kg. Times of animal treatment were 1, 12, 24 and 36 h before LPS instillation and 12, 24 and 36 h after LPS challenge. Animals from normal group and model group were treated with the same volume of normal saline at corresponding times. The mice were killed by cervical dislocation. Left and right part of lungs were separated, harvested, immediately frozen in liquid nitrogen, and then stored at -80°C until used.

Histological analysis of lung

Left lungs were fixed in 10% formaldehyde and embedded in paraffin. Specimens were cut with 4 µm thick sections those were stained by routine hematoxylin and eosin (H&E) for histological analysis. Sections were analyzed under light microscope (Olympus BX50, Barcelona, Spain).

Real-time PCR assay for TNF-α, IL-2, IL-6, IL-8, IL-4 and IL-10 mRNA level in the lung

Total RNA was extracted from lung tissues using the Trizol reagent according to the manufacturer's protocol. Quality of RNA was monitored by 1.2% agarose gel electrophoresis. First-strand cDNA was synthesized using M-MLV reverse transcriptase.

To determine levels of TNF-α, IL-2, IL-6, IL-8, IL4 and IL10 transcript derived from lungs, real-time quantitative PCR was performed following the manufacture instruction of SYBR Premix Ex Taq™. Primers used in this study were listed in **Supplemental Table 1**. PCR was performed using an ABI 7500 Real-Time Detection System (Applied Biosystems, USA) in accordance with manufacture. Based on standard curve constructed, the expression levels of TNF-α, IL-2, IL-6, IL-8, IL-4 and IL-10 were calculated by $2^{-\Delta\Delta CT}$.

Table 1. Primers used in the experiment.

Primer	Sequence (5'–3')
TNF-α-F	ACTGGGCATCCTTTGTCTCGG
TNF- α-R	CTGTTGGTGAAGAGGTGCGG
IL-2-F	ATCGTGGGTGGCGTTGAG
IL-2-R	GTTCTGGCGGATGTTGTGG
IL-6-F	CTCAGGACGGTAAGTGTCGCTT
IL-6-R	TGGTAGAACTGGAAGGACGGCT
IL-8-F	GGGTTTCCTTACTCCGCTTTTCG
IL-8-R	GTCAGTGCCTTGGTTTCAGGTCTG
IL-4-F	TTGAAGAAAGCAGACGCCCC
IL-4-R	TGGCAGGTTGAGTGGCAGC
IL-10-F	ATCGCAACCCATCAAACCTCG
IL-10-R	ACAATCGTCGCAGACACGGT
β-actin-F	CATCCACGAGACCACCTACAAC
β-actin-R	GAAATACTGCCTCGCTCCCTC

ELISA assay for the TNF- α , IL-2, IL-6, IL-8, IL4 and IL10 protein levels in the lung

Right lungs were weighted, perfused with chilled normal saline, then cut into small pieces, placed in 0.2 M phosphate buffer (pH 7.4) and homogenized using homogenizer to obtain 20% homogenate. Resultant homogenate was centrifuged (5000x g for 15 min, 4°C) and the supernatant was sub packaged to sterile eppendorf tubes. Levels of TNF- α , IL-2, IL-6, IL-8, IL-4 and IL-10 in lung were measured by ELISA kits. All the procedures were performed in accordance with the manufacturer's instructions.

Statistical analysis

Original data were analyzed by Statistical Package for Social Sciences software (version 14.0) (SPSS, Inc., Chicago, IL). Results were expressed as mean $\pm$ SEM. Multiple sets of data comparison were carried out by one-way analysis of

variance (ANOVA). Comparison of data between groups were treated by least significance difference (LSD) combined with Tamhane's way. Values with $P < 0.05$ were considered as statistically significant.

RESULTS

Histopathologic analysis

Histopathological examination of normal animals showed normal cellular architecture with distinct alveolar cell (**Figure 1A**). The lung sections of model group exhibited an obvious damage characterized by focal hemorrhage, distortion and alveolar thickening (**Figure 1B**). Compared with that of model group, injury derived from LPS was ameliorated by Dex treatment and different-dose 20E treatment (**Figures 1C-1F**). Furthermore, obvious protecting-role was observed in Dex and high-dose group of 20E (**Figures 1C and 1F**).




Figure 1. Effects of 20E on LPS-induced lung histopathological change (hematoxylin and eosin stain, magnification: 400). (A) Normal group, (B) LPS group, (C) Dex group, (D) Low-dose group of 20E, (E) Middle-dose group of 20E, (F) High-dose group of 20E.

Effects of 20E on the TNF- α , IL-2, IL-6, IL-8, IL-4 and IL-10 mRNA level in the lung

Compared with that of normal group and Dex group, the mRNA levels of TNF- α , IL-2, IL-6 and IL-8 were significantly increase in the model group (**Figure 2**). Administration of different-dose of 20E could result in a

dose-dependent decline of TNF- α , IL-2, IL-6 and IL-8 level. In the high-dose group of 20E, mRNA level of TNF- α , IL-2, IL-6 and IL-8 decreased 58.1% ($P < 0.01$), 61.1% ($P < 0.01$), 52.1% ($P < 0.01$) and 47.7% ($P < 0.01$) compared with that of model group (**Figure 2**), respectively.




Figure 2. Effects of 20E on the TNF- α , IL-2, IL-6 and IL-8 mRNA level in the lung.

Note: $n=5$ /each tissue. ^a $P<0.05$, ^b $P<0.01$ vs control group at normal group. ^c $P<0.05$, ^d $P<0.01$ vs control group at the model group

Compared with that of normal group and Dex group, mRNA levels of IL-4 and IL-10 were significantly decreased in the model group (Figure 3). But, in high-dose group of 20E, mRNA level of IL-4 and IL-10 increased 80.6% ($P<0.01$) and 116.8% ($P<0.01$) in contrast with that of model group, respectively (Figure 3).




Figure 3. Effects of 20E on the IL-4 and IL-10 mRNA level in the lung.

Note: $n=5$ /each tissue. ^a $P<0.05$, ^b $P<0.01$ vs control group at normal group. ^c $P<0.05$, ^d $P<0.01$ vs control group at the model group

Effects of 20E on the TNF- α , IL-2, IL-6, IL-8, IL-4 and IL-10 expression in the lung

Compared with that of normal group and Dex group, in a significant increase of TNF- α , IL-2, IL-6 and IL-8 level was observed in the model group ($P<0.01$) (Table 2). Expression levels of TNF- α , IL-2, IL-6 and IL-8 showed a decline trend in different 20E treated groups compared with that of model group. In the high-dose group of 20E, level of TNF- α , IL-2, IL-6 and IL-8, respectively decreased 48.3% ($P<0.01$), 66.3% ($P<0.01$), 26.9% ($P<0.05$) and 45.7% ($P<0.01$) (Table 2), but the level of IL-4 and IL-10 increased 1.9 times ($P<0.01$) and 2.9 times ($P<0.01$) in contrasted with that of model group (Table 2).

Table 2. Effects of 20E on the TNF- α , IL-2, IL-6, IL-8, IL-4 and IL-10 protein level in the lung.

Group	TNF- α	IL2	IL-6	IL8	IL-4	IL-10
Normal group	0.64 $\pm$ 0.17	1.17 $\pm$ 0.45	0.71 $\pm$ 0.13	0.88 $\pm$ 0.14	3.15 $\pm$ 0.35	1.89 $\pm$ 0.41
Model group	2.89 $\pm$ 0.27 ^b	3.43 $\pm$ 0.73 ^b	1.70 $\pm$ 0.21 ^b	2.13 $\pm$ 0.34 ^b	0.93 $\pm$ 0.27 ^b	0.70 $\pm$ 0.25 ^b
Dex group	1.11 $\pm$ 0.14 ^{ad}	1.48 $\pm$ 0.51 ^d	0.98 $\pm$ 0.11 ^d	1.26 $\pm$ 0.31 ^d	2.45 $\pm$ 0.32 ^d	1.33 $\pm$ 0.38 ^{ad}
Low-dose group	2.08 $\pm$ 0.33 ^b	3.12 $\pm$ 0.40 ^b	1.69 $\pm$ 0.40 ^b	2.43 $\pm$ 0.23 ^b	0.94 $\pm$ 0.22 ^b	0.76 $\pm$ 0.31 ^b
Middle-dose group	1.47 $\pm$ 0.21 ^{bd}	2.44 $\pm$ 0.34 ^{bd}	1.55 $\pm$ 0.33 ^b	1.91 $\pm$ 0.30 ^b	1.69 $\pm$ 0.31 ^{ac}	1.21 $\pm$ 0.18 ^{ac}
High-dose group	1.21 $\pm$ 0.16 ^{ad}	1.34 $\pm$ 0.51 ^d	0.81 $\pm$ 0.27 ^d	1.11 $\pm$ 0.29 ^d	2.71 $\pm$ 0.34 ^d	1.53 $\pm$ 0.27 ^d

Note: $n=5$ /each tissue. ^a $P<0.05$, ^b $P<0.01$ vs control group at normal group. ^c $P<0.05$, ^d $P<0.01$ vs control group at the model group

DISCUSSION

In the current study, administration of 20E resulted in a decrease of the alveolar cell damage and decline of the TNF- α , IL-2, IL-6 and IL-8 expression in the ALI mice, which suggested a protective-role of 20E against LPS injury is closely associated with suppression of TNF- α , IL-2, IL-6 and IL-8 production. Inflammation cascade is initiated by the innate immune system, in which TNF- α , IL-2, IL-6 and IL-8 play a key role in the inflammatory response [13]. Generally, inflammatory response is taken as a direct pathogenic-signal in ALI [14]. It has been demonstrated that the over-expression of these pro-inflammatory mediators contribute to ALI or lung cancer [15]. Thus, down-regulation of TNF- α , IL-2, IL-6 and IL-8 expression derived from 20E is beneficial in reducing the lung damage of mice. Similar studies are also reported in ALI cases. Administration of dexamethasone-21-acetate can dramatically reduce the concentration of cytokines TNF- α , IL-2, IL-6 against the damage of LPS on lung [16]. In addition, 1 α , 25-dihydroxyvitamin D3 and vitamin D3 analogue can attenuate the innate immunity response in a mouse model of ALI through suppression of IL-8 production [17,18]. Notably, wide applications of steroids are observed in medicine besides dexamethasone-21-acetate, 1 α , 25-dihydroxyvitamin D3 and vitamin D3 analogue [19-21]. Thus, these studies should open a new road to explore of 20E functions.

Further research showed the expressions of IL-4 and IL-10 were obviously induced in the middle-dose group of 20E and high-dose group of 20E, suggesting multiple targets acted on by 20E are involved in the protection event against ALI. Inflammatory response of organs is coordinated by the interplay of anti-inflammatory cytokines and pro-inflammatory cytokines, where a low concentration of pro-inflammation cytokines and/or a high concentration of anti-inflammatory cytokines contribute to protection of tissues and cells against damage and sustain homeostasis of body [22,23]. Meanwhile, the IL-4 and IL-10, as anti-inflammatory cytokines, their expression levels that is induced by 20E is helpful in inhibiting the expression of pro-inflammatory cytokines [24,25]. For example, IL-10 works as a key player to reduce the release of pro-inflammatory mediators during ALI, such as IL-6 and inducible nitric oxide synthase [24,26]. Therefore, we postulate that the relative level of anti-inflammatory cytokines and inflammatory cytokines play a crucial role in organs against injury [22,23].

In the current study, protective-role of 20E against ALI in the mice showed a dose-dependent effect suggesting that different targets acted on by 20E are likely involved in inflammation of ALI. In the insect and crustacean, 20E, as an important hormone that plays a key role in regulating development, growth, reproduction and innate immunity by acting on different targets [27,28]. Considering here, the

study of TNF- α , IL-2, IL-6 and IL-8, IL-4 and IL-10 temporal and spatial expressions derived from 20E activation is an interesting work. Meanwhile, regulation of TNF- α , IL-2, IL-6 and IL-8, IL-4 and IL-10 derived from different-dose 20E treatment groups is complex event. Therefore, great efforts are required to elucidate the mechanisms of 20E in ALI.

ACKNOWLEDGEMENT

This research was funded by the National Natural Science Foundation of Henan (No. 18A330004, PXY-BSQD-2018009, 17A180010, PXY-PYJJ-2018005) and China Postdoctoral Science Foundation Funded Project (2016M590143).

CONFLICT OF INTEREST

We declare that there is no conflict of interest.

REFERENCES

1. Nakagomi T, Kitada O, Kuribayashi K, Yoshikawa H, Ozawa K, et al. (2004). The 150 kDa oxygen-regulated protein ameliorates lipopolysaccharide-induced acute lung injury in mice. *Am J Pathol* 165: 1279-1288.
2. Sachs UJ (2011) Recent insights into the mechanism of transfusion-related acute lung injury. *Curr Opin Hematol* 18: 436-442.
3. Ebihara I, Hirayama K, Nagai M, Kakita T, Sakai K, et al. (2011) Angiopoietin balance in septic shock patients with acute lung injury: Effect of direct hemoperfusion with polymyxin B-immobilized fiber. *Ther Apher Dial* 15: 349-354.
4. Wilson CN, Vance CO, Doyle TM, Brink DS, Matuschak GM, et al. (2012). A novel post-exposure medical counter measure L-97-1 improves survival and acute lung injury following intratracheal infection with *Yersinia pestis*. *Innate Immun* 18: 373-389.
5. Hu LX, Du YY, Zhang Y, Pan YY (2012) Lack of association between interleukin-8-251 target; A polymorphism and colorectal cancer risk: A meta-analysis based on 3,019 cases and 3,984 controls. *Asian Pac J Cancer Prev* 13: 5075-5079.
6. Ju ST, Sharma R, Gaskin F, Fu SM (2012) IL-2 controls trafficking receptor gene expression and Th2 response for skin and lung inflammation. *Clin Immunol* 145: 82-88.
7. Zhang CR, Lin JC, Xu WM, Li M, Ye HS, et al. (2013) Interleukin-12 and interleukin-2 alone or in combination against the infection in invasive pulmonary aspergillosis mouse model. *Mycoses* 56: 117-122.
8. Hung TJ, Chen WM, Liu SF, Liao TN, Lee TC (2012) 20-Hydroxyecdysone attenuates TGF- β 1-induced renal

- cellular fibrosis in proximal tubule cells. *J Diabetes Complicat* 26: 463-469.
9. Jian CX, Liu XF, Hu J, Li CJ, Zhang G, et al. (2013) 20-hydroxyecdysone-induced bone morphogenetic protein-2-dependent osteogenic differentiation through the ERK pathway in human periodontal ligament stem cells. *Eur J Pharmacol* 698: 48-56.
 10. Kumpun S, Girault JP, Dinan L, Blais C, Maria A, et al. (2011) The metabolism of 20-hydroxyecdysone in mice: Relevance to pharmacological effects and gene switch applications of ecdysteroids. *J Steroid Biochem* 126: 1-9.
 11. Foucault AS, Mathé V, Lafont R, Even P, Dioh W, et al. (2011) Quinoa extract enriched in 20-hydroxyecdysone protects mice from diet-induced obesity and modulates adipokines expression. *Obesity* 20: 270-277.
 12. Xia XC, Yuan GQ, Zhang QY, Liu RZ, Wang ZX, et al. (2014) Effects of 20-hydroxyecdysone on improving memory deficits in streptozocin-induced type 1 diabetes mellitus in rat. *Eur J Pharmacol* 740: 45-52.
 13. Sharma R, Sung SS, Gaskin F, Fu SM, Ju ST (2012) A novel function of IL-2: chemokine/chemoattractant/retention receptor genes induction in Th subsets for skin and lung inflammation. *J Autoimmun* 38: 322-331.
 14. Fang Y, Xu P, Gu C, Wang Y, Fu XJ, et al. (2011) Ulinastatin improves pulmonary function in severe burn-induced acute lung injury by attenuating inflammatory response. *J Trauma* 71: 1297-1304.
 15. Yao HY, Zhang LH, Shen J, Shen HJ, Jia YL, et al. (2011) Cytoporus polysaccharide prevents lipopolysaccharide-induced acute lung injury associated with down-regulating toll-like receptor 2 expression. *J Ethnopharmacol* 137: 1267-1274.
 16. Yubero S, Manso MA, Ramudo L, Vicente S, De Dios I (2012) Dexamethasone down-regulates the inflammatory mediators but fails to reduce the tissue injury in the lung of acute pancreatitis rat models. *Pulm Pharmacol Ther* 25: 319-324.
 17. Takano Y, Mitsuhashi H, Ueno K (2011) 1 α ,25-dihydroxyvitamin D₃ inhibits neutrophil recruitment in hamster model of acute lung injury. *Steroids* 76: 1305-1309.
 18. Takano Y, Mitsuhashi H, Ishizuka S, Takahashi K, Chokki M, et al. (2012) TEI-A00114: A new vitamin D₃ analogue that inhibits neutrophil recruitment in an acute lung injury hamster model while showing reduced hypercalcemic activity. *Steroids* 77: 1535-1542.
 19. Jules-Elysee KM, Wilfred SE, Memtsoudis SG, Kim DH, YaDeau JT, et al. (2012) Steroid modulation of cytokine release and desmosine levels in bilateral total knee replacement: A prospective, double-blind, randomized controlled trial. *J Bone Joint Surg Am* 94: 2120-2127.
 20. Wigenstam E, Jonasson S, Koch B, Bucht A (2012) Corticosteroid treatment inhibits airway hyper responsiveness and lung injury in a murine model of chemical-induced airway inflammation. *Toxicology* 301: 66-71.
 21. Chazenbalk G, Singh P, Irge D, Shah A, Abbott DH (2013) Androgens inhibit adipogenesis during human adipose stem cell commitment to pre-adipocyte formation. *Steroids* 78: 920-926.
 22. Sadr K, Walker RH (2012) The balancing act of the inflammatory cascade after bilateral total knee arthroplasty commentary on an article by Kethy M. Jules-Elysee, MD, et al.: "steroid modulation of cytokine release and desmosine levels in bilateral total knee replacement. A prospective, double-blind, randomized controlled trial". *J Bone Joint Surg Am* 94: e1791-1792.
 23. Yang R, Yang L, Shen X, Cheng W, Zhao B, et al. (2012) Suppression of NF- κ B pathway by crocetin contributes to attenuation of lipopolysaccharide-induced acute lung injury in mice. *Eur J Pharmacol* 674: 391-396.
 24. Sun L, Cornell TT, LeVine A, Berlin AA, Hinkovska-Galcheva V, et al. (2013) Dual role of interleukin-10 in the regulation of respiratory syncytial virus (RSV)-induced lung inflammation. *Clin Exp Immunol* 172: 263-279.
 25. Wechsler ME (2013) Inhibiting interleukin-4 and interleukin-13 in difficult-to-control asthma. *N Engl J Med* 368: 2511-2513.
 26. Ma HJ, Huang XL, Liu Y, Fan YM (2012) Sulfur dioxide attenuates LPS-induced acute lung injury via enhancing polymorphonuclear neutrophil apoptosis. *Acta Pharmacol Sin* 33: 983-990.
 27. Flatt T, Heyland A, Rus F, Porpiglia E, Sherlock C, et al. (2008) Hormonal regulation of the humoral innate immune response in *Drosophila melanogaster*. *J Exp Biol* 16: 2712-2724.
 28. Wang JL, Liu XS, Zhang Q, Zhao HB, Wang YF (2012) Expression profiles of six novel C-type lectins in response to bacterial and 20E injection in the cotton bollworm (*Helicoverpa armigera*). *Dev Comp Immunol* 7: 221-232.