

Effect of Carbon Dioxide Concentration on Cell Growth and Oil Yield of Freshwater and Marine Microalgae

Mariam Al Hattab¹ and Abdel Ghaly^{1*}

¹Department of Process Engineering and Applied Science, Faculty of Engineering, Dalhousie University, Halifax, Nova Scotia, Canada

Received March 26, 2015, Accepted April 6, 2015, Published April 15, 2015

ABSTRACT

The *Chlorella saccharophila* (freshwater) and *Tetraselmis suecica* (marine) microalgae were investigated for their potential as feedstock for the production of biodiesel. Assessment was based on their ability to produce biomass and lipid accumulation. Varying concentrations of CO₂ (3, 6 and 9%) were used at 24 h light exposure. The results indicated that *T. suecica* produced higher cell yields compared to the *C. saccharophila* under all parameters tested. Statistical analysis indicated that the biomass yields achieved using CO₂ at varying concentrations were significantly different from one another. However, varying CO₂ concentrations over the range of 3 to 9% did not significantly affect the oil yields for both species over the elapsed time. Thus, from an economic stand point it is much more suitable to use CO₂ at a concentration of 3% as opposed to higher concentrations. The use of NaHCO₃ as a carbon source on the biomass and oil yields were also evaluated using the same experimental parameters. Results indicated that the CO₂ carbon source resulted in higher biomass and oil yields for the marine microalgae species when compared to NaHCO₃ carbon source. However, the *Chlorella saccharophila* species resulted in higher biomass and lipid yields using the sodium bicarbonate carbon source. This suggests that different species have a preferred carbon source. Some are better at the uptake of one source over the other. Lower oil yields were achieved using the 3% CO₂ as the carbon source compared to NaHCO₃ for the *Chlorella saccharophila* species. The *Tetraselmis suecica* species resulted in a slight increase in oil yield using 3% CO₂ as opposed to NaHCO₃. The optimal growth conditions for *Chlorella saccharophila* are the combination of nutrients, with 24 h light exposure and NaHCO₃ as a carbon source and those for *Tetraselmis suecica* are the ammonium nitrate, the 24 h light exposure and 3% CO₂ as a carbon source.

Keywords: Freshwater, Marine Water, Microalgae, Cell yield, Oil Yield, NaHCO₃, CO₂, Carbon Source.

INTRODUCTION

The demonising fossil fuel reserves and the environmental concerns associated with burning fossil fuels have accelerated the need for a renewable energy source that is environmentally friendly. Increased carbon dioxide emissions have been correlated with the amount of fossil fuel being burnt [1]. Biofuels such as biodiesel and bioethanol are promising substitution for petroleum fuel source [2]. Numerous feedstocks can be used as biomass for biofuel generation which include food waste, agricultural waste, municipal waste and both edible and nonedible oilseeds [3]. Currently, the best crops for biofuel production are oilseeds, but they are considered a food source for many people around the world [2,3]. However, microalgae have been noted to store oil that is 10 folds higher than the leading plant crop [4].

Microalgae are abundant microorganisms in nature, able to convert carbon dioxide into biomass which can then be used for biodiesel production via a transesterification reaction process [5]. Microalgae as an alternate fuel source is ideal

because of their high growth rates and their ability to store higher lipids than the leading crop plants [6]. In addition, the waste that is generated after oil extraction can be used for other value added products, such as animal feed, organic fertilizers, and other biofuel products such as methane and ethanol via fermentation [7]. The amount of lipids stored in

Corresponding author: Professor A. E. Ghaly, Department of Process Engineering and Applied Science, Faculty of Engineering, Dalhousie University, Halifax, Nova Scotia, Canada; [Tel:902-494-6014](tel:902-494-6014); Email: abdel.ghaly@dal.ca

Citation: Al Hattab M & A Ghaly (2015) Effect of carbon dioxide concentration on cell growth and oil yield of freshwater and marine microalgae. International Journal of Bioprocess and Biotechnological Advancements, 1(1):31-44

Copyright: ©2015 Al Hattab M, Ghaly A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the microalgal cells can be manipulated by changing the environmental parameters such as temperature, pH, nutrient source, carbon source and light duration [8-10].

Biodiesel is a liquid fuel that is biodegradable and nontoxic [11]. It generates the same amount of energy (calorific value) as that produced using petroleum diesel without the resale of harsh compounds such as NO_x, SO_x and hydrocarbons into the atmosphere [3,12]. Biodiesel can be used in existing diesel engines without the need for much modification [2]. It is for these reasons that biodiesel is regarded as the best renewable energy source that is environmentally friendly and a viable source for replacement of the currently used petroleum source.

OBJECTIVES

The aim of this study was to investigate the possibility of increasing the microalgae cell growth rate and oil yield by exposing the microalgae to various carbon sources (NaHCO₃ and CO₂) in a specially designed pilot scale open pond system. The specific objectives were: (a) to study the effect of CO₂ concentration in the air on biomass yield and oil content at three levels (3:97, 6:94 and 9:91 v/v CO₂ to air) and (b) to evaluate the effect of carbon source (NaHCO₃ and CO₂) on the microalga biomass yield and oil content.

MATERIALS AND METHODS

Experimental Apparatus

A fully automated multiple open pond system (Figure 1) consisted of a frame, 18 open pond units, a cooling unit, a lighting unit, a supernatant collection unit and control unit was used in this study.

The frame (244 cm in width x 41 cm in depth x 283 cm in height) consisted of three shelves (76 cm apart) and housed the open pond, light, cooling, water collection and control units. Each shelf was divided vertically into two sides by a 1.2 cm thick plywood sheet to provide a better control of light and feed. The open pond unit consisted of six ponds, each was made of galvanized steel and was divided into three compartments (each was 38 cm in length x 38 cm in width x 12.5 cm in height and can hold up to 18 L). The lighting unit provided 430 hectolux of illumination per shelf (480 μmol m⁻² s⁻¹) using a mixture of fluorescent and incandescent lamps (six 40 W cool white fluorescent lamps 122 cm in length and four 100 W incandescent bulbs) mounted on each shelf, that sit 100 cm away from the ponds. A cooling unit was designed to continuously remove the heat produced by the lamps to avoid heating of the algae on the upper and middle shelves. A 5 cm diameter PVC pipe (having 6 mm diameter holes spaced 6 cm apart and facing out) was placed under the backside of the ponds. Two metal blocks placed under each pond provided a 5 cm space between the pond and the lighting system of the shelf below it. A 5 cm diameter PVC pipe was attached vertically to the left side of the frame and acted as a manifold through which air was blown by means of a motor




Figure 1. Experimental Apparatus.

driven fan (Model AK4L143A Type 821, Franklin Electric, Bluffton, Indiana, United States of America). The supernatant from each tray was collected in a separate container (2.7 L each) located at the bottom of the system. The outlets were connected to plastic tubes of 1 cm outside diameter, which were passed through a solenoid valve.

A computer was used to operate and control the various components of the open pond system and record the various measurements. The light intensity was measured using a Quantum Sensor, SQ-316 Series (Apogee, Logan, Utah, United States of America). The pH was measured using pH electrodes (EW-59001-65, Cole Parmer, Montreal, Quebec, Canada). The temperature was measured using thermocouples (WD-08541-12, Nova-Tech International, Houston, Texas, United States of America). A basic computer program (BASIC Stamp Editor v2.5) allowed the configuration of the operating frequency and duration of the light, aeration unit and collection system. The computer was connected to a data coordinator (cDAQ-9178, National Instruments) which had 24 digital output ports and 24 digital input ports. The digital output ports were connected to electronic circuits which were responsible for the lighting, cooling and collection systems.

Microalgae

One freshwater microalgae (*Chlorella saccharophila*) and one marine (*Tetraselmis suecica*) were selected based on their ability to yield high biomass and store lipids [13].

The freshwater strain *Chlorella saccharophila* was selected for study because of its high lipid content (45%). This strain is capable of achieving a biomass yield of 3.88 g/L, which is not the highest among the freshwater species, but can however be offset by the fact that it achieves the highest lipid content. This results in a lipid yield of 1.75 g/L. The highest biomass yielding algae *Scenedesmus obliquus* of 4.34 g/L only achieves a lipid content of 38%, which intern results in a lipid yield of 1.69 g/L. *Chlorella saccharophila* is a green unicellular microalga belonging to the *Chlorella* genus [14]. The cells have an average size of 7.3 μm [15]. The cells contain a single chloroplast enclosed in a spherical or subspherical form. These cells reproduce asexually through production of non-motile autospores [16]. This species is able to use glucose [17], bicarbonate and carbon dioxide as the carbon source for growth [18]. The optimal temperature and pH for growth are 20-24°C and 7.5-8, respectively.

The marine microalgae strain *Tetraselmis suecica* was selected for this study because of its high biomass yield of 4.48 g/L and comparatively high lipid content. This species achieves a lipid content of 23% which is not the highest among the other species but can, however, be offset by the fact that it achieves the highest biomass yield. This results in a lipid yield of 1.03 g/L, while the *Chaetoceros muelleri*, with the highest lipid content of 34%, only achieves a biomass yield of 0.98 g/L, which results in a lipid yield of 0.33 g/L. *Tetraselmis suecica* grows as single cells. They are motile and can be compressed or curved, but they are never twisted

[19]. The cells are spherical or elliptic with a length of 35 μm and a width of 14 μm . This species is able to use both sodium bicarbonate [20] and carbon dioxide [21] as the carbon source for growth. The optimal temperature and pH for growth are 18-24°C and 7-9, respectively [22].

Experimental Design

Two set of experiments were carried out. In the first set of experiments, the selected freshwater (*Chlorella saccharophila*) and marine (*Tetraselmis suecica*) microalgae species were grown in an open pond system using NaHCO_3 as a carbon source. The sodium bicarbonate (NaHCO_3) was administered at a concentration of 1300 mg/L. Ammonium nitrate was used for the marine microalgae and a combination of nutrients (ammonium nitrate, ammonium sulfate and ammonium phosphate) was used for the freshwater algae as sources of nitrogen at the optimum light exposure (24 h) as recommended by [13]. The light intensity was kept at 480 $\mu\text{mol/m}^2 \text{s}^{-1}$ and the nitrogen content, pH and temperature were kept constant at 70 mg/L, 8.3-8.9 and 22°C, respectively. In the second set of experiments the effects of carbon dioxide concentration on the algae biomass and oil content were evaluated. Carbon dioxide was administered at concentrations of 3, 6 and 9% (v/v in air). The algae were exposed to full light exposure (24 h and the light intensity was kept at 480 $\mu\text{mol/m}^2 \text{s}^{-1}$). A combination of nutrients (ammonium nitrate, ammonium phosphate and ammonium sulfate) was used as nutrient for the freshwater microalgae and ammonium nitrate was used for the marine microalgae. The nitrogen content, pH and temperature were the same as in the first set of experiment. The best results obtained with CO_2 were compared with those obtained with NaHCO_3 .

Preparation of Liquid Medium for Inoculum Growth

The freshwater microalgae medium was prepared on algal proteose medium (ATCC Catalog Medium No. 847, American Type Culture Collection, Manassas, Virginia, United States of America) and was made up by adding 1 g of proteose peptone (Difco 0120) to 1 L of Bristols solution (Table 1). Bristols solution was prepared by adding the following amounts from the prepared stock solutions: 10 mL of NaNO_3 , 10 mL of CaCl_2 , 10 mL of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 10 mL of K_2HPO_4 , 10 mL of KH_2PO_4 , 10 mL of NaCl , 0.05 mL of FeCl_3 and 940 mL of distilled water. The stock solutions were prepared as follows: 10 g of NaNO_3 in 400 mL of distilled water, 1g of CaCl_2 in 400 mL of distilled water, 3 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ in 400 mL of distilled water, 3 g of K_2HPO_4 in 400 mL of distilled water, 7 g of KH_2PO_4 in 400 mL of distilled water and 1 g of NaCl in 400 mL of distilled water.

The marine microalgae medium was prepared in F/2 medium [23]. The trace element liquid medium stock solution (Table 2) was prepared by the addition of 4.16 g of Na_2EDTA , 3.15 g of $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, 0.01 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.022 g of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.18 g of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ and 0.006 g of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ into 1 L of

to 125 mL Erlenmeyer flask containing 25 mL of F/2 liquid media and then left to grow at room temperature for 2 weeks at a photocycle of 14 h light and 10 h dark. The mixture was then transferred to a 500 mL Erlenmeyer flask containing 250 mL of F/2 liquid media and was left to grow for 2 weeks at a photocycle of 14 h light and 10 h dark. Finally, the media was transferred from the 500 mL flask into a 30 L bioreactor containing 25 L of F/2 liquid media and left to grow for 2 additional weeks at a cycle of 14 h light and 10 hour dark.

Preparation of Algae Production Media

The freshwater production medium is a modification of the Fitzgerlad medium [24]. The preparation of the stock solutions for this media is shown in Table 3. The medium was made up by the addition of 1 mL of each of the stock solutions A, B, C and D to 1 L distilled water (Table 4).

A modification of the F/2 media [23] was used as the production medium for the marine microalga. The medium was modified by eliminating the addition of sodium nitrate. The medium consists primarily of autoclaved ocean water (Halifax Waterfront, Halifax, Nova Scotia, Canada). Table 5 shows the elemental analysis of the components present in the marine water which was performed at the Mineral Engineering Center of Dalhousie University.

Experimental Protocol

To each compartment in the open pond system a total of 4.75 L of freshwater production media was added. The amount of nutrient was added to the production medium. This solution was enriched with the desired carbon source (1.3 g/L of sodium bicarbonate, 3% CO₂, 6% CO₂ or 9% CO₂). To this, 250 mL of *Chlorella saccharophila* inoculum was added to each compartment. The cells were exposed to 24 h light and left to grow for 10 days. Every other day, 100 mL sample was taken for experimental analyses. The samples were analyzed for pH and biomass yield. At the end of the run the biomass was harvested from the liquid media using a Sorvall T1 Centrifuge (Thermo Scientific, Marietta, Ohio, United States of America). The supernatant from the centrifuge tubes was decanted and the cells were collected for biomass yield and oil content analyses. The marine medium was used with marine algae and the same procedure was followed using the ammonium nitrate nutrient system.

Microalgae Biomass Determination

The freshwater microalga yield was determined by measuring the optical density at 484 nm from a standard curve between the cell count and optical density. The number of Colony Forming Units (CFU) for *Chlorella saccharophila* was determined using a series of dilutions. A test tube containing 9 mL of autoclaved distilled water and a 1 mL aliquot sample was added to the tube. The contents of the tube were vortexed (Thermolyne Maxi Mix, Thermolyne Corporation, Hampton, New Hampshire, United States of America) to distribute the cells. A 1 mL aliquot of this solution was added to another tube that had been autoclaved

Table 3. Formulation of stock solutions for *Chlorella saccharophila* production medium.

Stock Solutions (per 200 mL)	Composition
A	24.648 g MgSO ₄ •7H ₂ O
B	1.360 g KH ₂ PO ₄
	8.700 g K ₂ HPO ₄
C	1.392 g FeSO ₄ •7H ₂ O
	1.864 g EDTA tri Na
D	0.620 g H ₃ BO ₃
	0.340 g MnSO ₄ •H ₂ O
	0.057 g ZnSO ₄ •7H ₂ O
	0.018 g (NH ₄) ₆ Mo ₇ O ₂₄ •4H ₂ O
	0.027 g CoCl ₂ •6H ₂ O
	0.024 g KBr
	0.017 g KI
	0.023 g CdCl ₂ •5/2H ₂ O
	0.091 g Al ₂ (SO ₄) ₃ (NH ₄) ₂ SO ₄ •24H ₂ O
	0.040 mg CuSO ₄ •5H ₂ O
0.560 mL H ₂ SO ₄ (97%)	

Table 4. Components of freshwater production medium.

Component	Amount (mL)
A	1
B	1
C	1
D	1
Distilled Water	996

Table 5. Elemental analysis of autoclaved ocean water used for marine production media.

Element	Amount (mg/L)
Na	10 254.00
Mg	1 078.00
S	1 010.00
K	395.00
Ca	386.00
Sr	6.79
Si	2.80
P	0.10
Ba	0.05
Al	0.05
Ni	0.04
Zn	0.02
Mo	0.01
Cd	0.01
Co	0.01
Cu	0.01

with 9 mL of distilled water. This tube was again vortexed to distribute the cells. This was repeated 7 times to obtain dilutions of 1:1, 1:10, 1:100, 1:1000, 1:10 000, 1:100 000, 1:1 000 000. For each of the dilutions made, 0.1 mL of the solution was added to a petri dish containing solid freshwater medium. The plates were sealed with parafilm, inverted and incubated at room temperature (~24°C) at a photocycle of 14 hours light and 10 hours dark for 3 days. The plates were then

Microalgae Oil Content Using CO₂ as a Carbon Source

The oil yield results are depicted in Table 6. Analysis of the variance (ANOVA) was performed on the oil yield data as shown in Table 9. The effects of microalgae type on oil yield were significant at the 0.003 level. However, the effect of CO₂ concentration and the interactions between microalgae type and CO₂ concentration were not significant. Tukey's grouping was used to test the differences among the levels of each parameter as shown in Table 10. The two microalgae *Chlorella saccharophila* and *Tetraselmis suecica* were significantly different from one another at the 0.05 level. The highest mean oil yield (4.18%) was obtained from the freshwater microalgae species. The CO₂ concentrations were not significantly different from one another at the 0.05 level. The highest mean oil yield (3.16%) was obtained with the 3% CO₂ concentration.

Effect of Microalgae Type

The effect of the microalgae type on the oil content is illustrated in Figure 7. *Chlorella saccharophila* achieved the highest oil yields at all CO₂ concentration. It produced average oil yields of 4.71, 3.90 and 3.59% while *Tetraselmis suecica* produced average oil yields of 2.09, 1.01 and 2.43% at the 3%, 6% and 9% CO₂ concentrations, respectively. These results are similar to those of Pittman et al. [47] which indicated that the marine microalgae produce much lower oil yields compared to freshwater microalgae. The oil yields obtained from *Chlorella saccharophila* were 4 times higher than those obtained from *Tetraselmis suecica*, despite the higher biomass yields obtained from the *Tetraselmis suecica* species.

Sharma et al. [48] stated that the occurrence and extent to which lipids are produced by microalgae is species/strain specific. Pittman et al. [47] stated that different species use their energy for different metabolic pathways. In this study a trade-off between cell generation and lipid accumulation was seen among the species. The marine species used most of its energy for cell generation as opposed to lipid while the freshwater species used most of its energy for oil accumulation as opposed to cell generation.

Demirbas [7], Moheimani, [33], Sobczuk et al., [49], Sukenik et al., [50], Wagenan et al., [51] and Pagnanelli et al., [52] reported a lipid content in the range of 36-47% and 15-23% for *Chlorella saccharophila* and *Tetraselmis suecica* species, respectively. The differences in the lipid content are

attributed to the different nutrient systems used and the culture age before harvest.

Effect of Carbon Dioxide Concentration

The effect of carbon dioxide concentration on the oil yield is illustrated by Figure 8. As the carbon dioxide concentration was increased from 3% to 9%, the oil yield decreased from 4.7% to 3.6% for the freshwater microalgae (*Chlorella saccharophila*). On the other hand as the carbon dioxide concentration was increased from 3% to 6% the oil yield decreased from 2.1% to 1.0% for the marine microalgae (*Tetraselmis suecica*). However, a further increase in CO₂ concentration to 9% increased the oil yield to 2.43%.

It should be noted that the trends in Figure 9 are the opposite of the trends of cell yield shown in Figure 5. The higher oil yield obtained for the freshwater microalgae at the 3% CO₂ concentration can be attributed to the trade-off of lower cell generation, and the lower oil yields obtained when the CO₂ concentration was increased to 9% is a result of increased cell division. Similarly, the variation in lipid content for the marine (*Tetraselmis suecica*) microalgae species can also be attributed to the variation in biomass yield caused by variation in CO₂ concentration and the tolerance of the species to the acidity caused by higher CO₂ concentrations. The results showed that from an economic stand point, the 3% CO₂ concentration is the most optimal condition for lipid accumulation in both species.

Similar results were reported in the literature. Widjaja et al. [36] noted that the microalgae species *Chlorella vulgaris* in lipid yields of 20%, 28% and 25% at the CO₂ concentration of 0, 0.33 and 0.83%, respectively. Huang and Su [53] noted that for the microalgae species *Chlorella vulgaris* grown using 0%, 15% and 50% CO₂ concentrations resulted in lipid yields of 34%, 35% and 36%, respectively. The findings are similar to those obtained in this study since they indicate that varying the CO₂ concentration does not significantly influence the lipid content. The variation in oil yield can be attributed to the varying cultivation periods, variation in nutrient systems and the effectiveness of the oil extraction methods used.

Effect of Carbon Source

Biomass

The cell yields shown in Table 6 for *Chlorella saccharophila* species which was obtained using NaHCO₃ as

Table 9. Analysis of the variance for oil yield using CO₂ as the carbon source.

Source	DF	SS	MS	F	P
Total	17	58.206			
Model					
Algae Species (S)	1	30.147	30.147	14.26	0.003
CO ₂ Concentration (C)	2	1.464	0.732	0.35	0.714
S*C	2	1.228	0.614	0.29	0.753
Error	12	25.367	2.114		

DF: Degree of freedom; SS: Sum of square; MS: Mean of square; R²= 56.42%

Table 10. Tukey’s grouping on oil yield by algae species and CO₂ concentration.

Factors	Level	N	Mean Yield	Tukey’s Grouping*
Species	Marine Water	9	1.590	A
	Freshwater (v/v)	9	4.178	B
CO ₂ :Air	3:97	6	3.164	A
	6:94	6	2.994	A
	9:91	6	2.493	A

*Groups with the same letter are not significantly different from each other at the 0.05 level.




Figure 7. Effect of microalgae type on the oil yield at varying CO₂ concentrations (CS-*Chlorella saccharophila*, TS- *Tetraselmis suecica*).






Figure 8. Effect of CO₂ concentration on the oil yield of freshwater and marine microalgae (CS-*Chlorella saccharophila*, TS-*Tetraselmis suecica*).

a carbon source (0.689x10⁶ cells/mL) were 74% higher than those achieved using 3% CO₂ (0.181x10⁶ cells/mL). The cell yield for *Tetraselmis suecica* species that resulted while using NaHCO₃ (0.750x10⁶ cells/mL) were 53% lower than those achieved using 3% CO₂ (1.148x10⁶ cells/mL). These results can be attributed to the cells ability to convert the carbon source into the preferred form for uptake and the abundance of the carbon source. The direct uptake of bicarbonate through an active transport system has only been noted in certain species [54]. In addition, some species have better extracellular carboanhydrase activities which allow them to convert the carbon source into different forms [54-56].



(a) Cell Yield



(b) Oil Yield

Figure 9. Effect of carbon source on the cell yield and oil yield of freshwater and marine microalgae (CS-*Chlorella saccharophila*, TS- *Tetraselmis suecica*).

Devgoswami et al. [46] studied the *Chlorella* microalgae species and noted biomass productivity of 82 and 189 mg/L/d using sodium bicarbonate and CO₂, as the carbon source, respectively. Moheimani [33] noted that the *Chlorella* sp. and *Tetraselmis suecica* grown using CO₂ as a carbon source resulted in biomass yields that were 6 and 23% higher than those obtained using NaHCO₃ as the carbon source. Goswami et al. [35] reported that the *Selenastrum* sp. grown using NaHCO₃ (20-100 mg/L) and CO₂ (4.4-8.2 g/L) resulted in biomass productivities in the range of 689-1102 mg/L/d and 667-889 mg/L/d, respectively. In this study, the biomass productivity achieved using NaHCO₃ and 3% CO₂ for *Chlorella saccharophila* was 106.8 mg/L/d and 28.1 mg/L/d, respectively. The biomass productivity achieved using NaHCO₃ and 3% CO₂ for *Tetraselmis suecica* was 116.3 mg/L/d and 177.9 mg/L/d, respectively. Variation in the values achieved in this study and those of the literature are attributed to the different species and different cultivation methods used.

Oil yield

The oil yields obtained with NaHCO₃ and 3% CO₂ as carbon sources are shown in Figure 9. The oil yield achieved

13. Alhattab, M. and A. Ghaly, 2014. Effects of light exposure and nitrogen source on the production of oil from freshwater and marine water microalgae. American Journal of Biochemistry and Biotechnology, In Press.
14. Lewis, L. A. 1997. Diversity and phylogenetic placement of *Bracteacoccus Tereg* (Chlorophyceae, Chlorophyta) based on 18S ribosomal RNA gene sequence data. J. Phycol. 33:279–85.
15. Bock, C., L. Krienitz and T. Proschold, 2011. Taxonomic reassessment of the genus *Chlorella* (Trebouxiophyceae) using molecular signatures (barcodes), including description of seven new species. Fottea: Journal of the Czech Psychological Society, 11(2): 293-312. ISSN: 1802-5439.
16. John, D.M., 2002. The freshwater algal flora of the British Isles: An identification Guide to freshwater and terrestrial algae. Cambridge University Press, 1:1890.
17. Singh, D., M. Puri, S. Wilkens, A.S. Mathur, D.K. Tuli and C.J. Barrow, 2013. Characterization of a new zeaxanthin producing strain of *Chlorella saccharophila* isolated from New Zealand marine waters. Bioresource Technology, 143:308-314. <http://dx.doi.org/10.1016/j.biortech.2013.06.006>
18. Matsuda, Y. and B. Colman, 1996. Active uptake of inorganic carbon by *Chlorella saccharophila* is not repressed by growth in high CO₂. Journal of Experimental Botany, 47(305): 1951-1956.
19. Acuna, J.L. and M. Kiefer, 2000. Functional response of the appendicularian *Oikopleura dioica*. Limnology and Oceanography, 45, 608–618. DOI:10.4319/lo.2000.45.3.0608
20. White, D.A., A. Pagarette, P. Rooks, S.T. Ali, 2012. The effect of sodium bicarbonate supplementation on growth and biochemical composition of marine microalgae cultures. Journal of applied Phycology, 25(1): 153-165. ISSN: 0921-8971
21. De Castro Araujo, S. and V.M.T. Garcia, 2005. Growth and biochemical composition of the diatom *Chaetoceros cf. wighamii* brightwell under different temperature, salinity and carbon dioxide levels. I. Protein, carbohydrates and lipids. Aquaculture, 246(1-4): 405-412. ISSN 0044-8486
22. Lavens, P. and P. Sorgeloos, 1996. Manual on the production and use of live food for aquaculture. FAO Fisheries Technical Paper. No. 361. Rome, FAO, pp. 295. ISBN: 92-5-103934-8
23. Guillard, R.R.L. and J.H. Ryther, 1962. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervaceae* (Cleve) Gran. Canadian Journal of Microbiology, 8, 229-239. DOI:10.1139/m62-029
24. Hughes, E. O., P. R. Gorham, and A. Zehnder, 1959. Toxicity of a unialgal culture of *Microcystis aeruginosa*. Canadian Journal of Microbiology, 4, 225. DOI:10.1139/m58-024
25. Araujo, G.S., L.J.B.L. Matos, J.O. Fernandes, S.J.M. Cartaxo, L.R.B. Goncalves, F.A.N. Fernandes and W.R.L. Farias, 2013. Extraction of lipids from microalgae by ultrasound application: Prospection of the optimal extraction method. Ultrasonics Sonochemistry, 20: 95-98. <http://dx.doi.org/10.1016/j.ultsonch.2012.07.027>
26. Isleten-Hosoglu, M., I. Gultepe and M. Elibol, 2012. Optimization of carbon and nitrogen sources for biomass and lipid production by *Chlorella saccharophila* under heterotrophic conditions and development of Nile red fluorescence based method for quantification of its neutral lipid content. Biochemical Engineering Journal, 61:11-19. DOI:10.1016/j.bej.2011.12.001
27. Herrera-Valencia, V., P.Y. Contreras-Pool, S.J. Lopez-Adrain, S. Peraza-Echeverria and L.F. Barahona-Perez, 2011. The green microalga *Chlorella saccharophila* as a suitable source of oil for biodiesel production. Current Microbiology, 63:151-157. DOI: 10.1007/s0028-011-9956-7
28. Bondioli, P., L. Della Bella, G. Rivolta, G. Chini Zittelli, N. Bassi, L. Rodolfi, D. Casini, M. Prussi, D. Chiaramonti, and M.R. Tredici, 2012. Oil production by the marine microalgae *Nannochloropsis sp.* F&M-M24 and *Tetraselmis suecica* F&M-M33. Bioresource Technology, 114: 567–672. DOI: 10.1016/j.biortech.2012.02.123
29. Danquah, M.K., R. Harun, R. Halim and G.M. Forde, 2010. Cultivation medium design via elemental balancing for *Tetraselmis suecica*. Chemical and Biochemical Engineering Quarterly, 24(3):361-369. ISSN:0352-9568
30. Michels, M.H.A., P.M. Slegers, M.H. Vermue and R.H. Wijffels, 2013. Effect of biomass concentration on the productivity of *Tetraselmis suecica* in a pilot-scale tubular photobioreactor using natural sunlight. Algal Research. <http://dx.doi.org/10.1016/j.algal.2013.11.011>
31. Chinnasamy, S., A. Bhatnagar, R.W. Hunt, K.C. Das, 2010. Microalgae cultivation in a wastewater dominated by carpet mill effluents for biofuel applications. Bioresource Technology, 101:3097–3105. DOI: 10.1016/j.biortech.2009.12.026
32. Hempel, N., I. Petrick and F. Behrendt, 2012. Biomass productivity and productivity of fatty acids and amino acids of microalgae strains as key characteristics of suitability for biodiesel production. Journal of Applied Phycology, 24:1407-1418. DOI: 10.1007/s10811-012-9795-3.
33. Moheimani, N.R., 2013. Inorganic carbon and pH effect on growth and lipid productivity of *Tetraselmis suecica* and *Chlorella sp.* (Chlorophyta) grown outdoors in bag photobioreactors. Journal of Applied Phycology, 25:387-398. ISSN: 1573-5176
34. Kaewkannetra, P., P. Enmak and T. Chiu, 2012. The effect of CO₂ and salinity on the cultivation of *Scenedesmus obliquus* for biodiesel production. Biotechnology and Bioprocess Engineering, 17:591-597. DOI 10.1007/s12257-011-0533-5
35. Goswami, R.C.D, N. Kalita and M.C. Kalita, 2012. A study on growth and carbon dioxide mitigation by microalgae *Selenastrum sp.* its growth behavior under

