

Anesthesia Management in a Patient with a History of Multidrug Anaphylaxis

Serdar Epözdemir*

*Department of Anesthesiology and Reanimation Medipol University Camlica Hospital, Bulgurlu Neighborhood Alemdag Street, Turkey.

Received July 30, 2021; Revised August 26, 2021; Accepted August 29, 2021

ABSTRACT

Perioperative anaphylaxis is a life-threatening systemic allergic reaction. Allergic reactions occurring during anesthesia were found to be between 1/6000-1/20000 in retrospective studies. The mortality rate is between 3-5%. The causes of perioperative anaphylaxis are neuromuscular blocking agents, antibiotics, latex, hypnotic induction agents, chlorhexidine, opioids, and colloids. Risk factors for perioperative anaphylaxis include female gender, mast cell disorders, allergic diseases (asthma, eczema, and hay fever), drug allergy, and atopy. We aimed to present the anesthetic management in our case that had to have knee replacement surgery, had co-morbidities and had a history of anaphylaxis against multiple drugs.

Keywords: Anaphylaxis, Knee replacement, Co-morbidities, Anesthesia management

Abbreviation: CSF: Cerebrospinal Fluid

INTRODUCTION

Approximately 260 million people in the world undergo surgery in a year. There is nothing more natural than the fact that these people of all races, genders, and different co-morbidities, who need surgery for different reasons, encounter anesthesia and are under various risks as a result. We anesthetists try to do our duty by minimizing the risks and feeling that we have to create a safe surgical process first and then a comfortable and aesthetic situation. Allergy and Anaphylaxis are two important perioperative issues that continue even when we start the day with us and finish our work in our daily practice [1-6].

CASE PRESENTATION

A 67-year-old female patient, height: 154 cm, weight: 105 kg, diagnosed with hypertension, obesity, hypothyroidism. Preoperative evaluation for left knee prosthesis surgery was performed by us. Chest X-ray, electrocardiography, hemogram, biochemical analysis, thyroid function test: Normal. In her anamnesis, she had a history of angioedema, fainting, burning in the stomach, facial tingling, glottis edema after the use of levofloxacin, radiocontrast agent and meloxicam at different times. Recommendation of department of pulmonology, immunology and allergy diseases, ASA III patient, who was administered 40 mg prednol 1x1 7 h before surgery, and 40 mg Avil 1x1 1 h before surgery, was administered as premedication 3 mg dormicum as an intravenous infusion in 100 ml saline before the operation. He was monitored after he was taken to the operating room, vital signs were recorded as BP: 160/88, heart rate: 102

beats/min, SpO₂: 95%. Regional anesthesia preparations were made, he was brought to a sitting position, and the intrathecal interval was approached with a gray spinal needle (27 Gauge) from the L 3-4 interval by following the rules of asepsis-antisepsis. After observing the arrival of CSF fluid, 10 mg of Heavy Marcaine was administered, it was kept in the left lateral position for a while, anesthesia Control was done and surgery was started. Preoperatively, 80 mg of prednol, 8 mg of ondansetron, and 2 g of iespor as prophylactic antibiotic were administered intravenously in 100 ml of physiological saline. The entire 100 mg contramal IV infusion was administered to the patient without any problems. Contramal IV PCA was prepared for postoperative analgesia. No complications and anaphylactoid reactions were observed during this period.

DISCUSSION

Ensuring the safety of patients in the perioperative period is the most important responsibility of the anesthetist. During this period, many events, allergies and anaphylactic reactions, which are related or not related to anesthetic methods and

Corresponding author: Serdar Epözdemir, Department of Anesthesiology and Reanimation Medipol University Camlica Hospital, Bulgurlu Neighborhood Alemdag Street, Turkey, Tel: +902166813030, E-mail: epzdemir@yahoo.com

Citation: Epözdemir S. (2022) Anesthesia Management in a Patient with a History of Multidrug Anaphylaxis. Int J Anaesth Res, 5(2): 178-179.

Copyright: ©2022 Epözdemir S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

agents, can be seen. Although the frequency of these reactions varies according to geographical regions (1/10,000 to 1/30,000), they may result in severe cardiovascular collapse, bronchospasm, and severe skin reactions [7,8].

In an international study conducted by Kroigaard [9] it was reported that an allergic reaction due to nondepolarizing muscle relaxants followed by latex and antibiotic allergies developed frequently during anesthesia. Latex allergy can occur as Type I (anaphylactic type) and Type IV (late type). Type I reactions may occur after 30-60 min, while Type IV reactions may occur after 24-48h. Latex-containing products are frequently used in the health sector. This can be life-threatening for people with a latex allergy. Allergy history should be questioned in the preoperative period. Findings such as urticaria, rhinitis, respiratory distress, facial edema due to exposure to these substances should raise suspicion for latex allergy. In our patient, there was a history of allergic reaction that developed 3 times in the perioperative period. Latex allergy was detected in the later allergy test. Although it has been shown that preoperative pharmacological prophylaxis does not prevent the occurrence of anaphylaxis [10], we applied H1-H2 antagonists and steroids in order to prevent or reduce possible reactions related to nonspecific histamine release [8].

CONCLUSION

We think that complications can be prevented by making the preoperative preparations of patients with a history of anaphylaxis thoroughly with a good anamnesis, choosing non-allergenic drugs in the intraoperative period and optimal use of analgesic drugs for the postoperative period.

REFERENCES

1. Lobera T, Audicana MT, Pozo MD, Blasco A, Fernandez E, et al. (2008) Study of hypersensitivity reactions and anaphylaxis during anesthesia in Spain. *J Investig Allergol Clin Immunol* 18: 350-356.
2. Mertes PM, Laxenaire MC, Alla F (2003) Anaphylactic and anaphylactoid reactions occurring during anesthesia in France in 1999-2000. *Anesthesiology* 99: 536-545.
3. Chong YY, Caballero MR, Lukawska J, Dugue P (2008) Anaphylaxis during general anesthesia: One-year survey from a British allergy clinic. *Singapore Med J* 49: 483-487.
4. Mertes PM, Laxenaire MC, Alla F (1981) Groupe d'Etudes des Reactions Anaphylactodes Peranesthésiques. Anaphylactic and anaphylactoid reactions occurring during anesthesia interaction with compound 48/80. *Agents Actions* 11: 196-203.
5. Nel L, Eren E (2011) Peri-operative anaphylaxis. *Br J Clin Pharmacol* 71: 647-658.
6. Grosman N (1981) Histamine release from isolated rat mast cells: Effect of morphine and related drugs and their interaction with compound 48/80. *Agents Actions* 11: 196-203
7. Laxenaire MC, Mertes PM (2001) Anaphylaxis during anesthesia. Results of a two-year survey in France. *Br J Anaesth* 87: 549-558.
8. Mertes PM, Laxenaire MC (2002) Allergic reactions occurring during anesthesia. *Eur J Anaesthesiol* 19: 240-262.
9. Kroigaard M, Garvey LH, Gillberg L (2007) Scandinavian Clinical Practice Guidelines on the diagnosis, management and follow-up of anaphylaxis during anesthesia. *Acta Anaesthesiol Scand* 51: 655-670.
10. Setlock MA, Cotter TP, Rosner D (1993) Latex allergy: Failure of prophylaxis to prevent severe reaction. *Anesth Analg*. 76: 650-652.