









correlated with many cancer types including breast cancer especially in the metastasis stage [32]. Results of docking are elucidated in **Table 3**.


**Table 3.** PatchDock as well as FireDock results of the best anticancer peptides.

Peptides	PatchDock			FireDock				
	Score	Area	ACE	Global Energy	Attractive VdW	Repulsive VdW	ACE	HB
EGFR								
Peptide 5	7600	992.4	-122.52	-19.35	-22.85	25.33	0.22	-2.13
Peptide 7	6794	831.8	-126.24	-56.01	-24.31	5.60	-12.86	-1.25
Peptide 10	7726	918.1	-43.34	-10.70	-22.41	9.27	11.81	-4.82
ER- $\alpha$								
Peptide 5	7834	999.2	-275.33	-63.55	-27.06	9.59	-17.47	-0.25
Peptide 7	7200	964.3	-420.73	-23.91	-29.78	72.17	-21.71	-1.8
Peptide 10	7022	945.7	-449.44	-63.35	-23.21	7.73	-20.42	-0.35
MMP-3								
Peptide 5	6330	867.4	-337.62	-52.11	-25.57	10.42	-9.55	-1.93
Peptide 7	6220	695	-58.32	-38.91	-23.04	4.12	-4.92	-3.42
Peptide 10	6542	803.2	-181.46	-38.16	-19.16	8.64	-9.15	-0.34

ACE: Atomic Contact Energy; VdW: Van der Waal; HB: Hydrogen Bonds

As shown in **Table 3**, peptide 7 greatly blocks EGFR with a global energy of -56.01 kcal/mole while the remaining peptides are not good enough for EGFR inhibition. With respect to ER- $\alpha$ , peptide 5 as well as peptide 10 are superior inhibitors with a global energy -63.55 and -63.35 kcal/mole. Of the top 3 peptides, peptide 5 are good candidate for inhibition of MMP-3 (global energy -52.11 kcal/mole) whilst peptides 7 and 10 had a global energy of -38.91 and -38.16 kcal/mole. Collectively, peptide 7 is a good candidate for blockade of EGFR whereas peptide 5 can be used as

inhibitor of both ER- $\alpha$  and, along with peptide 10, MMP-3. Peptide-protein interactions of best poses are shown in **Figure 2**. These data suggest that the studied peptides fragmented from *N.sativa* can be utilized as ACP against breast cancer receptors. Surprisingly, peptide 7 and peptide 5 are even better than the reference inhibitors of EGFR and ER- $\alpha$  (Erlotinib and TAM) in terms of global energy estimated by FireDock (-50.90 and -60.2 kcal/mole; data not shown).



**Figure 2.** Peptide-protein docking 3D interaction visualized by discovery studio 2021. Docking results of peptide 7 against EGFR (A), peptide 5 against ER- $\alpha$  (B), and peptide 5 against MMP-3 (C).

EGFR interacted with peptide 7 through VdW interactions imposed by two lysine residues within the active site pocket and one arginine residue through H-bonds. Similarly, Asp 351 and Met 528 formed H-bonds with peptide 5 and the rest interacting residues interacted via VdW forces with ER- $\alpha$ . Regarding MMP-3, peptide 5 formed electrostatic attractions with Asp 111 (Figure 2).

Accordingly, the present work demonstrates *in-silico* the anti-cancer activity of the peptides fragmented from some *N.sativa* proteins as predicted by ACPred and mACPred webservers and then validated by peptide-protein docking through PatchDock-FireDock platform. This candidates *N.sativa* as a superior source as therapeutic nutraceutical option against breast cancer theoretically.

## CONCLUSION

In conclusion, 6 *N.sativa* proteins after proteolysis using trypsin enzyme gave rise to 23 peptides for which physico-chemical properties were calculated. 11 of which had ACP bioactivity as predicted by mACPred and ACPred platforms. Among them, peptides 5 (HGSCNYK), peptide 7 (CICYEC), and peptide 10 (TCSGLCGCK) were the best in terms of the possibility score as ACP against 3 types of breast cancer receptors, EGFR, ER- $\alpha$  and MMP-3 as demonstrated by PatchDock and FireDock servers which necessitate *in vitro* assays confirmation. However, further analysis of the top peptides against wide array of other cancer types should be addressed. Also, the peptides with low anti-cancer activity should be assessed for different bioactivities.

## REFERENCES

- Khan MA, Chen H, Tania M, Zhang D (2011) Anticancer activities of *Nigella sativa* (black cumin). Afr J Tradit Complement Altern Med 8: 226-232.
- Majdalawieh AF, Fayyad MW, Nasrallah GK (2017) Anti-cancer properties and mechanisms of action of thymoquinone, the major active ingredient of *Nigella sativa*. Crit Rev Food Sci Nutr 57: 3911-3928.
- Ahmad A, Husain A, Mujeeb M, Khan SA, Najmi AK, et al. (2013) A review on therapeutic potential of *Nigella sativa*: A miracle herb. Asian Pac J Trop Biomed 3: 337-352.
- Butt MS, Sultan MT (2010) *Nigella sativa*: Reduces the risk of various maladies. Crit Rev Food Sci Nutr 50: 654-665.
- Gali-Muhtasib H, Roessner A, Schneider-Stock R (2006) Thymoquinone: A promising anti-cancer drug from natural sources. Int J Biochem Cell Biol 38: 1249-1253.
- Nickavar B, Mojab F, Javidnia K, Amoli MAR (2003) Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. Zeitschrift Für Naturforschung C 58: 629-631.
- Salem ML (2005) Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed. Int Immunopharmacol 5: 1749-1770.
- Majdalawieh AF, Fayyad MW (2016) Recent advances on the anti-cancer properties of *Nigella sativa*, a widely used food additive. J Ayurveda Integr Med 7: 173-180.
- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, et al. (2021) Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 71: 209-249.
- Rizvi M, Alam MS, Mehdi SJ, Ali A, Batra S (2012) Allelic loss of 10q23. 3, the PTEN gene locus in cervical carcinoma from Northern Indian population. Pathol Oncol Res 18: 309-313.
- Rahmani AH, Alzohairy M, Babiker AYY, Khan AA, Aly SM, et al. (2013) Implication of androgen receptor in urinary bladder cancer: A critical mini review. Int J Mol Epidemiol Genet 4: 150.
- Jo Y, Choi N, Kim K, Koo H-J, Choi J, et al. (2018) Chemoresistance of cancer cells: Requirements of tumor microenvironment-mimicking *in vitro* models in anti-cancer drug development. Theranostics 8: 5259.
- Lee Y-W, Chen T-L, Shih Y-RV, Tsai C-L, Chang C-C, et al. (2014) Adjunctive traditional Chinese medicine therapy improves survival in patients with advanced breast cancer: A population-based study. Cancer 120: 1338-1344.
- Rahmani AH, Alzohairy MA, Khan MA, Aly SM (2014) Therapeutic implications of black seed and its constituent thymoquinone in the prevention of cancer through inactivation and activation of molecular pathways. Evid Based Complement Alternat Med 2014: 724658.
- Tyagi A, Tuknait A, Anand P, Gupta S, Sharma M, et al. (2015) CancerPPD: A database of anticancer peptides and proteins. Nucleic Acids Res 43: D837-D843.
- Thundimadathil J (2012) Cancer treatment using peptides: Current therapies and future prospects. J Amino Acids 2012: 967347.
- Karami M, Chaleshgar M, Salari N, Akbari H, Mohammadi M (2022) Global Prevalence of Anemia in Pregnant Women: A Comprehensive Systematic Review and Meta-Analysis. Matern Child Health J 26(7): 1473-1478.
- Chiangjong W, Chutipongtanate S, Hongeng S (2020) Anticancer peptide: Physicochemical property,

- functional aspect and trend in clinical application (Review). *Int J Oncol* 57: 678-696.
19. Consortium U (2019) UniProt: A worldwide hub of protein knowledge. *Nucleic Acids Res* 47: D506-515.
  20. Gasteiger E, Hoogland C, Gattiker A, Wilkins MR, Appel RD, et al. (2005) Protein identification and analysis tools on the ExPASy server. *The Proteomics Protocols Handbook 2005*: 571-607.
  21. Garg VK, Avashthi H, Tiwari A, Jain PA, Ramkete PW, et al. MFPPI-multi FASTA ProtParam interface. *Bioinformation* 12: 74.
  22. Boopathi V, Subramaniam S, Malik A, Lee G, Manavalan B, et al. (2019) mACPpred: A support vector machine-based meta-predictor for identification of anticancer peptides. *Int J Mol Sci* 20: 1964.
  23. Schaduangrat N, Nantasenamat C, Prachayasittikul V, Shoombuatong W (2019) ACPred: A computational tool for the prediction and analysis of anticancer peptides. *Molecules* 24: 1973.
  24. Lamiable A, Thévenet P, Rey J, Vavrusa M, Derreumaux P, et al. (2016) PEP-FOLD3: Faster de novo structure prediction for linear peptides in solution and in complex. *Nucleic Acids Res* 44: W449-W454.
  25. Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ (2005) PatchDock and SymmDock: Servers for rigid and symmetric docking. *Nucleic Acids Res* 33: W363-W367.
  26. Andrusier N, Nussinov R, Wolfson HJ (2007) FireDock: Fast interaction refinement in molecular docking. *Proteins* 69: 139-159.
  27. Sutradhar KB, Amin M (2014) Nanotechnology in cancer drug delivery and selective targeting. *Int Sch Res Notices* 939378: 1-12.
  28. Hayashi MA, Ducancel F, Konno K (2012) Natural peptides with potential applications in drug development, diagnosis, and/or biotechnology. *Int J Pept* 2012: 757838.
  29. Grisoni F, Neuhaus CS, Gabernet G, Müller AT, Hiss JA, et al. (2018) Designing anticancer peptides by constructive machine learning. *ChemMedChem* 13: 1300-1302.
  30. Liu Y, Ma H, Yao J (2020) ER $\alpha$ , A Key Target for Cancer Therapy: A Review. *Onco Targets Ther* 13: 2183-2191.
  31. Sigismund S, Avanzato D, Lanzetti L (2018) Emerging functions of the EGFR in cancer. *Mol Oncol* 12: 3-20.
  32. Suhaimi SA, Chan SC, Rosli R (2020) Matrix Metalloproteinase 3 Polymorphisms: Emerging genetic Markers in Human Breast Cancer Metastasis. *J Breast Cancer* 23: 1-9.