

glaucoma and ocular hypertension [6]. Astigmatism is a type of refractive error where rays of light form line focus image at different meridians instead of point focus. A previous study found that patients with with-the-rule astigmatism having purely corneal astigmatism over 4D resulted in an under estimate of intraocular pressure and against the rule astigmatism, an overestimated of intraocular pressure by 1 mm of Hg, but the etiology was still unknown [7].

Furthermore, blood pressure and its co-relation with intraocular pressure in patients with refractive errors are of clinical interest. Blood pressure is the pressure of circulating blood on the walls of blood vessels. It is expressed in terms of the systolic pressure over diastolic pressure and is measured in millimeters of mercury (mm Hg). Normal resting blood pressure in an adult is approximately 120 mm of mercury systolic and 80 mm of mercury diastolic abbreviated as "120/80 mm Hg". There is correlative not necessarily causal, relationship between glaucoma and systemic hypertension and the risk of developing systemic hypertension in hypermetropic patients is higher compared to myopia [8]. Long standing hypertension may cause micro vascular damage whereas low systemic blood pressure may reduce local perfusion, particularly in the present of existing intraocular pressure elevation or poor auto regulation [9].

Therefore, this study would help us to know whether the change in refractive status of eye may have any co-relation between IOP and blood pressure or not.

MATERIALS AND METHODOLOGY

The experimental design of this study was hospital-based, observational prospective and crosses-sectional. The study sample consisted of patients, between 18 to 60 years of age. Permission was obtained from ethical committee of the institute. An informed consent was obtained from patients willing to participate in the study. Patients with ocular diseases, systemic diseases like Diabetes, asthma, COPD, pathological myopia >-10.00 D, pathological hypermetropia $>+10.00$ D and those under any medications except for controlled hypertension were excluded.

All patients underwent baseline examination to rule out any abnormality in the eyes. A detailed history was taken as per

the proforma. Unaided, aided and pinhole visual acuity was recorded for distance at 6 m and near at 33 cm with Snellen's distance and near visual acuity chart. Refraction was performed objectively and subjectively with undilated and dilated state using GR100 Grand Seiko Autorefractometer (Japan). Refractive error was stratified into three categories: Low myopia (≤ -3.00 D), moderate myopia (-3.00 D to -6.00 D), high myopia (>-6.00 D), low hypermetropia ($\leq +2.00$ D), moderate hypermetropia ($+2.00$ D to $+5.00$ D), high hypermetropia ($>+5.00$ D). Astigmatism was defined by cylinder at least ± 0.50 DC.

Slit lamp evaluation was done to rule out any anterior segment abnormality. Non-contact tonometry by using Keeler's non-contact tonometer was performed on both eyes, with the right eye measured first, between 10.00 AM to 3.00 PM. Measurements adhered to protocols; the patient was seated, with eyes in the primary position, eyelashes or eyelids did not obscure the applanation circle and the alignment spot was centered and focused in the alignment circle before applanation. Detailed examination of the posterior segment was carried out with direct ophthalmoscope by using $+20.00$ D lens. Then, systolic and diastolic blood pressure was measured using a standard mercury sphygmomanometer after 5 min of rest in the sitting position.

RESULTS

In this study, 344 eyes of 180 patients (36 simple myopic, 68 simple hyperopic, 75 simple astigmatic, 117 compound astigmatic and 48 myopic astigmatic) were examined of which 91 (51%) were male and 89 (49%) were female. Following assessments were performed to find out the correlation between IOP, refractive errors and blood pressure. **Tables 1 and 2** showed comparison between refractive errors and blood pressure. By using ANOVA test, P -value was <0.05 therefore there was statistically significant co-relationship found between systolic blood pressure and refractive errors and no co-relation found between refractive errors and diastolic blood pressure. Compound astigmatism group of patients were likely to have lesser systolic blood pressure (**Figure 1**) compared to other types of refractive errors.

Table 1. Correlation between refractive errors and systolic blood pressure.

Refractive Error Group	Number of eyes	Systolic Blood Pressure		P-value
		Mean	SD	
Myopia	36	120.00	15.31	0.003
Hyperopia	68	123.24	14.50	
Simple Astigmatism	75	120.53	14.69	
Compound Astigmatism	117	117.18	11.44	
Mixed Astigmatism	48	125.21	11.67	

Table 2. Correlation between refractive errors and diastolic blood pressure.

Refractive Error Group	Number of eyes	Diastolic Blood Pressure		P-value
		Mean	SD	
Myopia	36	85.83	13.55	0.283
Hyperopia	68	84.97	10.65	
Simple Astigmatism	75	85.60	12.38	
Compound Astigmatism	117	82.99	9.63	
Mixed Astigmatism	48	86.46	9.34	




Figure 1. Mean systolic blood pressure and diastolic blood pressure with refractive error type.

By using Pearson co-relation coefficient test, **Tables 3 and 4** showed no statistically co-relation between intraocular pressure and systolic blood pressure in patients with simple myopia, simple hyperopia, simple astigmatism and compound astigmatism but poor positive co-relation found between diastolic blood pressure and intraocular pressure in


patients with mixed astigmatism ($P=0.023$). Slight increase in systolic blood pressure was observed in patients with mixed astigmatism compared to other refractive errors. Likewise slight increase in diastolic pressure was observed in patients with mixed astigmatism which did not show any clinical significance (**Figure 2**).

Table 3. Co-relation between refractive errors and intraocular pressure.

Refractive Error Group	Number of eyes	IOP		P-value
		Mean	SD	
Myopia	36	16.22	3.18	0.301
Hyperopia	68	16.06	3.50	
Simple Astigmatism	75	15.23	3.27	
Compound Astigmatism	117	15.68	3.05	
Mixed Astigmatism	48	16.38	3.72	

Table 4. Co-relation between intraocular pressure and blood pressure with respect to refractive errors.

	Number of patients	Pearson correlation coefficient (r) between	
		SBP vs. IOP	DBP vs. IOP
Myopia	36	0.065	-0.12
P-value		0.708	0.404
Hyperopia	68	-0.033	0.006
P-value		0.788	0.958
SA	75	0.009	0.02
P-value		0.941	0.865
CA	117	0.162	0.108
P-value		0.082	0.245
MA	48	0.047	0.327
P-value		0.75	0.023*

**Figure 2.** Mean IOP with refractive error type.

DISCUSSION

In this report we presented a detailed study of intraocular pressure, refractive errors and blood pressure in an 18 to 60 years old population. We took this age group as the period between these ages is one of the refractive stability. Additionally, we studied correlation between refractive errors and blood pressure which has not been discussed in any study.

Refractive errors are very common ocular problems in the population worldwide. Myopia is one of the risk factors for glaucoma that is commonly mentioned in various studies [10,11]. Study done by Abdalla et al. [10] showed high IOP in myopic individuals compared to non-myopic individuals suggesting that relationship between glaucoma and myopia may be pressure mediated. The possible mechanism for raised intraocular pressure in myopic is the shearing forces exerted by scleral tension across the lamina cribrosa and may be important in pathogenesis of pressure damage [10].

However study done by Lee et al. [12] in Singapore, assessed association between intraocular pressure and refractive errors in children of age group 9-11 years. Refractive errors were categorized into 4 groups: hypermetropia, low myopia, emmetropia and high myopia. The result of this study showed no significant intraocular pressure differences between different types of refractive errors. They said that elevated intraocular pressure and myopia might not be linked in these children. Our study also did not provide sufficient evidence to conclude that intraocular pressure was related to refractive errors, but slight increase in intraocular pressure was observed in mixed astigmatic patients compared to other types of refractive errors in older age groups. Different methods and samples may be the reason for the inconsistent result with different studies.

Our results compared to Keihanian et al. study [13], by showing significant correlation between changes of systolic

blood pressure and intraocular pressure ($P=0.005$), while the relationship between diastolic blood pressure and intraocular pressure changes were not seen ($P>0.005$). But result of our study indicated a direct relation between systolic blood pressure and refractive errors, as the systolic blood pressure showed a statistically significant result but no clinically significant result in those patients and only weak or poor positive co-relation found between diastolic blood pressure and intraocular pressure with respect to mixed astigmatism [14].

No significant correlation was found in the measured parameters with age or sex. A large sample size, high refractive error group and measurement of corneal biomechanical parameters would have added to the study's strengths.

CONCLUSION

In summary, our study had shown weak co-relation between refractive errors, blood pressure and intraocular pressure which did not show any clinical significance. The observations need confirmation by study with age matched emmetropes as controls.

REFERENCES

- Pan CW, Ramamurthy D, Saw SM (2012) Worldwide prevalence and risk factors for myopia. *Ophthalmic Physiol Opt* 32: 3-16
- Haskemi H, Khabazkhoob M, Emamian MH, Shariati M, Yekta A, et al. (2016) Distribution of intraocular pressure and its determinants in an Iranian adult population. *Int J Ophthalmol* 9: 1207-1214.
- Manny R, Mitchell LG, Cotter SA, Jones-Jordan LA, Kleinstein RN, et al. (2011) Intraocular pressure, ethnicity and refractive error. *Optom Vis Sic* 88: 1445-1453.
- Quigley HA (1987) Reappraisal of the mechanism of glaucomatous optic nerve damage. *Eye* 1: 318-322.
- Wong TY, Wang JJ, Rochtchina E, Klein R, Mitchell P (2004) Does refractive error influence the association of blood pressure and retinal vessel diameters? The Blue Mountains Eye Study. *Am J Ophthalmol* 137: 1050-1055.
- Shen L, Melles RB, Metlapally R, Barcellos L, Schaefer C, et al. (2016) The association of refractive error with glaucoma in a multiethnic population. *Ophthalmology* 123: 92-101.
- Boyd K, Edmond JC (2018) Astigmatism. *American Academy of Ophthalmology. Eye Smart® Eye health.*
- Yassin AS, Elham R, Tamimi AL (2016) Age, gender and refractive error association with IOP in healthy Saudi participants. *Saudi J Ophthalmol* 30: 44-48.
- Choi AJ, Han K, Yong-MP, Chan PK (2014) Age related association of refractive error with IOP in the Korea. National Health and Nutrition Examination survey. *PLoS One* 9: e111879.
- Abdullah MI, Hamdi M (1970) Applanation ocular tension in myopia and emmetropia. *Br J Ophthalmol* 54: 122-125.
- Marcus MW, de Vries MM, Junoy Montolio FG, Jansonius NM (2011) Myopia as a risk factor for open angle glaucoma: A systematic review and meta-analysis. *Ophthalmology* 118: 1989-1994 e2.
- Lee AJ, Saw SM, Gazzard G, Cheng A, Tan DTH. (2004) Intraocular pressure associations with refractive error and axial length in children. *Br J Ophthalmol* 88: 5-7.
- Keihanian F, Saeidinia A, Tahere M, Keihanian F, Akbari M, et al. (2016) Relationship between intraocular pressure and some risk factors in Northern Iran. *Int J Med Res Health Sci* 4: 104-110.
- Dusek WA, Pierscionek BK, McClelland JF (2012) Age variation in intraocular pressure in a cohort of healthy Austrian school children. *Eye* 26: 841-845.