

Gender Based Violence (GBV) in Emergency Medicine

Tiziana Ciarambino*, Gennaro Sansone and Mauro Giordano

*Department of Internal Medicine, Hospital of Marcanise, ASL Caserta, Italy.

Received March 15, 2019; Accepted March 27, 2019; Published July 12, 2019

ABSTRACT

Gender Based Violence (GBV) is any act ‘that results in, or is likely to result in, physical, sexual or psychological harm or suffering’ that is directed against a person because of their biological sex, gender identity or perceived adherence to socially defined norms of masculinity and femininity. GBV includes intimate partner violence and can be physical, sexual, emotional, economic or structural where that violence targets someone because of their gender or non-compliance with gender norms. The aim of the guidelines is to provide an adequate and integrated intervention in the treatment of the physical and psychological consequences that in particular male violence produces on women's health.

Keywords: Gender based violence, Woman health

INTRODUCTION

Gender Based Violence (GBV) is any act ‘that results in, or is likely to result in, physical, sexual or psychological harm or suffering’ that is directed against a person because of their biological sex, gender identity or perceived adherence to socially defined norms of masculinity and femininity. GBV includes intimate partner violence and can be physical, sexual, emotional, economic or structural where that violence targets someone because of their gender or non-compliance with gender norms. It can be experienced by women and girls, men and boys, and transgender and intersex people of all ages and has direct consequences on health, social, financial and other aspects of their lives [1].

GBV takes on many forms and can occur throughout the lifecycle, from the prenatal phase through childhood and adolescence, the reproductive years and old age [2].

Types of GBV

The WHO divided GBV in two groups:

- Gender-based violence:
 - Rape by strangers
 - Female genital mutilation
 - Sexual harassment in the workplace
 - Selective malnutrition of girls
- Domestic/Family violence:
 - Child abuse
 - Elder abuse

Intimate partner violence and sexual abuse of women and girls in the family are overlapped in these two groups [3].

Epidemiologies

The lifetime prevalence of rape by an intimate partner was an estimated 8.8% for women and an estimated 0.5% for men (**Figure 1**). An estimated 15.8% of women and 9.5% of men experienced other forms of sexual violence by an intimate partner during their lifetimes. Severe physical violence by an intimate partner was experienced by an estimated 22.3% of women and 14.0% of men during their lifetimes. Among female victims of completed rape, an estimated 78.7% were first raped before age 25 years, with 40.4% experiencing rape before age 18 years. Among male victims who were made to penetrate a perpetrator, an estimated 71.0% were victimized before age 25 years and an estimated 21.3% were victimized before age 18 years (**Figure 2**) [4].

Corresponding author: Tiziana Ciarambino, Department of Internal Medicine, Hospital of Marcanise ASL Caserta Italy, E-mail: tiziana.ciarambino@gmail.com

Citation: Ciarambino T, Sansone G & Giordano M. (2019) Gender Based Violence (GBV) in Emergency Medicine. J Pharm Drug Res, 2(4): 134-139.

Copyright: ©2019 Ciarambino T, Sansone G & Giordano M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. In this figure we reported the prevalence of GBV for women and for men.




Figure 2. In this figure we reported the prevalence of GBV for women and for men by age.

In one cross sectional study, 11.7% of women visiting a variety of American emergency departments were there because of acute injury or stress related to domestic violence. The lifetime prevalence rate for domestic violence of 54.2% in women attending Emergency Departments (ED) [5]

In a study population 12.6% of men attending emergency departments were victims of domestic violence by a current or former female intimate partner [6]. These data demonstrate how important GBV is to an emergency physician.

Risk factors

Individual factors:

- Young age
- Heavy drinking
- Depression
- Personality disorders
- Low academic achievement
- Low income
- Witnessing or experiencing violence as a child

Relationship factors:

- Marital conflict
- Marital instability
- Male dominance in the family
- Economic stress
- Poor family functioning

Community factors:

- Weak community sanctions against domestic violence
- Poverty
- Low social capital

Social factors:

- Traditional gender norms
- Social norms supportive of violence [7-10]

Health consequences

GBV has both physical and psychological consequences.

Physical:

- Headaches

- Back pain and other musculoskeletal pain
- Chest pain
- Gynecological disorders including menstrual disorders, pelvic pain and dyspareunia
- Sexually transmitted diseases, including HIV/AIDS
- Gastrointestinal disorders
- Urinary tract infection
- Acute respiratory infection

Psychological:

- Anxiety
- Substance use
- Tobacco use
- Family and social problems
- Depression [11]

MANAGEMENT IN ED

An American study identified that emergency department use was common in the two years before murder by a partner. In USA, a spouse or intimate partner perpetrated 33.6% of female homicides, while less than 8% of murders of men were perpetrated by a spouse or intimate partner [5].

a. How to detect GBV?

It seems that history taking and clinical examination is unsatisfactory for diagnosing domestic violence. While the evidence does not support screening for domestic violence, it is still important that emergency physicians know how to create the opportunity of a patient disclosing domestic violence, so that self-reported victims of domestic violence can be offered help. Ideally all consultations should take place in a private room with, initially, only the patient and the doctor. Simple, direct, non-judgmental questions are the best way to inquire about domestic violence if this is felt appropriate. The partner violence scale (PVS) consists of three questions (Have you been hit, kicked, punched or otherwise hurt by someone within the past year? Do you feel safe in your current relationship now? Is there a partner from a previous relationship who is making you feel unsafe now?) That have been compared with both the index spouse abuse (ISA) and the conflict tactics scale (CTS), two validated measures of domestic violence. The first question proved to be nearly as useful as all three questions at identifying victims. The sensitivity of the PVS compared with the ISA was 65%; specificity was 80% with a positive predictive value of 51% [5].

b. Management of an identified victim

As always the priority is to treat the physical injury. It should be explained to the victim that domestic violence is

unacceptable and against the law. Police contact should be offered while in the comparative safety of the emergency department. It is much easier to contact the police from the emergency department than a household that is shared with the perpetrator [5].

SPECIAL CASES

Elder abuse

The definition developed by Action on Elder Abuse in the United Kingdom and adopted by the International Network for the Prevention of Elder Abuse states that: "Elder abuse is a single or repeated act or lack of appropriate action, occurring within any relationship where there is an expectation of trust which causes harm or distress to an older person." Such abuse is generally divided into the following categories:

- Physical abuse – the infliction of pain or injury, physical coercion or physical or drug induced restraint.
- Psychological or emotional abuse – the infliction of mental anguish.
- Financial or material abuse – the illegal or improper exploitation or use of funds or resources of the older person.
- Sexual abuse – non-consensual sexual contact of any kind with the older person.
- Neglect – the refusal or failure to fulfil a caregiving obligation. This may or may not involve a conscious and intentional attempt to inflict physical or emotional distress on the older person.

The rate of abuse is 4-6% among older people if physical, psychological and financial abuse and neglect are all included. For older people, the consequences of abuse can be especially serious. Older people are physically weaker and more vulnerable than younger adults, their bones are more brittle and convalescence takes longer. Even a relatively minor injury can cause serious and permanent damage. Many older people survive on limited incomes, so that the loss of even a small sum of money can have a significant impact. They may be isolated, lonely or troubled by illness, in which case they are more vulnerable as targets for fraudulent schemes.

The medical profession has played a leading role in raising public concern about elder abuse. While it may be thought that doctors are best placed to notice cases of abuse – partly because of the trust that most elderly people have in them – many doctors do not diagnose abuse because it is not part of their formal or professional training and hence does not feature in their list of differential diagnoses. Most emergency departments do not use protocols to detect and deal with elder abuse, and rarely attempt to address the

mental health or behavioral signs of elder abuse, such as depression, attempted suicide, or drug or alcohol abuse [7].

There should be an investigation of a patient's condition for possible abuse if a doctor or other health care worker notices any of the following signs:

- Delays between injuries or illness and seeking medical attention;
- Implausible or vague explanations for injuries or ill-health, from either the patient or his or her caregiver;
- Differing case histories from the patient and the caregiver;
- Frequent visits to emergency departments because a chronic condition has worsened, despite a care plan and resources to deal with this in the home;
- Functionally impaired older patients who arrive without their main caregivers;
- Laboratory findings that are inconsistent with the history provided.

When conducting an examination the doctor or health care worker should:

- Interview the patient alone, asking directly about possible physical violence, restraints or neglect;
- Interview the suspected abuser alone;
- Pay close attention to the relationship between and the behavior of, the patient and his or her suspected abuser;
- conduct a comprehensive geriatric assessment of the patient, including medical, functional, cognitive and social factors;
- Document the patient's social networks, both formal and informal [12].

Child abuse

The WHO Consultation on Child Abuse Prevention drafted the following definition [8]: "Child abuse or maltreatment constitutes all forms of physical and/or emotional ill-treatment, sexual abuse neglect or negligent treatment or commercial or other exploitation, resulting in actual or potential harm to the child's health, survival, development or dignity in the context of a relationship of responsibility, trust or power" [13].

There are four types of child maltreatment by caregivers, namely:

- Physical abuse;
- Sexual abuse;
- Emotional abuse;

- Neglect

According to the World Health Organization, there were an estimated 57 000 deaths attributed to homicide among children under 15 years of age in 2000. Global estimates of child homicide suggest that infants and very young children are at greatest risk, with rates for the 0-4 year old age group more than double those of 5-14 year olds. World for children under 5 years of age living in high-income countries, the rate of homicide is 2.2 per 100 000 for boys and 1.8 per 100 000 for girls. In low- to middle-income countries the rates are 2-3 times higher – 6.1 per 100 000 for boys and 5.1 per 100 000 for girls [7]. Ill health caused by child abuse forms a significant portion of the global burden of disease. Importantly, there is now evidence that major adult forms of illness – including ischemic heart disease, cancer, chronic lung disease, irritable bowel syndrome and fibromyalgia – are related to experiences of abuse during childhood. The apparent mechanism to explain these results is the adoption of behavioral risk factors such as smoking, alcohol abuse, poor diet and lack of exercise. Research has also highlighted important direct acute and long-term consequences [14,15].

Other survivors have serious psychiatric symptoms, such as depression, anxiety, substance abuse, aggression, shame or cognitive impairments. Finally, some children meet the full criteria for psychiatric illnesses that include post-traumatic stress disorder, major depression, anxiety disorders and sleep disorders [16]. Studies in various countries have highlighted the need for the continuing education of health care professionals on the detection and reporting of early signs and symptoms of child abuse and neglect [17]. In the United States the American Medical Association and the American Academy of Pediatrics have produced diagnostic and treatment guidelines for child maltreatment and sexual abuse [18].

Domestic violence among male patients

In a study population 12.6% of men attending emergency departments were victims of domestic violence by a current or former female intimate partner. In 1978, Steinmetz described "the battered husband syndrome." She suggested that violence committed by women against their male partners had been largely ignored for several reasons. First, there is a stigma associated with being a man beaten by a woman. Therefore, men are unlikely to admit that they have been assaulted by their partners. Second, injuries inflicted by women on men tend to be less severe and presumably less visible. Finally, there has been little research or media attention on the subject. The most common types of abuse involved unarmed physical assaults and throwing of objects. The use of weapons was less common. Few victims sought help of any type, whether it is legal action, medical attention or counseling [6].

ITALIAN LEGISLATION ABOUT GBV

The first significant legislative innovation in the field of sexual violence, in Italy, took place with the approval of the Law of 15 February 1996, n. 66, which began to consider violence against women as a crime against personal freedom, innovating the previous legislation, which placed it among the crimes against public morality and good morals.

With the Law 4 April 2001, n. 154 new measures are introduced aimed at combating cases of violence within the home with the removal of the violent family member. In the same year the Laws n. 60 and the Law of 29 March 2001, n. 134 on legal aid for women, without financial means, raped and/or ill-treated, a fundamental instrument to defend them and assert their rights, in collaboration with the anti-violence centers and the courts.

With the Law of 23 April 2009, n. 38 penalties for sexual violence have been exacerbated and the crime of persecutory acts or stalking is introduced.

Italy has taken a historic step in the fight against gender violence with the law of 27 June 2013 n. 77, approving the ratification of the Istanbul Convention, drawn up on May 11, 2011. The guidelines outlined by the Convention are in fact the track and the lighthouse to implement effective measures, at national level, and to prevent and combat this phenomenon. On October 15, 2013 Law 119/2013 was approved (in force since October 16, 2013) "Conversion into law, with amendments, of the decree-law 14 August 2013, n. 93, which contains urgent provisions on security and to combat gender-based violence". On 24 November 2017, "the National Guidelines for Aziende Sanitarie and Aziende Ospedaliere on the issue of relief and social-health assistance to women victims of violence" were approved.

CONCLUSION

The aim of the guidelines is to provide an adequate and integrated intervention in the treatment of the physical and psychological consequences that male violence produces on women's health. The provision provides, after the nursing triage, unless it is necessary to assign an emergency code (red or equivalent), that the woman is recognized a codification of relative urgency (yellow code or equivalent) to ensure a timely medical examination (time of maximum waiting time 20 min) and minimize the risk of voluntary repentance or departure.

The guidelines also provide for the continuous updating of operators and operators, essential for a good reception, taking charge, risk detection and prevention [19].

REFERENCES

1. Policy and Programme Guidance: HIV and gender-based violence. UN Women Regional Office for Asia and the Pacific. WHO.
2. Khan A (2011) Gender-based violence and HIV: A program guide for integrating gender-based violence prevention and response in PEPFAR programs. Arlington, VA: USAID's AIDS Support and Technical Assistance Resources, AIDSTAR-One, Task Order 1.
3. Ellsberg M, Heise L (2005) Researching violence against women: A practical guide for researchers and activists. Washington DC, United States: World Health Organization, PATH.
4. Breiding MJ (2014) Prevalence and characteristics of sexual violence, stalking and intimate partner violence victimization — National Intimate Partner and Sexual Violence Survey, United States, 2011. *MMWR Surveill Summ* 63: 1-18.
5. Boyle A, Robinson S, Atkinson P (2004) Domestic violence in emergency medicine patients. *Emerg Med J* 21: 9-13.
6. Mechem CC, Shofer FS, Reinhard SS (1999) History of domestic violence among male patients presenting to an urban emergency department. *Acad Emerg Med* 6.
7. Krug E, Dahlberg LL, Mercy JA (2002) World report on violence and health. Geneva, World Health Organization.
8. McCloskey LA, Lichter E, Ganz ML (2005) Intimate partner violence and patient screening across medical specialties. *Acad Emerg Med* 12: 712.
9. Crandall ML, Nathens AB, Kernic MA (2004) Predicting future injury among women in abusive relationships. *J Trauma* 56: 906.
10. Khalifeh H, Oram S, Trevillion K (2015) Recent intimate partner violence among people with chronic mental illness: Findings from a national cross-sectional survey. *Br J Psychiatry* 207: 207.
11. Bonomi AE, Anderson ML, Reid RJ (2009) Medical and psychosocial diagnoses in women with a history of intimate partner violence. *Arch Intern Med* 169: 1692-1697.
12. Lachs MS, Pillemer KA (1995) Abuse and neglect of elderly persons. *N Engl J Med* 332: 437-443.
13. WHO (1999) Report of the Consultation on Child Abuse Prevention. Geneva, World Health Organization.
14. Anda R (1999) Adverse childhood experiences and smoking during adolescence and adulthood. *J Am Med Assoc* 282: 1652-1658.
15. Felitti V (1998) Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. *Am J Prev Med* 14: 245-258.
16. Fergusson DM, Horwood MT, Lynskey LJ (1996) Childhood sexual abuse and psychiatric disorder in

young adulthood. II: Psychiatric outcomes of childhood sexual abuse. *Journal of the American Academy of Child and Adolescent Psychiatry* 35: 1365-1374.

17. Alpert EJ (1998) Family violence curricula in US medical schools. *Am J Prev Med* 14: 273-278.
18. American Academy of Pediatrics (1999) Guidelines for the evaluation of sexual abuse of children: Subject review. *Pediatrics* 103: 186-191.
19. <http://www.salute.gov.it/portale/donna>