

## Cadaverine Level and pH: Indicator of Spoilage in the Flesh of Fish (Carp, Mackerel and Jaw) Storage at Different Temperatures

Traore Moumouny<sup>1\*</sup>, Koffi Yao Fulgence<sup>2</sup>, Tine Alphonse<sup>3</sup>, Kital Khemesse<sup>4</sup>, Oladele Oyelakim<sup>5</sup> and Koussemon Marina Camara<sup>6</sup>

<sup>1,2</sup>*Peleforo Gon COULIBALY University, Department of Biochemistry-Genetics, BP 1328 Korhogo, Cote d'Ivoire*

<sup>3,4</sup>*Cheikh Anta DIOP University, Department of Science and Technology, Laboratory of Photochimie and Analysis, BP 5005 Dakar, Senegal*

<sup>5</sup>*University of Gambia, School of Arts and Sciences, Division of Physical and Natural Sciences, Serrekunda, Gambia*

<sup>6</sup>*Nangui Abrogoua University, Department of Food Science and Technology, Laboratory of Food Biotechnology and Microbiology, 02 BP 801 Abidjan 02, Cote d'Ivoire.*

Published February 28, 2024

### ABSTRACT

Cadaverine (CAD) is an indicator of spoilage in fish. Its level was determined by spectrofluorimetric method in the flesh of selected species of fish stored at -10, 4, 22 and 32°C. Average recovery rates obtained ranged from 87.9 to 101.4% for all species. It was found that CAD was present in all fish species used and its level was essentially constant during storage at -10°C for 32 days. At 4°C, the variation in CAD level was characterized by a stable phase during the first two weeks followed by a descending phase. Over a period of three of storage, CAD levels tripled at temperatures of 22°C and 32°C. Variations in CAD levels were also accompanied by a significant variation in the pH value. Slight pH variations were also observed when fish was stored at -10°C for 32 days and for 18 days at 4°C. Beyond these times, pH values increased significantly. The pH of the fish flesh increased sharply during storage at 22°C and 32°C. The Pearson treatment correlation study indicated a strong positive correlation (0.81 and 0.93) between CAD and pH for all samples stored at 22°C and 32°C, respectively.

**Keywords:** Cadaverine, Spectrofluorimetric method, pH variations

**Corresponding author:** Traore Moumouny, Peleforo Gon COULIBALY University, Department of Biochemistry-Genetics, BP 1328 Korhogo, Cote d'Ivoire, E-mail: traoremoumouny@yahoo.fr

**Citation:** Moumouny T, Fulgence KY, Alphonse T, Khemesse K, Oyelakim O, et al. (2024) Cadaverine Level and pH: Indicator of Spoilage in the Flesh of Fish (Carp, Mackerel and Jaw) Storage at Different Temperatures. *Int J Biopro Biotechnol Advance*, 10(S1): 06.

**Copyright:** ©2024 Moumouny T, Fulgence KY, Alphonse T, Khemesse K, Oyelakim O, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.