

Intraoperative Bradycardic Reactions During Facial and Oral Surgery in the Office: The Role of the Trigemino-Cardiac Reflex (TCR)

Claudio Melloni*

*Consultant in Anesthesia, Bologna, Italy.

Received December 11, 2018; Accepted December 28, 2018; Published December 30, 2018

ABSTRACT

We present a series of 4 patients undergoing blepharoplasty or multiple dental implants who exhibited bradycardia and hypotension during the procedure attributable to the trigemino-cardiac reflex (TCR). The immediate administration of atropine, vasopressors, crystalloids solved the problem.

We advocate the need for a continuous monitoring of all vital signs plus the dedicated presence of the anesthesiologist even during minor surgery since the reaction occur at least in 2% of cases and may evolve into major cardiac events.

Abbreviations: Im: Involuntary movements; LA: Local Anesthesia/Anesthetic; LOC: Loss of Consciousness; PONV: Postoperative Nausea and Vomiting

INTRODUCTION

Vasovagal reactions, resulting in bradycardia and/or hypotension in response to a number of stimuli (fear, pain, unusual sights and smells, anxiety, fatigue, fasting, venipuncture, local anesthesia, viscous distention, etc.) are frequently encountered in surgery and anesthesia.

Surgery can predispose patients to vasovagal episodes owing to the temporo-spatial coincidence of many stimuli; the reactions are usually self-limiting, but potentially life-threatening exceptions have been described up to asystole in different environments, starting with venipuncture [1], dentistry [2], Maxillo facial surgery [3], ophthalmic surgery [4], gastrointestinal endoscopy [5].

A complete review of this topic has been covered in the article [6]; but this excellent review failed to mention the possibility that perioperative bradycardia could originate from the trigeminal nerve. As a matter of fact the surgical procedures more frequently involved include, ophthalmology, maxillofacial, as mentioned above, neuro cranic [7-9] and dental, suggesting that the trigeminal nerve is responsible of the so called trigeminocardiac reflex (TCR) [10].

The TCR is the reflexive response of bradycardia, hypotension, and gastric hyper motility induced with mechanical stimulation in the distribution of the trigeminal nerve; the sensory nerve endings of the Vth cranial nerve arrive at the trigeminal nucleus and the efferent pathway passes through the reticular formation and connects to the

motor nucleus of the vagus nerve; the efferent fibers arrive at the myocardium imposing the typical negative chronotropic and inotropic responses on the heart.

This reflex is triggered usually by a strong stimulus applied in the area innervated by the trigeminal fibers, what is less known and common is the occurrence of vagal reactions intraoperatively, apparently without cause and not accompanied by pain or discomfort.

We present 4 cases of bradycardia and hypotension during conscious sedation done in the office for cosmetic and dental surgery; 3 reactions occurred in the middle of the case without any apparent discomfort: all were attributed to the TCR.

CASE REPORTS

All patients were premedicated 20-30 min before surgery; they were laying supine, ECG, SaO₂, side stream etCO₂ monitored continuously and NIBP every 5 min: vital signs were collected every 5 min. A peripheral vein was in cannulated and normal saline infused at a speed of 100-150 ml/h. Midazolam and fentanyl were administered 5-10 min

Corresponding author: Claudio Melloni, MD, PhD, Consultant in Anesthesia, Bologna, Italy, E-mail: mellonclaudio@libero.it

Citation: Melloni C. (2018) Intraoperative Bradycardic Reactions During Facial and Oral Surgery in the Office: The Role of the Trigemino-Cardiac Reflex (TCR). Int J Anaesth Res, 1(1): 18-21.

Copyright: ©2018 Melloni C. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

before LA; sedation depth assessed according to the Ramsay scale maintaining a score between 2 and 3 at all times.

In all cases local anesthesia was successful and patients did not complain of pain, did not lose consciousness (except perhaps FS), pallor and sweating were not noticed (**Table 1**).

Table 1. Patients and their surgery details.

Name	Age	Weight (kg)	Height (cm)	Asa	Surgery	Premedication (mg)	Sedation	Time of occurrence (min) from start of surgery and till last
SV	51	54	160	2 (micro citemia)	Blefaroplasty sup and inf	Diaz 0.6	5+halop 50	Midaz 2+Fent 30 min
AC	39	60	162	1	Blefaroplasty sup and inf	diaz 5+halop 1	50+Clon 60	Midaz 2+Fent 65 min
FS	68	75	149	2, hypertension, obesity	Multiple dental implants	diaz 0.6	5+ halop 30	midaz 3+mep 60 min
CP	55	83	180	1	Multiple dental implants	diaz 8	midaz 4+fent 80+halop 0.5	85 min

Diaz: Diazepam (mg); fent: fentanyl (μ g); clon: clonidine (μ g); mep=meperidine (mg); halop=haloperidol (mg)

FS was receiving an additional dental block when the reaction started; in the remaining patients surgery and sedation was proceeding smoothly.

SV: BP drop from 102/55 to 69/35, HR from 55 to 35. Therapy Atropine 0.5 mg, ephedrine 5 mg. SaO₂ 100: after 3 min BP 85/52, HR 82. NO pain, no loc, no IM. SaO₂ and et CO₂ stable, AC: BP drop from 103//69 to 74/45, HR from 65 to 45: atropine 0.7 mg, no loc, no pain, no IM, SaO₂ and etCO₂ stable.

FS, BP drop from 103/57 to 89/47, HR from 73 to 43, SaO₂ 93; LOC for 1 min; tonicoclonic movements of the right hand and head: atropine 0.7 mg+etilephrine 1, oxygen 2 L/min+dexamethasone 4 mg; BP to 108/51 and hr 89, within 2 min; SaO₂ 98: crystalloids 250 ml in 5 min.

Patient returned for completion of surgery 3 months later, premedicated with diazepam 6 mg+atropine 0.5 per os, +atropine 0.3 at the beginning of surgery, midazolam 3 mg+fentanyl 50 μ g, procedure uneventful.

CP: bradycardia+hot flush, fast response to atropine 0.8 mg and 250 ml crystalloids.

DISCUSSION

The occurrence of the vaso-vagal reactions during the operation, excluding cases where this type of reaction happens at the moment of venipuncture or during infiltration local anesthesia, could be estimated at around 0.8-1% since these cases were collected in the last 2 years of a private practice limited to dental and cosmetic surgery encompassing roughly 500 cases. This incidence is similar to that reported in larger series like Shalom et al. [11] who reported an incidence around 2% of syncopal episodes over 2600 minor dermatological procedures and Edmondson et al. [12] who described vasovagal episodes in dental surgery.

The etiology of the above reactions surely attributable to the TCR remain unknown, since in all cases the pharmacological sedation provided by the anesthesiologist and the local/truncal analgesia provided by the surgeon were deemed good and in no instances the patients complained of pain, with the exception of patient FS where the reaction occurred during a LA top up in the palate.

The link between the stimulation of any sensory part of the Vth cranial nerve with the hemodynamic irregularities has

been frequently reported in the field of maxillo-facial and neurosurgery [13,14].

The physiology of this reflex has been intensively studied in the most recent years with the aim to allow a more precise definition and classification [15,16]. More recently a chronic form of the reflex has been described [17] so that we may divide the TCR into acute and chronic variants and into peripheral or central according to the site of origin: oculocardiac and maxilla-mandibular being peripheral and any TCR episode due to the stimulation of the fifth nerve pathway beyond the Gasserian ganglion considered as the central TCR.

The primary pathophysiological hemodynamic mechanism involved in our patients remain unknown since the intermittent measurement of blood pressure as opposed to the continuous monitoring of ECG and pulse oximetry could not give any clue as far as the etiology, since the slowing of the ECG gave the first sign of the reaction and prompted the immediate injection of atropine, blood pressure being checked immediately after and supported with vasopressors and crystalloids if found low. Only in the patient FS the reaction could be attributed to sudden stimulation in the area supplied by the greater palatine nerve (V).

Haloperidol, a potent neuroleptic used in the treatment of severe agitation [18] and then in the treatment of schizophrenia and other psychotic disorders [19] was administered at a low dosage to reduce the possibility of PONV [20] in this outpatient high risk population; moreover his protective effect was enhanced by the administration of dexamethasone at the end the procedure according to a multimodal strategy [21,22].

It is possible that the administration of haloperidol, could have been contributory to the onset of hypotension since the drug possesses a moderate alpha adrenergic blocking effect, thereby reducing peripheral vascular resistance and hence inducing a moderate hypotension [23,24]; unfortunately the more specific antiemetic droperidol is not available in Italy outside the hospital.

Atropine remains the milestone for the treatment of acute bradycardia and its advantages have been emphasized during manipulation of percutaneous compression of the trigeminal ganglion for trigeminal neuralgia albeit at low dosage in order to avoid rebound tachycardia [25]; however the clinical situation of our cases is far different from neurosurgery and we did not observe any dangerous rebound increase in HR following its injection, recommending the administration of a full dose, at least 0.5 mg. In the context of conscious sedation we cannot try to abolish the reflex by deepening anesthesia with a bolus of propofol as suggested by Chowdhury et al. [26]

It since might impair the respiration and hemodynamic stability of the patient and cause a shift toward general

anesthesia with the loss of the protective reflexes of the airways.

In order to obtain the best outcome we suggest that beside the drugs to be used for sedation, the diligent anesthesiologist should always have a few prefilled labeled syringes at hand with at least 1 mg of atropine and a vasopressor like ephedrine 30-50 mg, diluted in 10 ml of NS. It is also important that no time should be wasted looking around for the injection port of the intravenous line: a 3 way stopcock situated in the extension set is the best choice for speed of access and prevention of needle stick injuries: at the same time the fully release of the roller clamp maximize the infusion of the crystalloid in use.

CONCLUSION

In conclusion, the sudden appearance of bradycardia and hypotension during the maintenance phase of any procedure around the face emphasizes the need for the dedicated and continuous vigilance from the part of the anesthesiologist; treatment must be immediate in order to prevent the cardiocirculatory collapse.

REFERENCES

1. Wakita R, Ohno Y, Yamazaki S, Kohase H, Umino M, et al. (2006) Vasovagal syncope with asystole associated with intravenous access. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 102: e28-e32.
2. Sharma PR, Hargreaves AD (2006) Malignant vasovagal syncope. *Dent Update* 33: 246-248.
3. Ruhl DS, Ramsey MJ, Ruffin DM (2012) A profound case of neurally mediated syncope with asystole after septoplasty. *J Clin Anesth* 24: 310-314.
4. Hadden PW, Scott RC (2006) Cardiac arrest during phacoemulsification using topical anesthesia in an unsedated patient. *J Cataract Refract Surg* 32: 369.
5. Sharma G, Boopathy Senguttuvan N, Juneja R, Kumar Bahl V (2012) Neurocardiogenic syncope during a routine colonoscopy: An uncommon malignant presentation. *Intern Med* 51: 891-893.
6. Kinsella SM, Tuckey JP (2001) Perioperative bradycardia and asystole: Relationship to vasovagal syncope and the Bezold-Jarisch reflex. *Br J Anaesth* 86: 859-868.
7. Jaiswal AK, Gupta D, Verma N, Behari S (2010) Trigeminocardiac reflex: A cause of sudden asystole during cerebellopontine angle surgery. *J Clin Neurosci* 17: 641-644.
8. Schaller BJ, Fills A, Buchfelder M (2007) Detection and prevention of the trigeminocardiac reflex during skull base surgery. *Acta Neurochir (Wien)* 149: 331.

9. Prabhakar H, Anand N, Chouhan RS, Bithal PK (2006) Sudden asystole during surgery in the cerebellopontine angle. *Acta Neurochir (Wien)* 148: 699-700.
10. Lang S, Lanigan DT, van der Wal M (1991) Trigeminocardiac reflexes: Maxillary and mandibular variants of the oculocardiac reflex. *Can J Anaesth* 38: 757-760.
11. Shalom A, Westreich M, Hadad E, Friedman T (2008) Complications of minor skin surgery performed under local anesthesia. *Dermatol Surg* 34: 1077-1079.
12. Edmondson HD, Gordon PH, Lloyd JM, Meeson JE, Whitehead FI, et al. (1978) Vasovagal episodes in the dental surgery. *J Dent* 6: 189-195.
13. Joshi UM, Munnangi A, Shah K, Patil SG, Thakur N, et al. (2017) Trigemino-cardiac reflex: A phenomenon neglected in maxillofacial surgery? *J Maxillofac Oral Surg* 16: 181-185.
14. Sandu N, Chowdhury T, Meuwly C, Schaller B (2017) Trigeminocardiac reflex in cerebrovascular surgery: A review and an attempt of a predictive analysis. *Expert Rev Cardiovasc Ther* 15: 203-209.
15. Meuwly C, Chowdhury T, Sandu N, Golanov E, Erne P, et al. (2017) Definition and diagnosis of the trigeminocardiac reflex: A grounded theory approach for an update. *Front Neurol* 8: 533.
16. Leon-Ariza DS, Leon-Ariza JS, Nangiana J, Vargas Grau G, Leon-Sarmiento FE, et al. (2018) Evidences in neurological surgery and a cutting edge classification of the trigeminocardiac reflex: A systematic review. *World Neurosurg* 117: 4-10.
17. Chowdhury T, Schaller B (2017) Chronic trigeminocardiac reflex: An underestimated truth. *Front Neurol* 8: 22.
18. Heydari F, Gholamian A, Zamani M, Majidinejad S (2018) Effect of intramuscular ketamine versus haloperidol on short-term control of severe agitated patients in emergency department: A randomized clinical trial. *Bull Emerg Trauma* 6: 292-299.
19. Dold M, Samara MT, Li C, Tardy M, Leucht S (2015) Haloperidol versus first-generation antipsychotics for the treatment of schizophrenia and other psychotic disorders. *Cochrane Database Syst Rev* 16: 1.
20. Brettner F, Janitza S, Prüll K, Weninger E, Mansmann U, et al. (2016) Gender-specific differences in low-dose haloperidol response for prevention of postoperative nausea and vomiting: A register based cohort study. *PLoS One* 11: e0146746.
21. Benevides ML, Oliveira SS, de Aguiar-Nascimento JE (2013) The combination of haloperidol, dexamethasone and ondansetron for prevention of postoperative nausea and vomiting in laparoscopic sleeve gastrectomy: A randomized double-blind trial. *Obes Surg* 23: 1389-1396.
22. Skledar SJ, Williams BA, Vallejo MC, Dalby PL, Waters JH, et al. (2007) Eliminating post-operative nausea and vomiting in outpatient surgery with multimodal strategies including low doses of non-sedating, off-patent anti-emetics: Is "zero tolerance" achievable? *ScientificWorldJournal* 7: 959-977.
23. Schaper WKA, Jagenku AHM, Bogaard JM (1963) Hemodynamic and respiratory responses to dihydrobenzoperidol b(R4749) a potent neuroleptic compound in intact anesthetized dogs. *Arzneimittelforschung* 13: 316.
24. Stanley TH (1978) Cardiovascular effects of droperidol during enflurane and enflurane nitrous oxide anaesthesia in man. *Can Anaesth Soc J* 25: 26.
25. Wang CM, Guan ZY, Cai CH, Zhang J, Wang RW, et al. (2016) Comparative study of atropine combined with sodium nitroprusside pretreatment to prevent trigemino cardiac reflex after trigeminal ganglion compression *J Clin Diagn Res* 10: 9-12.
26. Chowdhury T, Ahuja N, Schaller B (2015) Severe bradycardia during neurosurgical procedure: Depth of anesthesia matters and leads to a new surrogate model of the trigeminocardiac reflex: A case report. *Medicine (Baltimore)* 94: 1-2.