

Giant Follicular Dendritic Cell Sarcoma of Cervical Lymph Node: A Pediatric Case Report

Aboubakr Mabrouki* and Hay El Andalous Rue

*Faculty of Medicine and Pharmacy, Mohammed VI University Hospital Centre/Mohammed I University, Oujda, Morocco.

Received November 29, 2022; Revised December 16, 2022; Accepted December 19, 2022

ABSTRACT

Follicular dendritic cell sarcoma of the neck (FDSC) is a rare neoplasm with only 7 reported cases for pediatric patient.

A 13 year-old girl presented with a left neck tumor. Positron Emission tomography coupled with a cervical scan showed a large left cervical lymphadenopathy hypermetabolic superior jugulocarotid.

Fine-needle aspiration cytology are performed which confirmed a diagnosis of FDSC.

The definitive diagnosis was reached by the Immunohistochemical study that showed tumor cells expressing CD23+, CXCL13+, and CD68+. Some of them are experimenting with CD21+, others cells, rarer, experimenting with S100.

The therapeutic decision was a left cervicotomy associated excision of the mass with a homolateral cervical dissection.

INTRODUCTION

Follicular dendritic cell sarcoma (FDSC) is a rare neoplasm with only 7 reported cases for pediatric patient [1-3].

Follicular dendritic cells (FDSC) are nonlymphoid cells essential for antigen presentation presenting in B lymphofollicles. They are spindle-shaped multinucleate cells with dendritic projections. Some cells can be binucleated or multinucleated. Benign or malignant tumors originating from FDCs rare and only a limited number of cases have been reported [4,5].

The diagnosis of FDSC is made by its typical location in a lymph node, morphology, immunohistochemical findings, and in some cases, ultrastructural findings. We report a case of FDSC. The clinicopathological features and the diagnosis and treatment of this rare tumor is discussed.

AIM

Illustrate a clinical case for pediatric patient of Follicular dendritic cell sarcoma of the cervical lymph node by emphasizing the contribution of the fine need aspiration in the diagnosis and to discuss the different therapeutic means of this affection.

CASE REPORT

A 13-year-old girl, without significant pathological history, presented at the consultation with a left cervical mass without dysphagia, dyspnea or dysphonis.

The mass gradually increases in size over 4 months. Physical examination found a hard mass of 4cm, and the skin opposite was normal.

The blood count with HIV, toxoplasmosis and Epstein Barr virus serology was negative.

A cervical scanner showed a homogeneous upper left oval jugulocarotid lymphadenopathy without central necrosis measured 35mm in diameter (**Figure 1**).

Corresponding author: Aboubakr Mabrouki, Faculty of Medicine and Pharmacy, Mohammed VI University Hospital Centre/Mohammed I University, Oujda, Morocco, Tel: 212653186453; E-mail: mabroukimabrouki03@gmail.com

Citation: Mabrouki A & El Andalous RH. (2022) Giant Follicular Dendritic Cell Sarcoma of Cervical Lymph Node: A Pediatric Case Report. *J Infect Dis Res*, 5(1): 273-276.

Copyright: ©2022 Mabrouki A & El Andalous RH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. Sagittal cervical tomography showing a lymphadenopathy in the upper left jugular region.

Positron emission tomography coupled with a Cervical scan showed a large left cervical lymphadenopathy hypermetabolic superior jugulocarotid SUVmax = 7.2 measured 35mm without other secondary localizations (**Figure 2**).




Figure 2. Large left cervical lymphadenopathy hypermetabolic superior jugulocarotid SUVmax = 7.2.

Fine needle aspiration cytology was performed which showed destruction of the normal ganglion near architecture by spindle cells in a stroma rich in lymphocytes and including some B immunoblasts (**Figure 3**).




Figure 3. Photomicrograph showing a diffuse proliferation replacing normal lymph node architecture.

These spindle cells sometimes have a bilobed nucleus of the dendritic follicle cell type with a large central nucleolus (**Figure 4**).




Figure 4. Spindle cells with a bilobed nucleus of the dendritic follicle cell type with a large nucleolus.

These cells are organized in a basic manner and the phenotype CD3 +, CD21 +, CXCL13 +, D2-40 +, CD23+ (**Figure 5**).




Figure 5. Neoplastic cells express CD 23.




Figure 6. Neoplastic cells express CD20.

Immunostaining with anti-CD20 finds some residual follicles.

EBV search is negative.

The therapeutic decision was a left cervicotomy associated excision of the mass with a homolateral cervical dissection.

The histological of the excised mass and the lymph nodes showed spindle-shaped or ovoid tumor cells, their cytoplasm moderately eosinophilic, their nucleus is rounded rather monotonous with fine chromatin and a visible central nucleolus and absence of ganglionic tumor proliferation.

An immunohistochemical study showed tumor cells expressing CD23+, CXCL13+, and CD68+. Some of them are experimenting with CD21+, other cells, rarer, experimenting with S100.

CD3 and CD20 are positive on entangled cells and negative on tumor cells (**Figure 6**) the diagnosis of FDCS was confirmed.

DISCUSSION

FDCS is a rare neoplasm for patient pediatric with only 7 cases found in the literature. The first case was describe by Apostolos [1].

FDC sarcoma is a very rare neoplasm which was first characterized by Monda [6].

A review of the literature suggests that FDCS has a slight male predominance, and the median age of diagnosis is 40 (age 9-86) [1-5].

The authors discuss the use of fine-needle aspiration for the pre-surgical diagnosis of FDC sarcomas and eliminate other differential diagnoses such as paraganglioma or undifferentiated carcinoma that can be help on the management of this neoplasm [7,8].

Several radiological examinations are requested in case of a cervical mass such a resonance magnetic imaging or positron emission computed tomography but remain non-specific in case of follicular dendritic cell sarcoma.

Histologically, the tumor cells are spindly, ovoid, or polygonal in shape, have eosinophilic cytoplasm, and have indistinct borders, resulting in a syncytial appearance [5,9].

They contain oval to round nuclei with smooth borders and mild atypia.

Immunohistochemically the most sensitive and specific markers for FDCS are CD21, CD23, and CD35 [5,10].

Variably markers found in FDCS include S-100, and vimentin, whereas nonspecific include desmoplakin [4,11].

Phenotype CD68 are nonspecific markers for FDCS [12].

The chemokine CXCL13 and podoplanin (D2-40) produced by the neoplastic FDCS cells can be used as a biomarker to diagnose this tumor [13,14].

An immunohistochemical study on our fine need aspiration showed the positivity of phenotype CD21, CD23, CXCL13 and D2-40.

However, following markers CD23, CD68 and on rare cells, s100 were positive on the removed mass.

The management of FDCS is similar to that of other soft tissue sarcomas; surgical resection remains the standard treatment.

The role of adjuvant treatment in the management of FDSC remains uncertain.

Some authors have suggested adjuvant radiotherapy, and others have recommended chemotherapy or radiotherapy only when the tumor is aggressive, of high volume, and surgically unresectable [9,15,16].

A recent series compared patients who underwent surgical resection alone with those who underwent surgical resection with adjuvant radiotherapy. Their results showed that radiation therapy reduced the rate of recurrence, but much larger studies are needed to confirm this [17].

In the case that we have reported, the decision of the multidisciplinary meeting between surgeons, oncologists and radiotherapists was taken not to add radiotherapy, avoiding complications of radiation therapy such as radio necrosis of the mandible and neuropathy, to the treatment with close monitoring because the lymphadenopathy is completely removed with no capsular breakage and absence of positive lymph nodes recess performed.

CONCLUSION

Follicular dendritic cell sarcoma (FDSC) of the lymph node is a rare neoplasm for the patient pediatric that can be diagnosed on the fine needle aspiration.

The management of FDSC is similar to that of other soft tissue sarcomas, surgical resection remains the standard treatment.

REFERENCES

- Karligkiotis A, Contis D, Bella M, Machouchas N, Volpi L, et al. (2013) Pediatric follicular dendritic cell sarcoma of the head and neck: A case report and review of the literature. *Int J Pediatr Otorhinolaryngol* 77(7): 1059-1064.
- Andersen MJ, Kerr Da, Lisovsky M, Vaickus LJ, Linos K (2020) Fine needle aspiration of an intranodal follicular dendritic cell sarcoma: A case report with molecular analysis and review of the literature. *Diagn Cytopathol* 49(2): E65-E70.
- Fonseca R, Yamakawa M, Nakamura S, van Heerde P, Miettinen M, et al. (1998) Atypical follicular dendritic cell sarcoma and interdigitating reticulum cell sarcoma: A review. *Am J Hematol* 59(2): 161-167.
- Chan JK, Fletcher CD, Nayler SJ, Cooper K (1997) Follicular dendritic cell sarcoma. Clinicopathologic analysis of 17 cases suggests a malignant potential higher than currently recognized. *Cancer* 79(2): 294-313.
- Perez-Ordóñez B, Erlandson R, Rosai J (1996) Follicular dendritic cell tumor: Report of 13 additional cases of a distinctive entity. *Am J Surg Pathol* 20(8): 944-955.
- Monda L, Warnke R, Rosai J (1986) A primary lymph node malignancy with features suggestive of dendritic reticulum cell differentiation. A report of 4 cases. *Am J Pathol* 122(3): 562-572.
- Vicandi B, Jimenez-Heffernan JA, Lopez-Ferrer P, Viguer JM (2000) Fine needle aspiration cytology of follicular dendritic cell sarcoma. A case report. *Acta Cytol* 44(6): 1106-1110.
- Dusenbery D, Watson CG (1996) Fine-needle aspiration biopsy findings in a case of follicular dendritic cell tumor. *Am J Clin Pathol* 106(5): 689-692.
- Duan GJ, Wu F, Zhu J, Guo DY, Zhang R, et al. (2010) Extranodal follicular dendritic cell sarcoma of the pharyngeal region: A potential diagnostic pitfall, with literature review. *Am J Clin Pathol* 133(1): 49-58.
- Jabbour MN, Fedda FA, Tawil AN, Shabb NS, Boulos FI (2014) Follicular dendritic cell sarcoma of the head and neck expressing thyroid transcription factor-1: A case report with clinicopathologic and immunohistochemical literature review. *Appl Immunohistochem Mol Morphol* 22(9): 705-712.
- Desai S, Deshpande RB, Jambhekar N (1999) Follicular dendritic cell tumor of the parapharyngeal region. *Head Neck* 21: 164-167.
- O Soriano A, Thompson MA, Admirand JH, Fayad LE, Rodriguez AM, et al. (2007) Follicular dendritic cell sarcoma: A report of 14 cases and a review of the literature. *Am J Hematol* 82: 725-728.
- Vermi W, Lonardi S, Bosisio D, Uguccioni M, Danelon G, et al. (2008) Identification of CXCL13 as a new marker for follicular dendritic cell sarcoma. *J Pathol* 216(3): 356-364.
- Yu H, Gibson Ja, Pinkus GS, Hornick JL (2007) Podoplanin (D2-40) is a novel marker for follicular dendritic cell tumors. *Am J Clin Pathol* 128(5): 776-782.
- El-Naggar AK, Chan JKC, Grandis JR (2017) WHO Classification of Head and Neck Tumors: In WHO/IaRC Classification of Tumors pp: 9.
- Chan KCJ, Tsang WYW, Ng CS (1994) Follicular dendritic cell tumor and vascular neoplasm complicating hyaline-vascular Castleman's disease. *Am J Surg Pathol* 18(5): 517-525.
- Saygin C, Uzunaslan D, Ozguroglu M, Senocak M, Tuzuner N (2013) Dendritic cell sarcoma: A pooled analysis including 462 cases with the presentation of our case series. *Crit Rev Oncol Hematol* 88: 253-271.