

Alpha-Fetoprotein Producing Tumor of the Colon: A Case Report and Literature Review

Jui-Ten Wu¹ and Swei H Tsung^{2*}

¹Director, Department of Surgery, Camillian Saint Mary's Hospital, Luodong, Yilan, Taiwan

²Director, Department of Pathology, Camillian Saint Mary's Hospital Luodong, Yilan, Taiwan.

Received December 12, 2022; Revised December 17, 2022; Accepted December 26, 2022

ABSTRACT

We recently experienced a 57-year-old woman with pure yolk sac tumor in the colon. Colonoscopy biopsy showed poorly differentiated adenocarcinoma. A right hemicolectomy was performed. The tumor showed a polymorphous pattern, in which the tumor cells grew in solid sheets with papillary and glandular structures; In addition, reticular-microcystic pattern and Schiller-Duval body were also observed. Using different immunochemical stains for alpha-fetoprotein were positive; including Glypican-3, HepPar-1, AFP; beta HCG was negative, ruling out the possibility of containing choriocarcinoma. These morphologic and immunochemical characteristics confirmed the diagnosis of yolk sack tumor. Herein, we described this case, which might represent the first case of yolk sac tumor in the colon in the literature.

Keywords: Alpha-fetoprotein, Colon, Immunochemical stains, Yolk sac tumor

INTRODUCTION

No tumor marker is specific for a particular disease. Initially, elevated serum alpha-fetoprotein (AFP) signals the presence of hepatocellular carcinoma (HCC) or yolk sac tumor (YST) in patients. Later, elevation of serum AFP was also found in cancer from the stomach [1,2], esophagus [3], lung [4] and omentum [5]. There had been a plethora report on alpha-fetoprotein producing gastric carcinoma (AFP-GC) in the literature, predominantly from Japan and China [6,7], a few from Taiwan and other part of the world. Second to AFP-GC was alpha-producing colorectal cancer (AFP-CRC). Nevertheless, the reported AFP-CRC was traditional adenocarcinoma combined with choriocarcinoma or yolk sac tumor. Recently, we encountered a patient who was thought to have AFP-CRC, histological examination after surgery turned out to be pure yolk sac tumor, which is extremely rare. Herein, we report this case and review the literature.

CASE REPORT

A 57-year-old woman visited our outpatient clinic complaining of poor appetite abdominal discomfort nausea and bloody stool for the past two weeks. Laboratory examination revealed anemia, with Hemoglobin level of 11.1 g/dl, HCT of 32.7%. The white blood cells and the platelet count were within normal limits. The blood chemistry profile showed nothing of note. However, her tumor markers were abnormal; Alpha-fetoprotein (AFP) was 1319 ng/ml (normal <7 ng/ml). Serum levels of

carcinoembryonic antigen (CEA) and human gonadotropin (HCG) were within normal limits. She underwent a colonoscopy examination; a large tumor was found at the ascending colon near the hepatic flexure. Biopsy was done, and the diagnosis of poorly differentiated adenocarcinoma was rendered. Preoperative computer tomography was performed for staging purpose (**Figure 1**). There was a tumor near the hepatic flexure, with extramural extension. There was no distant metastasis. Subsequently, right hemicolectomy and lymph node dissection was then performed. The tumor was adhered to the duodenum and pancreas. Dissection was performed with somewhat difficult. There was no metastasis in the liver. Her Serum level of AFP was 370 ng/ml 9days after surgery. Port a tube was installed, unfortunately, she went to a medical center for further therapy.

Corresponding author: Swei H Tsung, M.D., Department of Pathology, Camillian Saint Mary's Hospital, 160 S Chong Chung Rd, Luodong, Yilan, Taiwan, Tel: 886-03-954-4106, ext. 7372, Fax: 886-03-957-2916; E-mail: tsung.sweihsiung@gmail.com

Citation: Wu J-T & Tsung SH. (2023) Alpha-Fetoprotein Producing Tumor of the Colon: A Case Report and Literature Review. *Int J Clin Case Stud Rep*, 5(1): 214-219.

Copyright: ©2023 Wu J-T & Tsung SH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1. Computer tomography of coronal view (A) as well as axial view (B) displaying thickening of the bowel wall indicating a large tumor partially obstructing the lumen.

Histopathology

The postoperative specimen consisted of a segment of colon attached with an appendix. On opening, there was a fungating tumor with central ulceration; measuring 4x3x2 cm, four cm from the proximal resection margin (**Figure 2**). The tumor was grayish white in color, and apparently has invaded through the muscle layer to the serosa. Viewing from the back side, the serosa appeared hemorrhagic with fibrous adhesive bands, indicating adhesion to the adjacent organs. Sections from the tumor displayed a polymorphous appearance, i.e. the tumor was composed of solid sheets with areas of glandular or papillary structure (**Figure 3**). The solid sheets consist of closely packed cells with eosinophilic or clear cytoplasm. In addition, they exhibited a variety of other patterns characteristic for YST: clear cell, hepatoid and reticular microcystic. A few Schiller-Duval bodies were observed (**Figure 4**). Immunohistochemical staining (IHCs) (**Figure 5**) showed the following results: HepPar 1 3+, diffuse; Glypican-3+, focal; AFP 3+, focal; CDX-3 +, diffuse. HCG was negative (not shown). Histological evaluation, in conjunction with IHC studies, we made this case as yolk sac tumor. Because the tumor invaded through the visceral peritoneum, with six lymph nodes positive out of seventeen examined. There was no radiological evidence of distant metastasis. This tumor was staged as pT4aN2a. Later, we submitted the whole tumor for histological examination, no conventional adenocarcinoma or choriocarcinoma components were found which led us to diagnose this tumor as pure yolk sac tumor of the colon.

DISCUSSION AND LITERATURE REVIEW

AFP-GC accounts for the majority of AFP-producing cancers. AFP-GC is considered as the subtype of gastric cancer, because it has propensity to metastasize to the liver and lymph node, resulting in poor prognosis. AFP-CRC behaves aggressively, similar to that of AFP-GC. The cellular and molecular mechanism responsible for their poor prognosis was not clearly understood. An early report indicated that AFP had a suppressive effect on lymphocyte transformation [8]. Another report indicated that AFP could enhance proliferative activity and increase angiogenesis due to positive expression of vascular endothelial growth factor VEGF [9]. In a study in 2000, Amemiya [10] found that hepatocyte growth factor (HGF) and c-met expression was higher in AFP-GC than that in AFP negative gastric cancer. Through the HGF and c-met pathway, tumor growth was promoted. Patients with AFP-CRC behaved poorly similar to patients with AFP-GC, perhaps through the same mechanism. AFP-CRC is extremely rare, since Nakajima et al, reported the first case of AFP-CRC in 1985 [11], only a few cases have been sporadically published in the literature [12], the largest series was from Ren [13]. They analyzed 20 patients on clinicopathological features and prognosis. None of their cases had yolk sac tumor. According to Otani [14], only four cases of colorectal adenocarcinoma with YST component have been reported in the English literature up to 2022, but they were traditional adenocarcinoma combined with YST, or with YST and choriocarcinoma. To the best of our knowledge, our present case is the first report of pure colonic YST.



Figure 2. The resected specimen displayed a fungating tumor with central ulceration, at the ascending colon near the hepatic flexure.




Figure 3. Histologic sections of the tumor displayed polymorphous patterns (H&E stains, x200). (A) Solid pattern. (B) Glandular pattern. (C) Papillary configuration. (D) Clear cell pattern. (E) Hepatoid arrangement. (F) reticular-microcyst.




Figure 4. Schiller-Duval body. (H and E stain, x400).




Figure 5. Immunochemical stains, x200. (A) HepPar 1: 3+, diffuse. (B)Glypican-3: 3+, focal. C.AFP 3+, focal. D. CDX-2 2+, diffuse.

It is worth mentioning that endodermal sinus and yolk sac tumor are used interchangeably. The pathogenesis of the yolk sac tumor in the colon cancer still remains unknown. One theory proposed a retrodifferentiation mechanism of mature cells into primitive pluripotent cells [15]. A case report by Otani [14] supported this theory. In their patient, the cells from the original adenocarcinoma and yolk sac tumor in the recurrence shared mutation in APC and TP53 genes. The second theory was that of the migration misplacement. During embryogenesis, the primitive gonadal cells migrated from the cranial cavity to the external genitalia. During their migration, some primitive germ cell might misplace at the migration pathway. These cells could transform into germ cell tumor [15]. Currently, there is no standard guideline to treat AFP-CRC patients with germ cell component. In the literature, they were usually treated according to traditional adenocarcinoma, receiving adjuvant chemotherapy after resection. It was promising that some patients could benefit from preoperative chemotherapy [16-20].

CONCLUSION

In conclusion, we reported the first case of pure yolk sac tumor in colon, in which primary tumor resection and lymph dissection were performed. It was unfortunate that our patient was transferred to a medical center for further therapy. We missed the opportunity to gain the knowledge about treating pure yolk sac tumor of the colon.

REFERENCES

- McIntire KR, Waldmann TA, Moertel CG, Go VL (1975) Serum alpha-fetoprotein in patients with neoplasms of the gastrointestinal tract. *Cancer Res* 35: 991-996.
- Ishukura H, Fukasawa Y, Ogasawara K, Natori T, Tsukada Y, et al. (1985) An AFP-producing gastric carcinoma with features of hepatic differentiation. A case report. *Cancer* 56: 840-848.
- Khalid Y, Dasu N, Reja, Suga H, Itidiare M, et al. (2020) An extremely rare presentation of an alpha-fetoprotein-producing esophageal adenocarcinoma. *Case Rep Gastroenterol* 14: 497-503.
- Miyake M, Ito M, Mitsuoka A, Taki T, Wada H, et al. (1987) Alpha-fetoprotein and human chorionic gonadotropin-producing lung cancer. *Cancer* 59: 227-232.
- Harano K, Ando M, Sasajima Y, Yunokawa M, Yonemori K, et al. (2012) Primary yolk sac tumor of the omentum: A case report and literature review. *Case Rep Oncol* 5: 671-675.
- Li XD, Wu CP, Ji M, Wu J, Lu B, et al. (2013) Characteristic analysis of a-fetoprotein-producing gastric cancer in China. *World J Surg Oncol* 11: 246e51.
- Takashi Y, Mai O, Orgino T, Ueda H, Sawaguchi K, et al. (1987) Clinicopathological study of AFP producing gastric cancer. *Jpn Surg Soc* 88: 696-700.
- Yachim S (1983) the immunosuppressive properties of alpha-fetoprotein. A brief review. *NY Acad Sci* 417: 105-107.
- Kamei S, Kono K, Amemiya H, Takahashi A, Sugai H, et al. (2003) Evaluation of VEGF and VEGF-C expression in gastric cancer cells producing a-fetoprotein. *J Gastroenterol* 38: 540-547.
- Amemiya H, Kono K, Mori Y, Takahashi A, Ichihara F, et al. (2000) High frequency of c-Met expression in gastric cancer producing alpha-fetoprotein. *Oncology* 59: 145-151.
- Nakajima T, Okazaki T, Morinaga S, Tsumuraya M, Shimosato Y, et al. (1985) A case of alpha-fetoprotein-producing rectal carcinoma. *Jpn J Clin Oncol* 15: 679-685.
- Karakuch N, Toyota K, Kajikawa R, Wanatabe A, Saito Y, et al. (2021) Alpha-fetoprotein-producing colon carcinoma with rapidly increasing liver metastasis that resulted in death: A case report. *J Anus Rectum Colon* 5: 100-106.
- Ren F, Weng W, Zhang Q, Tan C, Xu M, et al. (2019) Clinicopathological features and prognosis of AFP-producing colorectal cancer: A single center analysis of 20 cases. *Cancer Manag Res* 11: 4557.
- Otani T, Kanemura H, Kimura M, Matsmura N, Sato T, et al. (2022) Yolk Sac tumor in a recurrence of colonic adenocarcinoma with shared mutation in APC and TP53 Genesis: A case report. *Int J Surg Pathol* 30: 646-651.
- Nogales FP, Preda O, Nicolae A (2012) Yolk sac tumor revisited. A review of their many faces and names. *Histopathology* 60: 1023-1033.
- Nakamura Y, Matsuda K, Yokoyama S, Tamura K, Mitani Y, et al. (2018) Alpha-fetoprotein producing rectal cancer successfully respond to preoperative chemotherapy. *Surg Case Rep* 4: 11.
- Yu YY, Ogino T, Okada S (1992) an alpha-fetoprotein-producing carcinoma of rectum. *Acta Pathol Jpn* 42: 684-687.
- Kawahara M, Takada A, Tachibana A, Kodama T, Kobayashi H, et al. (2009) Germ cell tumor of the colon with an adenomatous component. *Int J Clin Oncol* 14: 537-540.
- Ostor AG, McNaughton WM, Fortune DW, Rischin D, Hill coat BL, et al. (1993) Rectal Carcinoma with germ cell elements treated with chemotherapy. *Pathology* 25: 243-246.

20. Petricek CM (2001) Colonic adenocarcinoma: Metastasizing as a germ cell neoplasm. A Case report and review of the literature. *Arch Pathol Lab Med* 125: 558-561.