

<i>Combretum paniculatum</i>	-	-	0.5	-	0.13±0.19 ^b
<i>Combretum tomentosum</i>	-	-	0.81	0.03	0.21±0.30 ^b
<i>Dalbergia melanoxylon</i>	0.07	0.71	0.81	1.64	0.81±0.42 ^b
<i>Diospyros mespiliformis</i>	43.63	9.71	2.62	15.68	17.91±12.86 ^u
<i>Entada Africana</i>	-	-	0.17	-	0.04±0.06 ^b
<i>Eucalyptus camaldulensis</i>	-	0.21	0.21	-	0.11±0.11 ^b
<i>Faidherbia albida</i>	35.91	65.69	18.57	44.34	41.13±13.89 ^f
<i>Feretia apodanthera</i>	2.18	0.97	1.23	4.05	2.11±1.01 ^j
<i>Ficus platiphylla</i>	27.48	6.17	10.82	28.24	18.18±9.68 ^q
<i>Ficus sycomoros</i>	11.5	-	34.89	26.41	18.20±12.45 ^q
<i>Ficus thonningii</i>	0.48	-	6.99	-	1.87±2.56 ⁱ
<i>Gardenia aqualla</i>	6.06	16.21	0.5	4.47	6.81±4.70 ^o
<i>Grewia flavescens</i>	0.23	-	-	-	0.06±0.09 ^b
<i>Haematostaphis barteri</i>	-	-	-	0.31	0.08±0.12 ^b
<i>Ipomoea carnea</i>	0.23	-	-	-	0.06±0.09 ^b
<i>Isobertina doka</i>	-	3.17	-	-	0.79±1.19 ^b
<i>Jatropha curcas</i>	-	0.19	0.2	0.06	0.11±0.08 ^b
<i>Lannea barteri</i>	6.43	7.37	1.6	0.96	4.09±2.81 ^r
<i>Lannea velunita</i>	0.37	-	-	-	0.09±0.14 ^b
<i>Parkia biglobosa</i>	-	-	0.16	-	0.04±0.06 ^b
<i>Piliostigma thonningii</i>	15.46	28.55	0.2	9.29	13.38±8.63 ^s
<i>Sclerocarya birrea</i>	5.23	3.3	6.35	12,71	6.90±2.91 ^o
<i>Steganotenia araliaceae</i>	-	-	0.03	-	0.01±0.01 ^b
<i>Sterculia setigera</i>	26.78	5.13	0.68	12.31	11.22±8.32 ^k
<i>Strychnos spinosa</i>	-	0.04	-	-	0.01±0.02 ^b
<i>Tamarindus indica</i>	47.16	30.74	0.81	67.99	36.68±20.90 ^t
<i>Terminalia glauscesens</i>	1.87	-	-	-	0.47±0.70 ^b
<i>Vernonia thomsoniana</i>	0.06	14.14	0.1	1.22	3.88±5.13 ^b
<i>Vitex doniana</i>	1.27	0.05	1.16	-	0.62±0.59 ^b
<i>Ximania americana</i>	-	-	-	1.95	0.49±0.73 ^b
<i>Ziziphus mauritiana</i>	4.46	26.91	0.36	9.3	10.26±833 ^m
<i>Ziziphus mucronate</i>	-	0.6	-	-	0.15±0.23 ^b
<i>Ziziphus spina-christi</i>	0.35	6.13	0.24	2.48	2.30±2.00 ^j
Mean±sd	8.58±10.93 ^a	8.57±10.75 ^a	3.15±4.22 ^b	11.64±14.91 ^c	7.98±2.41 ^p

sd: standard derivation; Number with the same letter in the same row or column are not statistically different at 5 % level of probability; CU1: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

Production of above-ground biomass

Table 2 shows the above-ground biomass of the different species. This biomass varies from one species to another ($p < 0.0001$) but not really between CU ($0.23 > 0.05$). Some species have a very high production capacity than others. Regarding the first group there are: *Anogeissus leiocarpus*

(3393.74 t/ha), *Acacia polyacantha* (3386.10 t/ha), *Faidherbia albida* (951.70 t/ha), *Acacia seyal* (488.99 t/ha) and *Acacia ataxacantha* (422 t/ha) while the second group includes among others: *Adansonia digitata* (0.62 t/ha), *Adenium obesum* (0.01 t/ha), *Bauhinia rufescens* (0.04 t/ha), *Capparis sepiaria* (0.12 t/ha).

Table 2. Variation in Above ground Biomass Production (t/ha).

Species	CU1	CU2	CU3	CU4	Mean±sd
<i>Acacia ataxacantha</i>	218.28	1175.20	3.25	291.77	422±376.54 ^a
<i>Acacia macrostachya</i>	1.40	2.11	-	378.48	95.50±141.49 ^b
<i>Acacia nilotica</i>	1113	24.59	0.10	595.91	157.93±218.99 ^c
<i>Acacia polyacantha</i>	32.50	0.15	11.17	13500.57	3386.10±5057.24 ^d
<i>Acacia seyal</i>	8.88	1907.87	0.02	39.18	488.99±709.44 ^e
<i>Acacia sieberiana</i>	0.30	-	60.34	-	20.21±26.75 ^f
<i>Adansonia digitata</i>	0.02	1.21	-	0.62	0.62±0.40 ^g
<i>Adenium obesum</i>	0.01	0.01	0.01	0.01	0.01±0.00 ^g
<i>Agave sisalana</i>	0.62	-	-	13.25	6.64±6.31 ^h
<i>Annona senegalensis</i>	8.15	573.81	4.71	0.01	146.67±213.57 ⁱ
<i>Anogeissus leiocarpus</i>	8946.01	838.94	179.27	3610.72	3393.74±2884.63 ^j
<i>Azadirachta indica</i>	60.95	29.72	0.38	78.47	42.38±27.33 ^k
<i>Balanites aegyptiaca</i>	662.69	196.84	110.02	1774.93	686.12±544.40 ^l
<i>Bauhinia rufescens</i>	-	0.02	-	0.15	0.04±0.05 ^g
<i>Borassus aethiopicum</i>	54.63	0.01	74.08	-	42.91±28.60 ^m
<i>Boswellia dalzielii</i>	0.42	-	-	-	0.21±0.21 ^g
<i>Calotropis procera</i>	-	148.59	1.40	0.25	50.08±65.67 ⁿ
<i>Capparis sepiaria</i>	-	0.01	-	0.35	0.12±0.15 ^g
<i>Cassia singueana</i>	8.28	7.00	0.01	55.98	17.82±19.08 ^o
<i>Celtis integrifolia</i>	-	0.37	0.06	1.65	0.69±0.64 ^g
<i>Combretum aculeatum</i>	0.11	11.64	2.90	0.51	3.79±3.93 ^p
<i>Combretum collinum</i>	67.01	3.33	0.17	12.04	20.64±23.19 ^f
<i>Combretum paniculatum</i>	-	-	0.03	-	0.03±0.00 ^g
<i>Combretum tomentosum</i>	-	-	0.08	-	0.04±0.04 ^g
<i>Dalbergia melanoxylon</i>	-	0.06	0.08	0.43	0.14±0.14 ^g
<i>Diospyros mespiliformis</i>	865.86	26.52	1.27	80.60	243.56±311.15 ^q
<i>Faidherbia albida</i>	551.12	2237.41	119.34	898.91	951.70±642.86 ^r
<i>Feretia apodanthera</i>	0.83	0.13	0.22	3.49	1.17±1.16 ^s
<i>Ficus platiphylla</i>	296.26	9.26	34.08	315.61	163.80±142.13 ^u
<i>Ficus sycomorus</i>	39.26	-	515.48	270.18	274.97±160.34 ^t
<i>Ficus thonningii</i>	0.02	-	12.37	-	6.20±6.17 ^h
<i>Gardenia aqualla</i>	8.88	87.07	0.03	4.38	25.09±30.99 ^u
<i>Haematostaphis barteri</i>	-	-	-	0.01	0.01±0.00 ^g
<i>Isoberlina doka</i>	-	1.98	-	-	1.98±0.00 ^s
<i>Lamnea barteri</i>	10.19	13.99	0.40	0.12	6.18±5.91 ^h
<i>Lamnea velunita</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Piliostigma thonningii</i>	78.00	323.71	-	23.93	106.41±108.65 ^v
<i>Sclerocarya birrea</i>	6.31	2.17	9.90	49.52	16.97±16.27 ^w
<i>Sterculia setigera</i>	279.04	6.03	0.06	45.98	82.78±98.13 ^x
<i>Tamarindus indica</i>	1037.14	384.26	0.08	2423.38	961±769.05 ^y
<i>Terminalia glauscesens</i>	0.58	-	-	-	0.58±0.00 ^g
<i>Vernonia thomsoniana</i>	-	63.41	-	0.22	15.91±23.75 ^z
<i>Vitex doniana</i>	0.24	-	0.19	-	0.14±0.10 ^g
<i>Ximenia americana</i>	-	-	-	0.64	0.64±0.00 ^g
<i>Ziziphus mauritiana</i>	4.36	282.19	0.01	23.99	77.64±102.28 ^u
<i>Ziziphus mucronate</i>	-	0.04	-	-	0.04±0.00 ^g
<i>Ziziphus spina-christi</i>	0.01	9.12	-	1.12	2.56±3.28 ^g
Mean±sd	308.36±489.35 ^a	220.23±313.77 ^b	25.94±41.41 ^c	612.43±957.32 ^d	291.74±168.66 ^f

Ba: Above ground biomass; sd: standard derivation; Number with the same letter in the same row or column are not statistically different at 5 % level of probability; CU1: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

Production of underground biomass

Underground biomass of different species is shown in **Table 3**. Among these species, those with a high production capacity of root biomass are: *Anogeissus leiocarpus* (431.45

t/ha), *Acacia polyacantha* (389.65 t/ha), *Faidherbia albida* (143.25 t/ha), *Balanites aegyptiaca* (106.17 t/ha). On the other hand, those with a low capacity for producing root biomass are the most numerous and in reality, constitute the

rest of the forest species. Among them, are: *Adenium obesum*, *Combretum paniculatum*, *Lannea velunita*, with substantially zero biomass value (0.01 t/ha), *Adansonia digitata* (0.22 t/ha), *Agave sisalana* (1.81 t/ha). The underground biomass production of the species thus demonstrated is closely linked to the aerial biomass, as shown by Cairns [14]. To this end, the same species known

for the high production of aboveground biomass are those known for the high production of underground biomass. This suggests that species with an enormous aerial part (trunk, branches, leaves) actually have well-developed root systems. The underground biomass (UGB) varies very significantly from one species to another ($P < 0.0001$) but not between CUs ($P = 0.19 > 0.05$).

Table 3. Variation in Underground Biomass Production (t/ha) between CUs.

Species	CU1	CU2	CU3	CU4	Mean±sd
<i>Acacia ataxacantha</i>	40.45	179.05	0.98	52.28	68.19±55.43 ^a
<i>Acacia macrostachya</i>	0.47	0.67	-	65.79	16.73±24.53 ^b
<i>Acacia nilotica</i>	2.92	5.88	0.04	98.26	26.77±35.74 ^c
<i>Acacia polyacantha</i>	7.52	0.06	2.93	1548.07	389.65±579.21 ^d
<i>Acacia seyal</i>	2.39	274.73	0.01	8.87	71.50±101.61 ^e
<i>Acacia sieberiana</i>	0.12	-	12.99	-	4.37±5.75 ^f
<i>Adansonia digitata</i>	0.01	0.41	-	0.23	0.22±0.14 ^g
<i>Adenium obesum</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Agave sisalana</i>	0.23	-	-	3.40	1.81±1.59 ^h
<i>Annona senegalensis</i>	2.21	95.03	1.36	0.01	24.65±35.19 ⁱ
<i>Anogeissus leiocarpus</i>	1076.15	132.93	33.99	482.73	431.45±347.99 ^j
<i>Azadirachta indica</i>	13.10	6.95	0.15	16.38	9.14±5.60 ^k
<i>Balanites aegyptiaca</i>	107.93	36.92	22.08	257.74	106.17±76.67 ^m
<i>Bauhinia rufescens</i>	-	0.01	-	0.06	0.02±0.02 ^g
<i>Borassus aethiopicum</i>	11.90	0.01	15.57	-	9.16±6.10 ^v
<i>Boswellia dalzielii</i>	0.16	-	-	-	0.08±0.08 ^g
<i>Calotropis procera</i>	-	28.80	0.47	0.10	9.79±12.67 ^k
<i>Capparis sepiaria</i>	-	0.01	-	0.14	0.05±0.06 ^g
<i>Cassia singueana</i>	2.25	1.94	0.01	12.16	4.09±4.03 ^f
<i>Celtis integrifolia</i>	-	0.14	0.03	0.54	0.24±0.20 ^g
<i>Combretum aculeatum</i>	0.05	3.04	0.89	0.19	1.04±1.00 ^h
<i>Combretum collinum</i>	14.25	1.00	0.07	3.13	4.61±4.82 ^f
<i>Combretum paniculatum</i>	-	-	0.01	-	0.01±0.00 ^g
<i>Combretum tomentosum</i>	-	-	0.04	-	0.02±0.02 ^g
<i>Dalbergia melanoxylon</i>	-	0.03	0.04	0.16	0.06±0.05 ^g
<i>Diospyros mespiliformis</i>	136.69	6.28	0.43	16.78	40.04±48.32 ^l
<i>Faidherbia albida</i>	91.70	316.26	23.73	141.29	143.25±86.51 ^m
<i>Feretia apodanthera</i>	0.29	0.06	0.09	1.05	0.37±0.34 ^g
<i>Ficus platiphylla</i>	52.99	2.48	7.84	56.04	29.84±24.68 ⁿ
<i>Ficus sycomorus</i>	8.88	-	86.44	48.84	48.06±26.12 ^o
<i>Ficus thonningii</i>	0.01	-	3.20	-	1.61±1.59 ^h
<i>Gardenia aqualla</i>	2.39	17.96	0.01	1.28	5.41±6.27 ^p
<i>Haematostaphis barteri</i>	-	-	-	0.01	0.01±0.00 ^g
<i>Isoberlina doka</i>	-	0.63	-	-	0.63±0.00 ^g
<i>Lannea barteri</i>	2.70	3.57	0.16	0.05	1.62±1.51 ^h
<i>Lannea velunita</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Piliostigma thonningii</i>	16.30	57.30	-	5.74	19.83±18.73 ^q
<i>Sclerocarya birrea</i>	177	0.69	2.63	10.91	4.00±3.45 ^f
<i>Sterculia setigera</i>	50.26	1.70	0.03	10.22	15.55±17.35 ^r
<i>Tamarindus indica</i>	160.33	66.68	0.04	339.38	141.61±108.25 ^s
<i>Terminalia glauscesens</i>	0.21	-	-	-	0.21±0.00 ^g
<i>Vernonia thomsoniana</i>	-	13.57	-	0.09	3.42±5.08 ^x
<i>Vitex doniana</i>	0.10	-	0.08	-	0.06±0.04 ^g
<i>Ximenia Americana</i>	-	-	-	0.23	0.23±0.00 ^g
<i>Ziziphus mauritiana</i>	1.27	50.76	0.01	5.75	14.45±18.16 ^t
<i>Ziziphus mucronata</i>	-	0.02	-	-	0.02±0.00 ^g
<i>Ziziphus spina-christi</i>	0.01	2.45	-	0.38	0.71±0.87 ^g
Mean±sd	42.05±64.27 ^a	34.42±47.36 ^b	4.92±7.36 ^c	79.71±119.46 ^d	40.27±20.60 ^u

Bs: Underground biomass; sd: standard derivation; Number with the same letter in the same row or column are not statistically different at 5 % level of probability; CU1: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

Total biomass production capacity

Total biomass production varies significantly between CUs ($P < 0.05$) and between species ($P < 0.0001$) (Table 4).

Anogeissus leiocarpus stands out from other species by its total biomass production which is equal to 3825.19 t/ha. This species is followed by *Tamarindus indica* (1102.82 t/ha),

Faidherbia albida (1094.94 t/ha), *Acacia ataxacantha* (490.32 t/ha), *Diospyros mespiliformis* (283.61 t/ha) and *Ficus platiphylla* (193.64 t/ha). However, other species have a substantially nil biomass production, namely *Acacia sieberiana*, *Calotropis procera*, *Adenium obesum* and *Celtis integrifolia*. We can also note *Adansonia digitata* (0.84 t/ha) and *Agave sisalana* (8.75 t/ha) to quote.

Table 4. Variation of Total Biomass Production Between CUs (t/ha).

Species	CU1	CU2	CU3	CU4	Mean±sd
<i>Acacia ataxacantha</i>	258.73	1354.25	4.24	344.05	490.32±431.97 ^a
<i>Acacia macrostachya</i>	1.86	2.78	-	444.27	112.23±166.02 ^b
<i>Acacia nilotica</i>	14.05	30.47	0.14	694.17	184.71±254.73 ^c
<i>Acacia polyacantha</i>	40.02	0.21	14.10	15048.65	3775.74±5636.45 ^d
<i>Acacia seyal</i>	11.27	2182.60	0.04	48.05	560.49±811.06 ^e
<i>Acacia sieberiana</i>	0.42	-	73.32	-	24.58±32.50 ^f
<i>Adansonia digitata</i>	0.04	1.63	-	0.85	0.84±0.53 ^g
<i>Adenium obesum</i>	0.02	-	-	-	0.02±0.00 ^g
<i>Agave sisalana</i>	0.85	-	-	16.65	8.75±7.90 ^h
<i>Annona senegalensis</i>	10.37	668.84	6.07	0.01	171.32±248.76 ⁱ
<i>Anogeissus leiocarpus</i>	10022.16	971.87	213.26	4093.45	3825.19±3232.62 ^j
<i>Azadirachta indica</i>	74.05	36.67	0.52	94.86	51.52±32.93 ^k
<i>Balanites aegyptiaca</i>	770.62	233.76	132.10	2032.67	792,29±620.19 ^p
<i>Bauhinia rufescens</i>	-	0.04	-	0.21	0.06±0.07 ^g
<i>Borassus aethiopum</i>	66.53	0.02	89.65	-	52.07±34.70 ^l
<i>Boswellia dalzielii</i>	0.58	-	-	-	0.29±0.29 ^g
<i>Calotropis procera</i>	-	177.39	1.86	0.35	59.87±78.35 ^q
<i>Capparis sepriaria</i>	-	0.02	-	0.49	0.17±0.21 ^g
<i>Cassia siamea</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Cassia singueana</i>	10.53	8.94	0.01	68.14	21.90±23.12 ^m
<i>Celtis integrifolia</i>	-	0.51	0.09	2.18	0.93±0.84 ^g
<i>Combretum aculeatum</i>	0.16	14.68	3.79	0.70	4.83±4.92 ⁿ
<i>Combretum collinum</i>	81.26	4.33	0.24	15.17	25.25±28.00 ^o
<i>Combretum paniculatum</i>	-	-	0.04	-	0.04±0.00 ^g
<i>Combretum tomentosum</i>	-	-	0,12	-	0.06±0.06 ^g
<i>Dalbergia melanoxylon</i>	-	0.09	0,12	0,59	0.20±0.20 ^g
<i>Diospyros mespiliformis</i>	1002.56	32.80	1,70	97,38	283.61±359.47 ^r
<i>Eucalyptus calaudulensis</i>	-	0.01	0,01	-	0.01±0.00 ^g
<i>Faidherbia albida</i>	642.82	2553.68	143,07	1040,20	109.94±729.37 ^s
<i>Feretia apodanthera</i>	1.12	0.18	0,31	4,53	1.54±1.50 ^u
<i>Ficus platiphylla</i>	349.24	11.74	41,93	371,65	193.64±166.81 ^v
<i>Ficus sycomorus</i>	48.15	-	601,93	319,02	185.93±185.93 ^w
<i>Ficus thonningii</i>	0.04	-	15,57	-	7.80±7.77 ^s
<i>Gardenia aqualla</i>	1127	105.02	0,04	5,66	30.50±37.26 ^y
<i>Grewia flavescens</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Haematostaphis barteri</i>	-	-	-	0,01	0.01±0.00 ^g
<i>Ipomoea carnea</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Isoberlina doka</i>	-	2.61	-	-	2.61±0.00 ^z
<i>Lannea barteri</i>	12.89	17.55	0,56	0,18	7.80±7.43 ^{uv}
<i>Lannea velunita</i>	0.02	-	-	-	0.02±0.00 ^g
<i>Piliostigma thonningii</i>	94.30	381.01	0,01	29,67	126.25±127.38 ^ù

<i>Sclerocarya birrea</i>	8.08	2.86	12,53	60,43	20.97±19.73 ^γ
<i>Sterculia setigera</i>	329.30	7.73	0,08	56,19	98.33±115.49 ^β
<i>Tamarindus indica</i>	1197.47	450.93	0,12	2762,76	1102.82±877.29 ^α
<i>Terminalia glauscescens</i>	0.80	-	-	-	0.80±0.00 ^ε
<i>Vernonia thomsoniana</i>	-	76.99	-	0,30	19.32±28.83 ^λ
<i>Vitex doniana</i>	0.33	-	0,27	-	0.20±0.13 ^ε
<i>Ximenia americana</i>	-	-	-	0,87	0.87±0.00 ^ε
<i>Ziziphus mauritiana</i>	5.64	332.95	0,02	29,74	92.09±120.43 ^μ
<i>Ziziphus mucronate</i>	-	0.06	-	-	0.06±0.00 ^ε
<i>Ziziphus spina-christi</i>	0.02	11.57	0,01	1,50	3.27±4.15 ^ε
Mean±sd	350.41±552.73a	254.65±361.00b	30.86±49.06c	692.14±1075.95d	332.02±189.26 ^ν

Bt: Total biomass; sd: standard derivation; Number with the same letter in the same row or column are not statistically different at 5 % level of probability; CU1: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

Carbon stock estimate

The total amount of carbon stored by all the CUs is estimated on average at 166.01 tC/ha (Table 5). It varies significantly ($P = 0.03 < 0.05$) between CUs and between species. CU4 is the one that has a quantity of carbon (346.07 tC/ha) greater than those of other CUs which do not differ significantly ($P > 0.05$) from each other (15.43-175.20 tC/ha). The difference between UC4 and other UCs is due to

the low logging activity due to the difficult accessibility of UC4 to neighboring populations. In terms of species in general, those that sequester the most carbon is: *Anogeissus leiocarpus* (1912.59 tC/ha), *Acacia polyacantha* (1887.87 tC/ha) *Tamarindus indica* (551.41 tC/ha), *Faidherbia albida* (547.47 tC/ha), *Balanites aegyptiaca* (396.14 tC/ha), *Ficus sycomorus* (161.52 tC/ha), and *Diospyros mespiliformis* (141.80 tC/ha).

Table 5. Variation in the Quantity of Carbon Stored (tC/ha).

Species	CU1	CU2	CU3	CU4	Mean±sd
<i>Acacia ataxacantha</i>	129.37	677.12	2.12	172.03	245.16±215.98 ^a
<i>Acacia macrostachya</i>	0.93	1.39	-	222.14	56.11±83.01 ^b
<i>Acacia nilotica</i>	7.03	15.23	0.07	347.08	92.35±127.37 ^c
<i>Acacia polyacantha</i>	20.01	0.11	7.05	7524.32	1887.87±2818.23 ^d
<i>Acacia seyal</i>	5.64	1091.30	0.02	24.03	280.25±405.53 ^e
<i>Acacia sieberiana</i>	0.21	-	36.66	-	12.29±16.25 ^f
<i>Adansonia digitata</i>	0.02	0.81	-	0.43	0.42±0.27 ^g
<i>Adenium obesum</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Agave sisalana</i>	0.43	-	-	8.33	4.38±3.95 ^e
<i>Annona senegalensis</i>	5.18	334.42	3.04	0.01	85.66±142.38 ^f
<i>Anogeissus leiocarpus</i>	5011.08	485.94	106.63	2046.73	1912.59±1616.31 ^g
<i>Azadirachta indica</i>	37.02	18.34	0.26	47.43	25.76±16.46 ^h
<i>Balanites aegyptiaca</i>	385.31	116.88	66.05	1016.34	396.14±310.10 ⁱ
<i>Bauhinia rufescens</i>	-	0.02	-	0.10	0.03±0.04 ^g
<i>Borassus aethiopum</i>	33.26	0.01	44.83	-	26.03±17.35 ^j
<i>Boswellia dalzielii</i>	0.29	-	-	-	0.15±0.15 ^g
<i>Calotropis procera</i>	-	88.69	0.93	0.17	29.93±39.17 ^k
<i>Capparis sepriaria</i>	-	0.01	-	0.25	0.08±0.11 ^g
<i>Cassia singueana</i>	5.26	4.47	0.01	34.07	10.95±11.56 ^l
<i>Celtis integrifolia</i>	-	0.26	0.04	1.09	0.46±0.42 ^g
<i>Combretum aculeatum</i>	0.08	7.34	1.89	0.35	2.42±2.46 ^m
<i>Combretum collinum</i>	40.63	2.17	0.12	7.59	12.63±14.00 ⁿ
<i>Combretum paniculatum</i>	-	-	0.02	-	0.02±0.00 ^g
<i>Combretum</i>	-	-	0.06	-	0.03±0.03 ^g

<i>tomentosum</i>					
<i>Dalbergia melanoxylo</i>	-	0.05	0.06	0.30	0.10±0.10 ^g
<i>Diospyros mespiliformis</i>	501.28	16.40	0.85	48.69	141.80±179.74 ^o
<i>Faidherbia albida</i>	321.41	1276.84	71.53	520.10	547.47±364.68 ^p
<i>Feretia apodanthera</i>	0.56	0.09	0.16	2.27	0.77±0.75 ^g
<i>Ficus platiphylla</i>	174.62	5.87	20.96	185.82	96.82±83.40 ^q
<i>Ficus sycomorus</i>	24.07	-	300.96	159.51	161.52±92.96 ^r
<i>Ficus thonningii</i>	0.02	-	7.79	-	3.90±3.88 ^s
<i>Gardenia aqualla</i>	5.64	52.51	0.02	2.83	15.25±18.63 ^y
<i>Haematostaphis barteri</i>	-	-	-	0.01	0.01±0.00 ^g
<i>Isobertina doka</i>	-	1.30	-	-	1.30±0.00 ^t
<i>Lannea barteri</i>	6.44	8.78	0.28	0.09	3.90±3.71 ^u
<i>Lannea velunita</i>	0.01	-	-	-	0.01±0.00 ^g
<i>Piliostigma thonningii</i>	47.15	190.51	-	14.83	63.12±63.69 ^v
<i>Sclerocarya birrea</i>	4.04	1.43	6.26	30.21	10.49±9.86 ^w
<i>Sterculia setigera</i>	164.65	3.87	0.04	28.10	49.16±57.74 ^x
<i>Tamarindus indica</i>	598.74	225.47	0.06	1381.38	551.41±438.65 ^y
<i>Terminalia glauscesens</i>	0.40	-	-	-	0.40±0.00 ^g
<i>Vernonia thomsoniana</i>	-	38.49	-	0.15	9.66±14.42 ^w
<i>Vitex doniana</i>	0.17	-	0.14	-	0.10±0.07 ^g
<i>Ximenia Americana</i>	-	-	-	0.44	0.44±0.00 ^g
<i>Ziziphus mauritiana</i>	2.82	166.48	0.01	14.87	46.04±60.22 ^y
<i>Ziziphus mucronate</i>	-	0.03	-	-	0.03±0.00 ^g
<i>Ziziphus spina-christi</i>	0.01	5.78	-	0.75	1.64±2.07 ^t
Mean±sd	175.20±27636 ^a	127,33±180.50 ^b	15,43±24.53 ^c	346.07±537.98 ^d	166.01±94.63 ^u

sd: standard derivation; Numbers with the same letter in the same row or column are not statistically different at 5 % level of probability; UCI: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

Carbon credit assessment in the different CUs

The average carbon credit value in all CUs is 1660.08 dollars (\$), equivalent to 457328.71 FCFA (Table 6).

Table 6. Carbon Credit of Species According to CU (\$).

Species	CU1	CU2	CU3	CU4	Mean±sd
<i>Acacia ataxacantha</i>	1293.66	6771.25	21.18	1720.26	2451.59±2159.83 ^a
<i>Acacia macrostachya</i>	9.31	13.89	0.01	2221.37	561.14±830.11 ^b
<i>Acacia nilotica</i>	70.25	152.35	0.70	3470.84	923.53±1276.65 ^c
<i>Acacia polyacantha</i>	200.08	1.07	70.49	75243.23	18878.72±28182.26 ^d
<i>Acacia seyal</i>	56.35	10913.02	0.18	240.26	2802.45±4055.28 ^e
<i>Acacia sieberiana</i>	2.08	-	366.62	-	122.90±162.48 ^f
<i>Adansonia digitata</i>	0.18	8.13	-	4.27	4.19±2.67 ^g
<i>Adenium obesum</i>	0.08	-	-	-	0.08±0.00 ^h
<i>Agave sisalana</i>	4.27	-	-	83.25	43.76±39.49 ^o
<i>Annona senegalensis</i>	51.83	3344.22	30.37	0.07	856.62±1243.80 ⁱ
<i>Anogeissus leiocarpus</i>	50110.80	4859.36	1066,32	20467.25	19125.93±16163.09 ^j
<i>Azadirachta indica</i>	370.25	183.36	2.61	474.28	257.62±164.62 ^k
<i>Balanites aegyptiaca</i>	3853.10	1168.80	660.49	10163.35	3961.44±3100.96 ⁿ
<i>Bauhinia rufescens</i>	-	0.19	-	1.04	0.31±0.37 ^h

<i>Borassus aethiopicum</i>	332.64	0.11	448.26	-	260.34±173.49 ^l
<i>Boswellia dalzielii</i>	2.92	-	-	-	1.46±1.46 ^m
<i>Calotropis procera</i>	-	886.94	9.31	1.73	299.33±391.74 [@]
<i>Capparis sepiaria</i>	-	0.08	0.01	2.46	0.85±1.07 ^h
<i>Cassia siamea</i>	0.03	-	-	-	0.03±0.00 ^h
<i>Cassia singueana</i>	52.64	44.70	0.07	340.68	109.52±115.58 ⁿ
<i>Celtis integrifolia</i>	-	2.57	0.44	10.92	4.64±4.18 ^o
<i>Combretum aculeatum</i>	0.79	73.39	18.94	3.51	24.16±24.62 ^p
<i>Combretum collinum</i>	406.29	21.67	1.21	75.85	126.25±140.02 ^q
<i>Combretum paniculatum</i>	-	-	0.21	-	0.21±0.00 ^h
<i>Combretum tomentosum</i>	-	-	0.61	-	0.31±0.30 ^h
<i>Dalbergia melanoxylon</i>	-	0.45	0.61	2.96	1.01±0.98 ^m
<i>Diospyros mespiliformis</i>	5012.78	163.98	8.49	486.89	1418.04±1797.37 ^r
<i>Entada Africana</i>	-	-	0.02	-	0.02±0.00 ^h
<i>Eucalyptus camaldulensis</i>	-	0.03	0.03	-	0.03±0.00 ^h
<i>Faidherbia albida</i>	3214.12	12768.39	715.35	5200.98	5474.71±3646.84 ^s
<i>Feretia apodanthera</i>	5.61	0.91	1.55	22.66	7.69±7.49 ^t
<i>Ficus platiphylla</i>	1746.22	58.70	209.63	1858.24	968.20±834.03 ^u
<i>Ficus sycomorus</i>	240.73	-	3009.63	1595.12	1615.16±929.65 ^v
<i>Ficus thonningii</i>	0.19	-	77.85	-	39.02±38.83 ^w
<i>Gardenia aqualla</i>	56.35	525.12	0.21	28.32	152.50±186.31 ^x
<i>Grewia flavescens</i>	0.04	-	-	-	0.04±0.00 ^h
<i>Haematostaphis barteri</i>	-	-	-	0.07	0.07±0.00 ^h
<i>Ipomoea carnea</i>	0.04	-	-	-	0.04±0.00 ^h
<i>Isobertina doka</i>	-	13.04	-	-	0.13±0.00 ^h
<i>Jatropha curcas</i>	-	0.02	0.03	-	0.02±0.01 ^h
<i>Lannea barteri</i>	64.44	87.77	2.80	0.89	38.98±37.13 ^y
<i>Lannea velunita</i>	0.11	-	-	-	0.11±0.00 ^h
<i>Parkia biglobosa</i>	-	-	0.02	-	0.02±0.00 ^h
<i>Piliostigma thonningii</i>	471.50	1905.06	0.03	148.33	631.23±636.92 ^z
<i>Sclerocarya birrea</i>	40.39	14.28	62.64	302.13	104.86±98.63 ^z
<i>Sterculia setigera</i>	1646.50	38.66	0.41	280.96	491.63±577.43 ^y
<i>Tamarindus indica</i>	5987.36	2254.67	0.61	13813.78	5514.11±4386.46 ^z
<i>Terminalia glauscesens</i>	3.98	-	-	-	3.98±0.00 ^u
<i>Vernonia thomsoniana</i>	-	384.93	0.01	1.52	96.62±144.16 ^z
<i>Vitex doniana</i>	1.67	-	1.36	-	1.01±0.67 ^m
<i>Ximenia americana</i>	-	-	-	4.37	4.37±0.00 ^z
<i>Ziziphus mauritiana</i>	28.18	1664.77	0.10	148.70	460.44±602.17 ^z
<i>Ziziphus mucronate</i>	-	0.31	-	-	0.13±0.00 ^h
<i>Ziziphus spina-christi</i>	0.09	57.84	0,04	7.50	16.37±20.73 ^z
Mean±sd	1752.04±2763.63 ^a	1273.26±1804.98 ^b	154.31±245.28 ^c	3460.70±5379.76 ^d	1660.08±946.29 ^e

Number with the same letter in the same row or column are not statistically different at 5% level of probability; CU1: wooded savannah; CU2: shrub savannah; CU3: degraded savannah; CU4: wooded savannah

It varies very significantly from one species to another ($0.000 < 0.0001$) but not significantly between CUs ($P=0.23 > 0.05$). Among these CUs, CU4 has a carbon credit value ($3460.70 \$ \approx 93026.41 \text{CFAF}$) higher than those of other CUs ($154.31-1752.04 \$$). What shows the preponderant value of CU4 (wooded savannah), it is composition as a blossoming tree, the trunks of the trees that compose it are well developed, while in other CU (wooded savannah, shrub savannah and degraded savannah) the trees there are small in diameter. In addition, the exploitation of natural resources is regular in these CUs, because they are easily accessible to residents.

As for the species, those with the most carbon credit is for example *Anogeissus leiocarpus* (19125.93 \$), *Acacia polyacantha* (18878.72 \$), *Acacia ataxacantha* (2451.59 \$), *Ficus platiphylla* (968.20 \$), *Annona senegalensis* (856.62 \$) and finally *Balanites aegyptiaca* (396144 \$), those with the least carbon credit are among others *Adansonia digitata* (4.19 \$), *Cassia siamea* (0.03 \$) and finally *Capparis sepiaria* (0.85 \$). The value of carbon credit evaluated in this work is higher than that of Kodji [17] who worked in the same area. This difference in value would probably be linked to the surface area of the Collection Units (15 ha) compared to the latter's study sites (12 ha), growth of other plants and human action. There is also the large number of large individuals (circumference) inventoried in this study.

DISCUSSION

Biomass production

What shows the difference in biomass production between individuals would be their density and size such as diameter (**Table 1 & Table 4**).

Among the CUs, CU4 is the one with the highest above-ground biomass (612.43 t/ha), followed by CU1 (308.36 t/ha), then CU2 (220.23 t/ha) and finally CU3 (25.94 t/ha).

The big difference between CU4 (wooded savannah) and other CUs in terms of biomass would be linked to the abundance of large woody plants compared to other CUs whose woody plants are much more exploited. The above-ground biomass of woody plants in CUs in the study area varies from 25.94 to 612.43 t/ha. In comparison to the values of dry areas such as that of Mozogo-Gokoro National Park. Sandjong [18] estimated an average variation of 19.02 to 35.61 t/ha or even that of West Africa. Valbuena [19] found values between 0.49 and 18.91 t/ha.

Total biomass production capacity based on Collection Units

The significant amount of biomass observed in some species can be explained by their well-developed vegetative part (trunks, roots, leaves, branches). With regard to CUs, CU4 has a higher total biomass production (692.14 t/ha) than that of other CUs (30.86-350.41 t/ha) which do not differ significantly between them. This large total biomass

production in UC4 would be linked to the conservation of woody phyto diversity due to low exploitation (**Table 4**). The other three CUs (CU1, CU2 and CU3) have undergone heavy exploitation, this because of the uneven terrain well accessible to operators compared to CU4 which is not only at altitude but seems less accessible. Ibrahim and Abib Fanta [20] in their work on estimating the carbon stock in the tree and shrub faces of the Sudano-Guinean savannahs of Ngaoundere, Cameroon and Tchobsala [21] on carbon sequestration of anthropized vegetation in Ngaoundere (Adamaoua-Cameroon) showed that the differences between the faces of the savannahs are due to the differences in anthropogenic activities. As Herintsitohaina [22] confirmed in his work on the potential for carbon storage in the plant-soil system of Eucalyptus plantations in the Malagasy highlands.

Carbon stock estimate

The amount of carbon stored by individuals is influenced by their abundance and size. Mbow [23] who studied the challenges and hopes of REDD+ in Africa and Tchobsala [6] worked on the impact of logging and carbon sequestration in the Guinean savannah of Ngaoundere Region of Adamaoua-Cameroon demonstrated that human activities negatively influence the amount of carbon stored in vegetation, so as Herintsitohaina [22] for its part, but the emphasis rather on the taxonomic side, i.e. the amount of carbon stored varies with the species. In our study, the two factors could explain the difference in carbon stock between CUs and between species.

Carbon credit assessment in the different CUs

The value of carbon credit evaluated in this work is higher than that of Kodji [17] who worked in the same area. This difference in value would probably be linked to the surface area of the Collection Units (15 ha) compared to the latter's study sites (12 ha) and growth of other plants and human action. There is also the large number of large individuals (circumference) inventoried in this study.

CONCLUSION

This study enabled us to enlighten everyone and raise awareness of the impact of vegetation exploitation of the Minawao zone on carbon stock. 7143 individuals divided into 59 species, 42 genera and 30 families were inventoried in the different CUs. The amount of carbon evaluated between these different CUs varies considerably between them. The quantity of carbon stored by all the Collection Units was estimated at an average of 166.01 tC/ha, or the equivalent of 1660.08 \$, corresponding to 457328.71 FCFA. UC4 is the one with a quantity of carbon (346.07 tC/ha) greater than those of other CUs between them (15.43-175.20 tC/ha), therefore it is the one with a carbon credit value ($3460.70 \$ \approx 93026.41 \text{FCFA}$) higher than those of other CUs ($154.31- 1752.04 \$$). It is very important to

protect our environment so as to permit the vegetation sequestrate more carbon.

ACKNOWLEDGMENT

We would like to thank the Lamido of Gawar with his notables who were useful to us for surveys. We cannot forget the willingness of the population to provide us with useful information for our work.

REFERENCES

- UNFCCC (1992) United Nations Framework Convention on Climate Change New York, pp: 9.
- UNFCCC (1997) Kyoto Protocol to the United Nations Framework Convention on Climate Change.
- Cop21 (2015) Conférence de Paris de 2015 sur les changements climatiques. pp: 5.
- Tchotsoua M, Mapongmetsem PM, Tago M (2000) Urbanisation, crise économique et dynamique de l'environnement en milieu soudanien d'altitude: le cas du plateau de Ngaoundéré. *Revue géographique du Cameroun. Société et environnement au Cameroun* 14(2): 225-249.
- Mbolo T (2013) Characterization and impact of wood logging on plant formations in Ngaoundéré District, Adamawa Region, Cameroon. *J Ecol Nat Environ* 5(10): 265-277.
- Tchobsala, Mbolo M, Souare K (2014) Impact of wood logging on the phytomass and carbon sequestration in the guinea savanna of Ngaoundéré, Adamaoua Region, Cameroon. *Glob Adv Res J Environ Sci Toxicol* 3(3): 038-048.
- UNHCR (2018) Profil du camp pp: 4.
- Tchobsala (2011) Impact des coupes de bois sur la végétation naturelle de la zone périurbaine de Ngaoundéré (Adamaoua). [Doctoral dissertation, Université de Yaoundé I, Cameroun.]
- Lanly JP (1981) Manuel d'inventaire forestier, avec références particulières 41 aux forêts tropicales hétérogènes. *Études FAO: Forêts 27*, FAO, Rome, Italie. pp: 9.
- Schreuder HT, Banyard SG, Brink GE (1987) Comparison of three sampling methods in estimating stand parameters for a tropical forest. *Forest Ecol Manag* 21(1-2): 119-127.
- Arbonnier, M. Bonnet, Grard P (2008) *Ligneux du Sahel. V.1.0 CIRAD*.
- Maydell HJV (1990) *Arbres et Arbustes du Sahel. Livre version française de Jean-Bernad Chappious*. pp: 267.
- Brown S, Gilespe AJR, Lugo AE (1997) Biomass estimation methods for tropical forest with application to forest inventory data. *For Sci* 35(4): 881-902.
- Cairns MA, Brown S, Helmer EH, Baumgardner GA (1997) Root biomass carbon estimates for 1980. ORNL/CDIAC-92, DCP-055, Carbon Dioxide Information FAO Forestry 111: 1-11.
- IPCC (2000) Land use, land-use change and forestry (LULUCF). Watson, R.T. et al. (eds). Cambridge University Press pp: 375.
- Ecosystems Marketplace (2019) State of the voluntary carbon market. *Ecosystem Services & Management* pp: 25.
- Kodji P (2017) Impacts de l'installation des réfugiés sur la dynamique du couvert végétal du camp de Minawao dans le Département du Mayo-Tsanaga (Extrême Nord-Cameroun): implication pour une gestion durable. Mémoire de Master II, Université de Ngaoundéré. pp: 47.
- Sandjong SRC (2018) Etude phytoécologique du Parc National de Mozogo-Gokoro dans l'extrême-Nord Cameroun: implication pour une gestion durable. [Doctoral dissertation, Université de Maroua].
- Valbuena R, Heiskanen J, Aynekulu E, Pitkänen S, Packalen P (2016) Sensitivity of Above-Ground Biomass Estimates to Height-Diameter Modelling in Mixed-Species West Africa Woodlands. *PLoS One* 11(7): 1-2.
- Ibrahim A, Fanta AC (2008) Estimation du stock de carbone dans les faciès arborées et arbustives des savanes soudano-guinéennes de Ngaoundéré, Cameroun. *J Experim Biol* 04(01): 1-11.
- Tchobsala, Dongock ND, Nyasiri J, Ibrahima A (2016) Carbon storage of anthropoid's vegetation on the Ngaoundere escarpment (Adamawa-Cameroon). *J Adv Biol* 9(2): 1-10.
- Herintsitohaina RR (2009) Potentialités de stockage des carbone dans le système plante-sol des plantations d'Eucalyptus des haute terres Malgaches. [Doctoral dissertation]. pp: 193.
- Mbow C, Skole D, Dieng M, Justice C, Kwesha D, et al. (2012) Challenges and Prospects for REDD+ in Africa: Desk Review Of REDD+ Implementation in Africa. GLP Report No. 3. GLP-IPO, Copenhagen; pp: 145.