

























65. Jo S, Kim H, Kim S, Shin DH, Kim MS (2019) Characteristics of flavonoids as potent MERS-CoV 3C-like protease inhibitors. *Chem Biol Drug Des* 94(6): 2023-2030.
66. Jo S, Kim S, Shin DH, Kim MS (2020) Inhibition of SARS-CoV 3CL protease by flavonoids. *J Enzyme Inhib Med Chem* 35(1): 145-151.
67. Ryu YB, Park SJ, Kim YM, Lee JY, Seo WD, et al. (2010) SARS-CoV 3CLpro inhibitory effects of quinone-methide triterpenes from *Tripterygium regelii*. *Bioorg Med Chem Lett* 20(6): 1873-1876.
68. Park JY, Kim JH, Kwon JM, Kwon HJ, Jeong HJ, et al. (2013) Dieckol, a SARS-CoV 3CLpro inhibitor, isolated from the edible brown algae *Ecklonia cava*. *Bioorg Med Chem* 21(13):3730.
69. Peng C, Zhu Z, Shi Y, Wang X, Mu K, et al. (2020) Yang Y, Zhang X, Xu Z. Exploring the binding mechanism and accessible angle of SARS-CoV-2 spike and ACE2 by molecular dynamics simulation and free energy calculation.
70. Ho TY, Wu SL, Chen JC, Li CC, Hsiang CY (2007) Emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction. *Antivir Res* 74(2): 92-101.
71. Bosch BJ, Van der Zee R, De Haan CA, Rottier PJ (2003) The coronavirus spike protein is a class I virus fusion protein: Structural and functional characterization of the fusion core complex. *J Virol* 77(16): 8801-8811.
72. Adem S, Eyupoglu V, Sarfraz I, Rasul A, Ali M (2020) Identification of potent COVID-19 main protease (Mpro) inhibitors from natural polyphenols: An in-silico strategy unveils a hope against CORONA.
73. Rolta R, Yadav R, Salaria D, Trivedi S, Imran M, et al. (2021) In silico screening of hundred phytochemicals of ten medicinal plants as potential inhibitors of nucleocapsid phosphoprotein of COVID-19: An approach to prevent virus assembly. *J Biomol Struct Dyn* 39(18): 7017-7034.