

Are the Ocular Biometric Factors Important for Intraocular Pressure Assessment?

Pragati Garg*

*Department of Ophthalmology, Era's Lucknow Medical College and Hospital, Lucknow, UP, India.

Received March 02, 2019; Accepted March 07, 2019; Published April 26, 2019

Glaucomatous optic neuropathy is characterized by raised intraocular pressure, visual field defects and optic disc changes and of these; intraocular pressure is the principle modifiable risk factors for the development and progression of glaucoma [1]. Regulation of intraocular pressure is a complex physiological trait that depends on the production of aqueous humor, resistance to aqueous humor outflow and episcleral venous pressure [2]. Intraocular pressure is the hydrostatic pressure exerted by the aqueous humor.

The complex physiology involved in aqueous humor formation and its maintenance indicates its dependence on other systemic, physical, physiological and environmental factors and as the intraocular pressure is also dependent on aqueous humor, therefore, indirectly the intraocular pressure is also dependent on these factors. Ocular biometric factors reported to be involved with the intraocular pressure are central corneal thickness [3,4] corneal curvature, axial length, anterior chamber depth, Shaffer's angle and grade and refractive error [5-7].

Opinions regarding relationship between central corneal thickness (CCT) and intraocular pressure (IOP) are quite controversial. A reduced corneal thickness of 0.45 mm could produce an underestimation of the intraocular pressure by up to 4.7 mm Hg, whereas increased central corneal thickness of 0.59 mm could cause an overestimation of 5.2 mm Hg when the actual intraocular pressure is 20 mm Hg by Goldman applanation tonometer [8].

In the current issue, Garg et.al. Have shown a weak positive and a significant correlation between central corneal thickness (CCT) and intraocular pressure (IOP) ($r=0.092$, $p<0.001$) as do other studies [9,10]. Mean CCT showed an incremental trend with the increasing intraocular pressure categories and mean intraocular pressure of those having central corneal thickness $>600 \mu\text{m}$ was significantly higher as compared to that of patients with $\text{CCT}<540 \mu\text{m}$.

As there are multiple variables that might affect the intraocular pressure, therefore, the association derived on univariate assessment seems to be vague and may be the effect of some confounder. In order to elaborate the role of independent factors associated with intraocular pressure, a

multivariate analysis is a must, as done by Garg et al. In their research the IOP is projected as a dependent variable with age, gender, central corneal thickness, axial length and anterior chamber depth as independent variables. Out of this younger age, male gender, thicker central corneal thickness, steeper corneal curvature and short anterior chamber depth were seen to be predictors of higher intraocular pressure, as has been shown [10]. But different researchers have viewed and assessed the association of IOP with many variables using a unique predictive model. Hence, there is a need to come up with a set of stronger predictors present in all the studies.

Therefore, it is recommended that as intraocular pressure is a very important and first line assessment for glaucoma diagnosis, it should be calculated keeping in mind the various ocular, anthropometric and systemic factors which have an association with it.

REFERENCES

1. Wu SY, Leske MC (1997) Associations with intraocular pressure in the Barbados Eye Study. *Arch Ophthalmol* 115: 1572-1576.
2. Allingham RR, Damji KF, Freedman S, Moroi SE, Shafranov G, et al. (2005) Cellular and molecular biology of aqueous humor dynamics. *Shields' Textbook of Glaucoma*. (5th Edn), Lippincott Williams & Wilkins, Philadelphia, pp: 5-35.
3. Wolfs RC, Klaver CC, Vingerling JR, Grobbee DE, Hofman A, et al. (1997) Distribution of central corneal thickness and its association with intraocular pressure: The Rotterdam Study. *Am J Ophthalmol* 123: 767-772.

Corresponding author: Dr. Pragati Garg, Professor and Head, Department of Ophthalmology, Era's Lucknow Medical College and Hospital, Lucknow, UP, India, E-mail: drpragati89@gmail.com

Citation: Garg P. (2019) Are the Ocular Biometric Factors Important for Intraocular Pressure Assessment? *Ophthalmol Clin Res*, 2(1): 40-41.

Copyright: ©2019 Garg P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

4. Ventura AC, Böhnke M, Mojon DS (2001) Central corneal thickness measurements in patients with normal tension glaucoma, primary open angle glaucoma, pseudo exfoliation glaucoma or ocular hypertension. *Br J Ophthalmol* 85: 792-795.
5. Wilson LB, Quinn GE, Ying GS, Francis EL, Schmid G, et al. (2006) The relation of axial length and intraocular pressure fluctuations in human eyes. *Invest Ophthalmol Vis Sci* 47: 1778-1784.
6. Lee AJ, Saw SM, Gazzard G, Cheng A, Tan DT (2004) Intraocular pressure associations with refractive error and axial length in children. *Br J Ophthalmol* 88: 5-7.
7. Leydolt C, Findl O, Drexler W (2008) Effects of change in intraocular pressure on axial eye length and lens position. *Eye (Lond)* 22: 657-661.
8. Ehlers N, Bramsen T, Sperling S (1975) Applanation tonometry and central corneal thickness. *Acta Ophthalmol (Copenh)* 53: 34-43.
9. Bilak S, Simsek A, Capkin M, Guler M, Bilgin B (2015) Biometric and intraocular pressure change after cataract surgery. *Optom Vis Sci* 92: 464-470.
10. Tomoyose E, Higa A, Sakai H, Sawaguchi S, Iwase A, et al. (2010) Intraocular pressure and related systemic and ocular biometric factors in a population-based study in Japan: The Kumejima study. *Am J Ophthalmol* 150: 279-286.