

Dengue Infection in Pregnancy - A Near Miss Case Managed by Multidisciplinary Approach

Priyanka Thakur¹, Pooja Patil^{2*}, Veena Acharya¹, Raghunath Anant Nagvekar³ and Manish Rajput⁴

¹Consultant, Rajasthan Hospital, Department - Obstetrics and Gynecology, Rajasthan Hospital, Jaipur, India

²Head of Department - Obstetrics and Gynecology, L.N. Medical College, Bhopal, India

³Department of Interventional Radiology, Rajasthan Hospital, Jaipur, India

⁴Consultant Interventional Radiology, Rajasthan Hospital Jaipur, India.

Received June 07, 2023; Accepted June 12, 2023; Accepted June 15, 2023

ABSTRACT

Introduction: Dengue infection during pregnancy is challenging due to complications associated with it especially the high risk of maternal bleeding.

Case report: A full term pregnant women was diagnosed with dengue with warning signs (DWS). Emergency Lower Segment Cesarean section (LSCS) was done during the critical phase of the disease due to fetal distress. The patient had post-partum hemorrhage which was managed by uterotonic drugs, blood transfusions, Fresh Frozen Plasma (FFP). After 8 h of LSCS, she had bleeding through scar site and subcuticular drain site for which blood and blood products given and woman managed in Intensive Care Unit (ICU) by multi-disciplinary team. On Day 2 of LSCS, intra-abdominal bleeding was diagnosed as woman had abdominal distension. Bilateral uterine artery embolization (UAE) was done along with supportive measures and gradually the bleeding stopped and the woman was discharged on D11 of LSCS.

Conclusion: Dengue hemorrhagic fever during pregnancy requires a multidisciplinary approach. Timely intervention and aggressive management are the key to its positive outcome. Uterine artery embolization is a lifesaving procedure in the patients where there is bleeding due to thrombocytopenia or dengue hemorrhagic fever.

Keywords: Thrombocytopenia, Postpartum hemorrhage, Uterine artery embolization

INTRODUCTION

Dengue is an *Aedes Aegypti* mosquito borne infection caused by Flavivirus serotypes 1,2,3,4. 40% of world's population lives in dengue prone area and World Health Organization (WHO) estimates that about 100 million infection and 2200 deaths occur every year [1]. In 2009, WHO revised the classification of dengue infection into dengue without warning sign (d-w), dengue with warning signs (d+w) and severe dengue fever (sdf) [2]. The diagnosis and management of dengue infection in pregnancy is challenging because the physiological hemodilution of pregnancy masks the hemoconcentration associated with dengue. Delay in diagnosis also occurs due to overlapping signs and symptoms like thrombocytopenia, impaired liver functions, fever, ascites which are also associated with other conditions in pregnancy. Serology helps in confirming the diagnosis [3]. Liver is not the primary target of dengue, but raised transaminase level, acute fulminate hepatitis and hepatic failure have been seen [4]. Dengue infection in

pregnant women may be associated with adverse outcomes such as maternal mortality, stillbirth, and neonatal mortality. There was no significant statistical association established between maternal Dengue infection and the outcomes of preterm birth, maternal bleeding, low birth weight in neonates, and risk of miscarriage [5]. Some studies have reported vertical transmission which may cause neonatal thrombocytopenia [6,7].

Corresponding author: Pooja Patil, Department - Obstetrics and Gynecology, L.N. Medical College, Bhopal, India, Tel: 9755536795; E-mail: poojagyneccpatil@gmail.com

Citation: Thakur P, Patil P, Acharya V, Nagvekar RA & Rajput M. (2023) Dengue Infection in Pregnancy - A Near Miss Case Managed by Multidisciplinary Approach. J Womens Health Safety Res, 7(2): 307-310.

Copyright: ©2023 Thakur P, Patil P, Acharya V, Nagvekar RA & Rajput M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CASE REPORT

23-year-old primigravida was referred from periphery with dengue NS1 antigen positive and platelet count of 49000. She had complaints of fever with nausea and vomiting and epigastric pain for 3 days. At the time of admission, she was

febrile (102°F), pulse rate was 114 per minute and blood pressure was 92/60mmHg. On abdominal examination - Uterus was term size, single live fetus with cephalic presentation. Non-stress test (NST) was normal. On per vaginal examination cervical os was closed (**Tables 1 & 2**).

Table 1. Laboratory Investigations.

	D1 of admission	D2 of admission	D3 of admission/ POD1	POD2	POD3
Hemoglobin (Hb) G/dl	10.6	10.2	9.8	6.0	8.1
Hematocrit (%)	29.4	29.0	31.2	16.1	21.9
TLC (/mm ³)	5510	6100	16910	21510	18,940
Platelets (/ul)	44,000	29,000	20,000	39,000	56,000
CRP (mg/l)	10.8	-	56.4	197.4	114.2.
Liver functions					
AST(u/l)	112.7		488.9	156.2	112.3
ALT(u/l)	75.3		136.4	66.3	59.4
ALP(u/l)	322		163.0	174.5	104.2
Coagulation profile					
APTT			47.2	39.4	33.6
PT/INR			15/1.12	13.7/1.01	15.4/1.15
D-dimer (ug/ml)	3907.13		1193.38	1157.9	987.6

POD 1: Post-operative Day 1; POD 2: Post-operative Day 2; TLC: Total Leukocyte Count; CRP: C-Reactive Protein; AST: Aspartate Transaminase; ALT: Alanine Transaminase; ALP: Alkaline Phosphatase; APTT: Activated Partial Thromboplastin Clotting Test; PT: Prothrombin Time; INR: International Normalized Ratio

Table 2. Blood & Blood products given.

Therapy given	D1 of admission	D2 of admission	D3 of admission/POD1	POD2	POD3
Packed cell (ml)	None	None	700	1400	700
Fresh frozen plasma (ml)	None	None	1000	1400	500
Cryoprecipitate (ml)	None	None	None	120	120
Thromocyte concentrate apheresis (unit)	None	2 unit	None	2	1

Laboratory investigations showed hemoglobin 10gm/dl, platelets 44000, elevated liver enzymes, B negative blood group. A diagnosis of full-term pregnancy with dengue with warning signs was made. The patient was admitted in Medical ICU. On Day 2 of admission her platelets dropped to 29000 for which 2 units of Thrombocyte concentrate were transfused. NST was reactive and woman had no fresh complaints. On Day 3 of admission, fetal distress developed as was diagnosed on NST. Her Platelet count was 29000, Coagulation Profile was normal with International Normalized Ratio (INR) 1.1 and other reports were normal. With high-risk consent of hemorrhage and with arrangement of 4U Packed Red Blood Cell (PRBC) and 8 U FFP, woman was taken for Emergency LSCS under General Anesthesia. A live male child weighing 2.8 kg was delivered and was shifted to Neonatal Intensive Care Unit (NICU). There was mild to moderate Postpartum Hemorrhage (PPH) intraoperatively which was managed by 600mcg of misoprostol, 20 units of oxytocin, Methergine 100mcg intramuscular. As her blood group was B negative, there was non-availability of Platelets of the same blood group. Thus, she was transfused 2 Units of PRBC and 6 Units of FFP intraoperatively. A subcutaneous drain was kept as there was oozing from the subcutaneous tissues and woman shifted to ICU for observation. 8 h after Cesarean section there was bleeding from stitch line and drain site. Her hemoglobin dropped to 6 g/dl, Total Leukocyte Count (TLC)-21510, platelets-29000, C-Reactive Protein (CRP)-197.0, procalcitonin-65, activated Partial Thromboplastin Clotting Time (APTT)-47.2, INR-1.12. Woman was actively and aggressively managed with multi-disciplinary approach. She was transfused 6 Units (120ml) of cryoprecipitate, 6 Units (1200ml) of FFP, 4 Units (1400 ml) of PRBC, 2 units of thrombocyte concentrate (Single Donor Platelets) over the period of 12 h. On postoperative Day 2, her abdomen was found to be distended and tender. Possibility of intra-abdominal bleeding was suspected. Laboratory investigations showed hemoglobin 8.1g/dl, TLC 18940, platelets 56000, CRP- 114, INR- 1.01. There was ascites, pedal edema. X-ray chest showed bilateral pleural effusion. 2D echo was normal. Urinary output was maintained. Diuretics were given to treat fluid overload. Extensive supportive treatment was given along with higher broad spectrum antibiotic Meropenem. Bilateral venous doppler of lower limbs was done on complaints of pain in left lower limb but there was no evidence of deep venous thrombosis. Along with the opinion of intervention radiologist, Uterine artery embolization was decided as the further management to stop intra-abdominal bleeding. Selective cannulization of each uterine artery was done with micro catheter. Distal embolization was done by gel foam thus preserving the fertility of woman. Proximal embolization of bilateral uterine artery was done by coils. Post embolization woman was transfused with 2 Units (700ml) of PRBC and 2 Units (500ml) of FFP. Gradually the bleeding from stitch line and drain site stopped. The vitals of the woman gradually improved. She was transferred from

Medical ICU to high dependency unit (HDU) and then to postnatal ward. During recovery phase she was afebrile and vital parameters remained stable. Laboratory investigations showed hemoglobin 10.3g/dl, WBC count 10840, platelets count 1.86 lakhs/cumm. Antibiotics were continued for 7 days. Subcuticular drain was removed on 7th postoperative day. Anti- D was given within 72 h of delivery. Woman and her baby were discharged on post-operative Day 11.

DISCUSSION

Dengue hemorrhagic fever in pregnancy requires a high index of suspicion and quick management. Dengue fever presents in 3 phases [1].

Acute febrile phase: with high grade fever with nausea and vomiting with sore throat. Rash may and may not be present. Tourniquet test is positive.

Critical phase: when the fever subsides, the phase of capillary leak occurs due to which hemorrhagic complication develops and, in this phase, multiple organ involvement happens. There is progressive fall in White Blood Cell (WBC) count, low platelets count, rise in Packed Cell Volume (PCV) / Hematocrit. Signs of shock may appear.

Recovery phase: as the patient survives the critical phase gradually the extracellular fluid reabsorption takes place. The patient becomes hemodynamically stable, WBC rises, platelets rises and PCV reduces due to fluid reabsorption. Complications in this phase are rare. Pulmonary edema and hypervolemia may occur at this time due to excessive fluid overload. In the event of thrombocytopenia during the critical phase, severe bleeding can complicate the delivery. Induction of Labour or caesarean section should be avoided during the critical phase. If delivery is inevitable blood and blood products should be properly cross matched and saved. If platelet count is less than 50,000/ul, platelet transfusion should be done before delivery [8]. Atony of uterus should be avoided by timely use of uterotonics. In our case general anesthesia was given due to thrombocytopenia present [9]. Regional anesthesia can also be considered if the platelets count is above 70,000/ul. In our case although prophylactic transfusion was done but bleeding was observed due to defect in hemostatic system secondary to dengue infection. The complications of infection depend on the interaction of virus with the host. The virus multiplies in the cells and causes an antigen antibody reaction [10]. Thrombocytopenia results from immune mediated destruction of platelets [11]. However, it co-relates poorly with bleeding manifestations as seen in our case. It can be used as a prognostic indicator rather than as a parameter for therapeutic interventions [12]. Plasma leakage and intrinsic coagulopathy are the fundamental pathological changes responsible for the morbidity and mortality associated with dengue hemorrhagic fever [13]. In our case in spite of transfusion of blood products and fresh frozen plasma there was life threatening intra-abdominal bleeding which did not subside on

conservative treatment. Uterine artery embolization is a possible intervention which can be considered in such cases. Uterine artery embolization was successfully carried out by using gel foam of about 4mm in diameter in size in our case. Generally, it takes about 5/6 weeks for the vessels to recanalize and the blood flow returns to normal [14].

Pregnancies after UAE can have higher rates of miscarriage, and are more likely to have a cesarean section and postpartum hemorrhage. Though such complications are mentioned after the UAE but here it was a lifesaving procedure and thus must be considered in such cases.

REFERENCES

- World health organization (2009) Dengue guidelines for diagnosis, treatment, prevention and control: new edition; World Health Organization. Geneva. Switzerland pp: 147.
- Dengue in Pregnancy (2015) Management protocol fogs. Available online at: <https://www.fogs. org-2015/11>
- Sharma S, Jain S, Rajaram S (2016) Spectrum of maternal fetal outcomes during dengue infection in pregnancy: An insight. *Infect Dis Obstet Gynecol* 16: 1-4.
- Sellahewa KH (2013) Pathogenesis of dengue hemorrhagic fever and its impact on case management. *Int Sch Res Notices* 2013: 571646.
- Rathore SS, Oberoi S, Hilliard J, Raja R, Yogesh V, et al. (2022). Maternal and fetal-neonatal outcomes of dengue virus infection during pregnancy-Systematic Review. *Trop Med Int Health* 27(7): 619-629.
- Gonsalves M, Belli A (2010) The role of interventional radiology in obstetric hemorrhage. *Cardiovasc Intervent Radiol* 33: 887-895.
- Phupong V (2001) Dengue fever in pregnancy: A case report. *BMC Pregnancy Childbirth* 1: 7.
- Friedman EE, Dallah F, Harville EW, Myers L, Breart G, et al. (2014) Symptomatic dengue infection during pregnancy and infant outcomes: A retrospective cohort study. *PLoS Negl Trop Dis* 8(10): e3226.
- Witaya TP (2003) Parturient and perinatal dengue hemorrhagic fever. *Southeast Asian J Trop Med Public Health* 34(4): 797-799.
- Martina BEE, Koraka P, Osterhaus ADME (2009) Dengue virus pathogenesis: An integrated view. *Clin Microbiol Rev* 22(4): 564-581.
- Sellahewa KH (2013) Pathogenesis of dengue hemorrhagic fever and its impact on case management. Hindawi. Available online at: <http://dx.doi.org/10.5402/2013/571646>
- Wills B, Tran VN, Nguyen TH, Truong TT, Farrar J, et al. (2009) Hemostatic changes in Vietnamese children with mild dengue correlate with the severity of vascular leakage rather than bleeding. *Am J Trop Med Hyg* 81(4): 638-644.
- Kaufman RM, Gary KS, Aryeh S, Babu GW, Edward CW, et al. (2015) Platelets transfusion: A clinical guideline from the AABB. *Ann Intern Med* 162(3): 205-213.
- Ooi ET, Ganesanathan S, Anil R, Kwok FY, Sinniah M (2008) Gastrointestinal manifestations of dengue infection in adults. *Med J Malaysia* 63(5): 401-405.