

Extraosseous Accumulation of Tc-99m MDP in Lymph Node Metastases from Colon Cancer

Ya-Wen Chuang¹, Chin-Chuan Chang¹, Ying-Fong Huang^{1,2}, Yu-Chang Tyan^{2,3,4,5,6,*}

¹Department of Nuclear Medicine, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan.

²Department of Medical Imaging and Radiological Sciences, College of Health Science, Kaohsiung Medical University, Kaohsiung, Taiwan

³Center for Infectious Disease and Cancer Research, Kaohsiung Medical University, Kaohsiung, Taiwan

⁴Institute of Medical Science and Technology, National Sun Yat-sen University, Kaohsiung, Taiwan

⁵Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

⁶Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

Received September 6, 2017; Accepted September 11, 2017; Published October 26, 2017

In colon cancer, extraosseous uptake of bone-seeking radionuclide in liver metastases is commonly seen on the bone scan. However, extraosseous deposition in lymph node metastasis is rare. The authors report a case in which metastatic pulmonary lymph node metastases from colon cancer were shown to contain concentrated Tc-99m methylene diphosphonate (MDP). Computed tomography (CT) of the chest showed enlarged calcified lymph nodes at the bilateral hilar regions, and should be considered for extraosseous Tc-99m MDP accumulation.

A 60-year-old female patient with descending colon cancer (mucinous adenocarcinoma), status post chemotherapy, was referred for a scintigraphic bone scan to evaluate for metastatic disease and assist in planning therapy. The bone scan was performed at three hours after the injection of 740MBq (20 mCi) Tc-99m MDP. Findings include intense radiotracer accumulation in the anterior aspect of the right 6th rib. Additionally, soft tissue uptake is noted in bilateral pulmonary hilar regions (Figure 1a). Unenhanced CT of the chest performed 10 days earlier demonstrated metastatic lymphadenopathy and enlarged calcified lymph nodes at the bilateral hilar regions (Figure 1b). Postenhanced CT demonstrated large metastases at the liver (Figure 1c). Although accumulation of Tc-99m MDP in liver metastases with no apparent calcification occurs frequently in colon cancer, no characteristic Tc-99m MDP uptake was observed within those metastatic lesions.

Extraosseous uptake of Tc-99m MDP has been reported in various pathologic conditions [1,2]. Mucin-producing tumors contain a glycoprotein that is biochemically similar to ossifying cartilage and binds calcium salts.

Classically, mucinous adenocarcinoma tumors of the gastrointestinal tract are associated with this mechanism of Tc-99m MDP deposition in the primary and metastatic tumors [3,4]. These findings mainly correspond in location to extraosseous uptake on the bone scan, and suggest that the accumulation of Tc-99m MDP in the present case is strongly related to the calcium deposition. It also appears that Tc-99m MDP may accumulate in a calcified metastatic lesion before the calcification appears on x-ray CT [5].

COMPETING INTERESTS

The authors declare they have no conflict of interests in publishing this case study.

ACKNOWLEDGMENTS

This work was supported by Research Grants MOST 103-2320-B-037-025 from the Ministry of Science and Technology, KMU-TP105E12, KMU-TP105PR06, KMU-M106029, 105KMUOR02 and KMU-O104003 (Aim for the Top 500 Universities Grant) from Kaohsiung Medical University, and NSYSUKMU106-P011 from NSYSU-KMU Research Project, Taiwan.

Corresponding Author: Yu-Chang Tyan, Department of Medical Imaging and Radiological Sciences, Kaohsiung Medical University, Kaohsiung, Taiwan. E-mail: yctyan@kmu.edu.tw

Citation: Chuang Y W, Chang C C, Huang Y F & Tyan Y C. (2017) Extraosseous Accumulation of Tc-99m MDP in Lymph Node Metastases from Colon Cancer. Int J Med Clin Imaging, 1(2): 15-18.

Copyright: ©2017 Chuang Y W, Chang C C, Huang Y F & Tyan Y C. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. (a)The bone scan was performed after the injection of Tc-99m MDP. Findings include intense radiotracer accumulation in anterior aspect of the right 6th rib, and the soft tissue uptake is noted in bilateral pulmonary hilar regions.




Figure 1. (b)Unenhanced CT of the chest performed 10 days earlier demonstrated metastatic lymphadenopathy and enlarged calcified lymph nodes at the bilateral hilar regions.



Figure 1. (c)Postenhanced CT demonstrated the large metastases at liver.

REFERENCES

1. Tsai SC, Kao CH, Lin WY, Waanng SJ. Intestinal accumulation of Tc 99m MDP on bone e s scan. Semin Nucl Med 1999;29:80-81.

2.Takahashi T, Machida K, Honda N, Hosono M, Oku S, Osada H, Murata O, Nishimura K, Ohno H. Extraosseous accumulation of 99mTc-MDP in lymph node metastases of small cell carcinoma of the esophagus. Annals of Nuclear Medicine 2004;18:157-160.

3. Freeman L, Zuckier L. Nonosseous, nonurologic uptake on bone scintigraphy: atlas and analysis. *Semin Nucl Med* 2010;40:242-256.
4. Shepherd TM, Idakoji IA, Pampaloni MH. Incidental Detection of Gastrointestinal Stromal Tumor by Tc-99m MDP Bone Scan. *Clin Nucl Med* 2012;37:198-199.
5. Senda M, Tamaki N, Torizuka K, Fujiwara Y, Kudo M, Tochio H, Ito H, Yamaguchi H, Saiki Y, Ikekubo K. Accumulation of Tc-99m Methylene Diphosphonate in Calcified Metastatic Lesions of the Liver from Colonic Carcinoma: Comparison with Calcification on X-ray Computed Tomogram. *Clin Nucl Med* 1985;10:9-12.