# **International Journal of Medical & Clinical** Imaging

IJMCI, 1(2): 19-22



www.scitcentral.com

**Original Case Report: Open Access** 

# **Demonstration of Uterine Leiomyoma on Bone Scan**

Ya-Wen Chuang<sup>1,+</sup>, Chien-Chin Hsu<sup>2</sup>, Chin-Chuan Chang<sup>1,+</sup>, Chia-Yang Lin<sup>1</sup>, Ying-Fong Huang<sup>1,3</sup>, Yu-Chang Tyan<sup>3,4,5,6,7\*</sup>

<sup>1</sup>Department of Nuclear Medicine, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan.

<sup>2</sup>Department of Nuclear Medicine, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Taiwan.

<sup>3</sup>Department of Medical Imaging and Radiological Sciences, College of Health Science, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>4</sup>Center for Infectious Disease and Cancer Research, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>5</sup>Institute of Medical Science and Technology, National Sun Yat-sen University, Kaohsiung, Taiwan

<sup>6</sup>Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>7</sup>Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

<sup>+</sup> These authors contributed equally to this work.

Received September 27, 2017; Accepted October 4, 2017; Published October 27, 2017

# ABSTRACT

Background: Uterine Leiomyoma is a benign neoplasm that occurs from the overgrowth of smooth muscle and connective tissue in the uterus. A nuclear medicine scan is a sensitive indicator of metastatic bone diseases or other pathology. The diagnosis of uterine leiomyomas using nuclear medicine is straightforward, given the common clinical manifestations and typical imaging features.

Case presentation: A 45-year-old woman with a history of nasopharyngeal cancer was referred for a bone scan to assess for bone metastasis. Abnormal accumulation of radiotracer was shown in the pelvic cavity.

Conclusion: The differential diagnosis of the abnormal radioactivity included a sacral metastasis and a pelvic tumor with uptake of bone-seeking tracers. The ultrasonography of the pelvic cavity showed a huge uterine myoma with calcifications and cystic components. After the testing, a total abdominal hysterectomy was performed and multiple leiomyomata were confirmed by histopathologic examination.

Keywords: Bone Scan, Uterine Leiomyoma, Pelvic Tumor

# **INTRODUCTION**

Uterine leiomyoma has been reported as an occasional cause of extraskeletal uptake of a bone-seeking agent in the pelvis [1-5]. The cause of tumor uptake of boneseeking tracers is thought to be related to dystrophic calcification [6]. In dystrophic calcification, the mechanism appears to be the loss of intracellular calcium in the injured cells resulting in an increased calciumbinding capacity.

# CASE REPORT

The bone scan was performed three hours after an intravenous injection of 740MBq (20mCi) Tc-99m methylene diphosphonate (MDP) into a 45-year-old woman with a history of nasopharyngeal cancer. Abnormal accumulation of radiotracer was shown in the pelvic cavity (arrow) and it was separated from a concave urinary bladder (Figure 1). The differential diagnosis of the abnormal radioactivity included sacral metastasis and

SciTech Central Inc. Int J Med Clin Imaging (IJMCI) a pelvic tumor with uptake of bone-seeking tracers. The patient presented with vaginal spotting, dyspareunia, dysmenorrhea, and urinary urgency. Physical examination showed pale conjunctiva and a palpable mass in the lower abdomen. Laboratory data revealed a reduction of hemoglobin to 8.6 g/dl (normal range, 12 to 14 g/dl).

Corresponding Author: Yu-Chang Tyan, Department of Medical Imaging and Radiological Sciences, Kaohsiung Medical University, Kaohsiung, Taiwan. E-mail: yctyan@kmu.edu.tw

Citation: Chuang Y W, Hsu C C, Chang C C, Lin C Y, Huang Y F, et al. (2017) Demonstration of Uterine Leiomyoma on Bone Scan. Int J Med Clin Imaging, 1(2): 19-22.

Copyright: ©2017 Chuang Y W, Hsu C C, Chang C C, Lin C Y, Huang Y F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### International Journal of Medical & Clinical Imaging, 1(2): 19-22

Then ultrasonography was arranged to evaluate the pelvic mass, which demonstrated a heterogeneous hypo-echoic mass with calcifications and cystic components. It originated from the uterus and was approximately  $11.0 \times 10.0 \times 9.7$  cm in size (**Figure 2**). The patient underwent total abdominal hysterectomy for the uterine mass. The uterus measured  $16.6 \times 12.1 \times 7.1$  cm and weighed 801 gm. The histopathologic diagnosis revealed multiple intramural and subserous leiomyomata of the uterus. The largest measured up to 7 cm in diameter. The bone scan was repeated after the hysterectomy. The previous extraskeletal uptake of MDP in the pelvic cavity disappeared and the shape of the urinary bladder was normal (**Figure 3**).

## DISCUSSION

Incidental extraskeletal uptake of bone-seeking tracers on bone scans is occasionally seen. Soft tissue uptake must be differentiated from bone pathology [7]. Additionally, normal renal uptake and excretion of bone-seeking agents allow for gross visualization of the genitourinary system and provide useful diagnostic information. Retroperitoneal or pelvic masses should be suspected if abnormalities in the location or contour of urinary bladder are present [7-11]. Obtaining a relevant clinical history and laboratory data is helpful to clarify the incidental extraskeletal uptake of bone-seeking tracers.

## **COMPETING INTERESTS**

The authors declare they have no conflict of interests in publishing this case study.

#### ACKNOWLEDGMENTS

This work was supported by Research Grants MOST 103-2320-B-037-025 from the Ministry of Science and Technology, KMU-TP105E12, KMU-TP105PR06, KMU-M106029, 105KMUOR02 and KMU-O104003 (Aim for the Top 500 Universities Grant) from Kaohsiung Medical University, and NSYSUKMU106-P011 from NSYSU-KMU Research Project, Taiwan.



Figure 1. Abnormal accumulation of radiotracer was showed in the pelvic cavity (arrow) and it was separated from a concave urinary bladder.

International Journal of Medical & Clinical Imaging, 1(2): 19-22

Chuang Y W, Hsu C C, Chang C C, Lin C Y, Huang Y F, et al



**Figure 2.** The ultrasonography of pelvic cavity demonstrated a heterogeneous hypo-echoic mass with calcifications and cystic components. It originated from the uterus and was approximately 11.0 x 10.0 x 9.7 cm in size.



Figure 3. The previous extraskeletal uptake of MDP in the pelvic cavity disappeared and the shape of urinary bladder was normal.

#### REFERENCES

1. Ell PJ, Breitfellner G, Meixner M. Technetium-99m-HEDP concentration in calcified myoma. J Nucl Med. 1976; 17:323-324.

2. Stadalnik RC. Non-skeletal accumulation of bone seeking agents: pelvis. Sem Nucl Med. 1988; 18:159-161.

3. Rohrer DG, Williamson BR, Teates CD. Calcified uterine leiomyoma simulating metastatic disease on bone scan. South Med J. 1988; 81:651-652.

4. Inoue H, Aizawa N, Mizuno T, et al. A large degenerated subserous leiomyoma of the uterus: uncommon scintigraphic and ultrasonographic findings. Ann Nucl Med. 1989; 3:55-57.

5. Buyukdereli G, Guney IB, Kibar M. Bone scan findings of sacrococcygeal chordoma and uterine leiomyoma. J Women's Imaging. 2004; 6:171-174.

6. Elgazzar A. H. Orthopedic Nuclear Medicine. Berlin Heidelberg. Springer-Verlag, 2004, pp197-198.

7. Loutfi I, Collier BD, Mohammed AM. Nonosseous abnormalities on bone scans. J Nucl Med Technol. 2003; 31:149-153.

8. Shanley DJ, Buckner AB, Alexander HG. Unusual pelvic activity found on bone scan. Clin Nucl Med. 1991; 16:453-454.

9. Ackerman L, Shirazi P, Van Drunen M. Bladder displaced by stool-distended rectosigmoid colon presenting on bone scan as a pelvic soft tissue mass. Clin Nucl Med. 1997; 22:733-734.

10. Al Naeem AN, Powe J, Bakheet S, et al. Displaced urinary bladder creating an unusual pattern on bone scan mimicking disease. Clin Nucl Med. 1999; 24:137-138.

11. Lin ST, Sohn MH, Park SA. Curious radioactivity in the lower abdomen on bone scintigraphy: displacement of the urinary bladder by an incidentally diagnosed uterine myoma. Clin Nucl Med. 2000; 25:824-825.