Journal of Infectious Diseases & Research

JIDR, 3(S3): 09 www.scitcentral.com



ISSN: 2688-6537

Abstract: Open Access

Immunity to Glycan Alpha-Gal and COVID-19: Possibilities for Disease Control and Prevention

José de la Fuente^{1*,2}, José Miguel Urra^{3,4}, Elisa Ferreras-Colino¹, Marinela Contreras⁵, Carmen M Cabrera^{3,4}, Isabel G Fernández de Mera¹, Margarita Villar^{1,6}, Alejandro Cabezas-Cruz⁷ and Christian Gortázar¹

*¹SaBio, Instituto de Investigación en Recursos Cinegéticos IREC-CSIC-UCLM-JCCM, Spain

²Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, USA

³Immunology, Hospital General Universitario de Ciudad Real, Spain

⁴Medicine School, Universidad de Castilla la Mancha (UCLM), Spain

⁵Interdisciplinary Laboratory of Clinical Analysis, Interlab-UMU, Regional Campus of International Excellence Campus Mare Nostrum, University of Murcia, Espinardo, Spain

⁶Biochemistry Section, Faculty of Science and Chemical Technologies, Regional Centre for Biomedical Research (CRIB), University of Castilla-La Mancha, Spain

⁷UMR BIPAR, INRAE, ANSES, Ecole Nationale Vétérinaire d'Alfort, Université Paris-Est, France.

Published December 12th, 2020

ABSTRACT

The coronavirus disease 19 (COVID-19) pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has affected millions of people worldwide. The characterization of the immunological mechanisms involved in disease symptomatology and protective response is important to advance in disease control and prevention. Humans evolved by losing the capacity to synthesize the glycan Galα1-3Galβ1-(3)4GlcNAc-R (alpha-Gal), which resulted in the alpha-Gal syndrome associated with tick salivary glycoproteins with alpha-Gal modifications that after tick bite can induce in some individuals the production of high levels of anti-alpha-Gal IgE antibodies that mediate delayed anaphylaxis to mammalian meat and immediate anaphylaxis to tick bites. In contrast, humans can develop of a protective response against pathogens containing this modification on membrane proteins mediated by anti-alpha-Gal IgM/IgG antibodies naturally produced in response to bacterial microbiota. In addition to anti-alpha-Gal antibody-mediated pathogen opsonization, this glycan induces various immune mechanisms such as B-cell maturation, macrophage response, activation of the complement system, innate immune pathway and TLR-mediated anti-inflammatory responses that have shown protection in animal models against infectious diseases. Based on these results, we hypothesized that the immune response to alpha-Gal may contribute to the control of COVID-19. To address this hypothesis, we characterized the antibody response to alpha-Gal in patients at different stages of COVID-19 and healthy control individuals. The results showed that while the inflammatory response and the anti-SARS-CoV-2 IgG antibody titers increased, reduction in anti-alpha-Gal IgE, IgM and IgG antibody titers and alteration of anti-alpha-Gal antibody isotype composition correlated with COVID-19 severity. These results suggested that the inhibition of the alpha-Gal-induced immune response may translate into more aggressive viremia and severe disease inflammatory symptoms. These results support our proposal of developing interventions such as probiotics/postbiotics based on commensal bacteria with alpha-Gal epitopes to modify the microbiota and increase the alpha-Galinduced protective immune response and reduce the severity of COVID-19.

Keywords: Coronavirus, Microbiota, Immune pathway, Inflammatory response

Corresponding author: José de la Fuente, SaBio, Instituto Investigación en Recursos Cinegéticos IREC-CSIC-UCLM-JCCM, Ronda Toledo s/n. 13005 Ciudad Real, Spain. jose_delafuente@yahoo.com

Citation: de la Fuente J, Urra JM, Ferreras-Colino E, Contreras M, Cabrera CM, et al. (2020) Immunity to Glycan Alpha-Gal and COVID-19: Possibilities for Disease Control and Prevention. J Infect Dis Res, 3(S3): 09.

Copyright: ©2020 de la Fuente J, Urra JM, Ferreras-Colino E, Contreras M, Cabrera CM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.