Journal of Ageing and Restorative Medicine

JARM, 1(1): 7-12 www.scitcentral.com Sc Tech a quantum to research.

Original Review: Open Access

Molecular Imaging of Stem Cell Therapy for Cerebral Ischemia in the Aged Brain

Andrei Gresita¹, Mircea Popescu¹, Andrei Olaru⁴, Denissa-Greta Olaru⁵, Otilia Margaritescu⁶ and Bogdan Catalin^{5*}

¹Department of Functional Sciences, Center of Clinical and Experimental Medicine, University of Medicine and Pharmacy of Craiova, Romania

²Department of Ophthalmology, University of Medicine and Pharmacy Craiova, Romania

³Medlife Clinic, Department of Ophthalmology, Craiova, Romania

Department of Neurosurgery, University of Medicine and Pharmacy Craiova, Romania

¹⁵ Experimental Research Centre for Normal and Pathological Aging, University of Medicine and Pharmacy of Craiova, Romania

Received February 22, 2018; Accepted April 2, 2018; Published June 17, 2018

ABSTRACT

Cell therapy has emerged as a promising approach to improve recovery after stroke. However, progress toward the development of efficient cell based therapies for ischemic stroke has been so far disappointing. The main problem remains the poor understanding of the dynamics of cellular interactions between host cells and administered cells for therapeutic purposes. The pathophysiological evolution of stroke events is driven by complex cellular interactions between many different cell types whose sequential recruitments have been insufficiently documented due to the lack of non-invasive imaging modalities. Specifically, the interplay between host neuroinflammation, which is considered to be a major obstacle to exogenous-mediated neuronal precursor cells, and exogenously administered stem cells, remains virtually unknown. Phagocytosis of dead and dying neurons and neuronal debris is beneficial in part because it reduces inflammation. However, microglia can also phagocytose live neurons, live neuronal progenitors or live stressed-but-viable neurons like those presumably occurring after transplantation causing death of the engulfed cell and compromising cell therapy using neuronal precursor cells. This lack of basic knowledge critically limits the optimization of timing and route of administration of stem cells for therapeutic purpose. In this review, we aim to decipherthe dynamic interplay between host neuroinflammation and therapeutic stem cells for regeneration after stroke by two non-invasive molecular imaging approaches (two photon microscopy and magnetic resonance imaging at high magnetic field) and their impact on behavioural outcome measures.

Keywords: Aging, Cell therapy, Imaging, Two photon microscopy, Magnetic resonance imaging

INTRODUCTION

Stroke is the third most common cause of death and the leading cause of disabilities worldwide. The damages after stroke generally have two different causes, the primary and secondary insult. The primary insult is mediated by the ischemic event itself and leads to oxygen and glucose deprivation, crucial substrates for the survival of the brain tissue, whereas the secondary insult is caused by the inflammatory response after stroke and mainly affects the penumbra, a brain area with a slightly compromised blood-supply that surrounds the ischemic core. Up to date, tPA (tissue plasminogen activator) is the only FDA approved therapeutic treatment for ischemic stroke [1]. But due to severe side effects and because of its very limited time window it is only applicable to less than 10 % of all stroke

patients [1], thus indicating an urgent need for alternative treatments.

Up to now, studies of many brain diseases have limited their

*Corresponding Author: Bogdan Catalin, MD, PhD, Department of Functional Sciences, Center of Clinical and Experimental Medicine, University of Medicine and Pharmacy of Craiova, Romania; Email: bogdan.catalin@yahoo.co.uk

Citation: Gresita A, Popescu M, Olaru A, Olaru D G, Margaritescu O et al., (2018) Molecular Imaging of Stem Cell Therapy for Cerebral Ischemia in the Aged Brain. J Ageing Restor Med, 1(1): 7-12.

Copyright: ©2018 Gresita A, Popescu M, Olaru A, Olaru D G, Margaritescu O et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

attention on the focal lesion, ignoring the wider impact of the directly affected territory on other brain regions. In this review we approach the brain as a rather horizontally organized, complex network where focal lesions will have far ranging effects on distal brain regions and their corresponding functions. This approach provides new insight into the mechanisms and reasons underlying functional disturbances and opens new avenues for future diagnosis. Specifically, through this novel approach we expect to make predictions with regard of the efficacy of stem cell therapy of ischemic stroke by gaining insight into of the time window of application and route of administration of stem cells on a background of neuroinflammation in aged subjects and to develop an experimental, pre-clinical rationale for the development of cell based therapeutic strategies for the treatment of stroke and neurodegenerative diseases.

Cell Therapy has Emerged as a Promising Approach to Improve Recovery after Stroke

However, progress toward the development of efficient cell based therapies for ischemic stroke has been so far disappointing. The main problem remains the poor understanding of the dynamics of cellular interactions between host cells and administered cells for therapeutic purposes. The pathophysiological evolution of stroke events is driven by complex cellular interactions between many different cell types whose sequential recruitments have been insufficiently documented due to the lack of non-invasive imaging modalities. Specifically, the interplay between host neuroinflammation, which is considered to be a major obstacle to exogenous-mediated neuronal precursor cells, and exogenously administered stem cells, remains virtually unknown.

Sub toxic insults such as inflammation, stressed but viable neurons may reversibly expose, eat-me signals, may expose the "eat-me" signal phosphatidylserine (PS) on neuronal surface. Activated microglia detect exposed 'eat-me' signals and engulfment of neurons or parts of neurons exposing such signals follows. This process has been coined primary phagocytosis or "phagoptosis" [2,3]. Toxic neuronal insults, such as dying neurons after stroke, irreversibly expose the "eat-me" signal recognized by primed microglia resulting in the phagocytosis if dead neurons or the so-called secondary phagocytosis [4]. There are "eat me" signals other than phosphatidylserine. During brain development and normal functioning in the adult, immune molecules, including complement proteins, C1q and C3, have emerged as critical mediators of synaptic refinement and plasticity via C3dependent microglial phagocytosis of synapses. Following an acute injury such as stroke the apoptotic neurons release a chemotactic signal such as fractalkine/CX3CL1 [5,6] and microglia expressing the fractalkine receptor(CX3CR1), promotes phagocytosis of apoptotic cells expressing CX3CL1 [5,7]. The complement components C1q and C3, which are produced by microglia and astrocytes, may induce phagocytosis by binding opsonized/altered neuronal surfaces. In this process C1q promotes the conversion of C3, expressed by microglia, to C3b. C3b then opsonizes neurons and is recognized through complement receptor 3 (CR3) expressed by activated microglia.

Phagocytosis of dead and dying neurons and neuronal debris is beneficial in part because it reduces inflammation. However, microglia can also phagocytose live neurons, live neuronal progenitors or live stressed-but-viable neurons like those presumably occurring after transplantation causing death of the engulfed cell and compromising cell therapy using neuronal precursor cells.

Mild neuroinflammation can be beneficial for regenerative events aimed at functional restoration after stroke [8]. Cell therapy itself can be used during the first week post-stroke to limit neuroinflammation in animal models [9-11]. However, persistent post-stroke neuroinflammation results in decreased proliferation of the newly born NPCs and ineffective integration into the circuitry of the re-organized brain area [12]. Moreover, a number of studies have demonstrated that neuroinflammatory processes can induce apoptosis in NPCs and immature neurons [13,14] and decrease the efficacy of both stroke-induced neurogenesis and exogenously supported neurogenesis. This hypothesis is supported by findings from studies using anti-inflammatory drugs such as indomethacin or minocyclin block microgliainduced apoptosis of NPCs in a pro-inflammatory milieu [15-17]. More recently studies directly implicated TNFa produced by lipopolysaccharide-activated microglia is as a key determinant in microglia induced-apoptosis in mouse NPCs in vitro and in vivo [18].

This lack of basic knowledge critically limits the optimization of timing and route of administration of stem cells for therapeutic purpose. In the present review, we aim to investigate the dynamic interplay between host neuroinflammation and therapeutic stem cells for regeneration after stroke by combining non-invasive molecular imaging modalities (two photon microscopy (2P-LSM) and magnetic resonance imaging (MRI) at high magnetic field) with in vivo neurophysiological and behavioural outcome measures. Up to now, studies of many brain diseases have limited their attention on the focal lesion, ignoring the wider impact of the directly affected territory on other brain regions. In the proposed project we approach the brain as a rather horizontally organized, complex network where focal lesions will have far ranging effects on distal brain regions and their corresponding functions. This approach provides new insight into the mechanisms and reasons underlying functional disturbances and opens new avenues for future diagnosis. Specifically, through this novel approach we expect to make significant improvements in the efficacy of stem cell therapy of ischemic stroke by optimizing the time window of application and route of administration of stem cells on a

background of neuroinflammation in aged subjects. Finally, we expect to decipher experimental, pre-clinical rationale for the development of cell based therapeutic strategies for the treatment of stroke and neurodegenerative diseases.

Role of the Immune Response on Tissue Protection and Recovery

We have explored the role of the polarization phenotype of microglia and macrophages for the neuronal survival and functional deficit, functional improvement respectively. Injecting the microRNA 124 (mir-124), we found a substantial shift from the pro- to the anti-inflammatory phenotype, combined with a significantly higher neuronal survival. In parallel, there was a tight correlation between decrease of neurological deficit and increase of anti-inflammatory (M2) phenotype [19,20].

Stem Cell Therapies in Preclinical Models of Stroke Associated with Aging

Stroke has limited treatment options, demanding a vigorous search for new therapeutic strategies. Initial enthusiasm to stimulate restorative processes in the ischemic brain by means of cell-based therapies has meanwhile converted into a more balanced view recognizing impediments related to unfavorable environments that are in part related to aging processes. Since stroke afflicts mostly the elderly, it is highly desirable and clinically important to test the efficacy of cell therapies in aged brain microenvironments. Although widely believed to be refractory to regeneration, recent studies done by us using both neural precursor cells and bone marrow-derived mesenchymal stem cells for stroke therapy suggest that the aged rat brain is not refractory to cell-based therapy, and that it also supports plasticity and remodeling [21,22]. Yet, important differences exist in the aged compared with young brain, i.e., the accelerated progression of ischemic injury to brain infarction, the reduced rate of endogenous neurogenesis and the delayed initiation of neurological recovery. Pitfalls in the development of cell-based therapies may also be related to age-associated comorbidities, diabetes e.g., or hyperlipidemia, which may result in maladaptive or compromised brain remodeling, respectively. These agerelated aspects should be carefully considered in the clinical translation of restorative therapies [23]. In the last 3 years, upon acquiring a 2P-LSM microscope, now the gold standard of microglia reaction [24-26], our focus has been on in vivo monitoring of the efficacy of anti-inflammatory therapies by quantifying microglia reaction to focal (Figure 1) or global lesions [27].

Functional Dynamics of Stem Cell Grafts

For almost 20 years, our research has focused on stem cell based regeneration of cerebral disorders. Based on the longstanding extensive work on stroke pathophysiology we have concentrated on various aspects of stem cell dynamics such migration, proliferation, vitality, and neuronal as differentiation. Using high resolution high field in vivo MRI at 7Tesla, we were the first to monitor directed stem cell migration towards the ischemic target in noninvasive longitudinal imaging studies [28]. Optimization of the bioluminescence imaging (BLI) for brain studies and identification of maximally sensitive luciferases allowed the in vivo assessment of stem cell graft vitality during longitudinal investigations [29,30]. Combination of this BLI protocol with 19F-MRI solved the controversial debate over the best suited location for graft vitality: it has been shown that the vitality of grafts in the peri-infarct area is equivalent to that of grafts in healthy tissue, thus allowing to safely implant stem cells close to but outside of the ischemic target zone [31]. From further studies, it was concluded that the graft size, but not the immune response to the grafting, influences the vitality. This resulted in recommendation of maximal graft size [32-34].

Differentiation and Integration of Neural Stem Cell Grafts

In a major project the temporal differentiation profile of human neural stem cells (NSCs) after grafting was identified. Stem cells were transduced to express imaging reporters under cell-specific control for the in vivo monitoring of selected stages of differentiation. For this purpose, luciferases for detection by BLI and fluorescent markers for immunohistological validation of in vivo observations by immunohistochemistry were chosen as imaging reporters. Setting the imaging reporters under gene control of early (Doublecortin) and late (Synapsin) neuronal differentiation, a time line was generated which permits for the first time the direct correlation between functional improvement, assessed by behavioral test, and the neuronal differentiation of the graft and its capacity to integrate. This time line further allows a reliable discrimination between discussed modes of action of stem cells for a functional improvement: bystander effect (during early phase before neuronal differentiation) and tissue replacement (after neuronal differentiation and during cell integration) [32]. In a further study the graft innervation by the host tissue was unraveled. Whole-brain light sheet microscopy of a translucent brain permitted at cellular resolution the afferent connection from far-ranging neuronal sites to the graft. The corresponding anatomical assignment of the connecting host tissue regions was achieved by coregistration of the wholebrain light sheet data set with high-resolution anatomical MRI and mouse brain atlas [35].



Figure 1. Two-Photon microscopy time laps images of an anesthetizes mouse right somatosensory cortex, taken over a period of three days(A)show green microglia of a transgenic mice (EGFP was bonded to exon1 of the CX3CR1 locus[7]), (B, C) reacting to a 2 mm deep stab wound injury done using a 30G needle. Every experiment has a 20 min baseline (-20 min) for every group investigated . After this baseline, the injury is done, and with the same needle a 1 µl red solution, obtained by adding Sulfurodamine B to vehicle is directly injected in the cortex. (D).Quantification of the number of microglia over time, after the lesion, shows a delay microglia proliferation in the treated animals compared with the controls. (Mean±SEM, *<0.05, **p<0.01).

Neuronal Networks and Post-Stroke Recovery

The understanding of cerebral diseases has moved from the focal attention on the lesion itself to more global concepts of far ranging effects in neuronal networks during the past years. Invasive approaches of experimental investigations are either limited to a few local recordings such as electrophysiology or permit only a description of a single time point because of invasiveness of the method. Modern noninvasive molecular imaging modalities not only allow the investigation of dynamic cellular processes (see above) but have recently begun to open new doors to generate temporal profiles of dynamic processes relating to both structural and functional connectivity networks in the brain. Animal studies in rodent models can now be investigated with impressive sensitivity and resolution, thus unravelling mechanisms and interactions between various components such as different cell types during lesion development and (therapeutically intervened) outcome. Thus, investigations by Dijkhuizen and colleagues have shown the disturbance of the functional networks after stroke induction, affecting the ischemic hemisphere but also extending the deficits into the transhemispheric connections of homotopic areas of the motor and sensory cortex [36,37].

REFERENCES

- 1. Tobin MK, Bonds JA, Minshall RD, Pelligrino DA, Testai FD, et al. (2014) Neurogenesis and inflammation after ischemic stroke: What is known and where we go from here. J Cereb Blood Flow Metab 34: 1573-1584.
- Fuhrmann M, Bittner T, Jung CK, Burgold S, Page RM, et al. (2010) Microglial cx3cr1 knockout prevents neuron loss in a mouse model of alzheimer's disease. Nat Neurosci 13: 411-413.
- Neher JJ, Neniskyte U, Zhao JW, Bal-Price A, Tolkovsky AM, et al. (2011) Inhibition of microglial phagocytosis is sufficient to prevent inflammatory neuronal death. J Immunol 186: 4973-4983.
- 4. Brown GC, Neher JJ (2014) Microglial phagocytosis of live neurons. Nat Rev Neurosci 15: 209-216.
- Noda M, Doi Y, Liang J, Kawanokuchi J, Sonobe Y, et al. (2011) Fractalkine attenuates excitoneurotoxicity via microglial clearance of damaged neurons and antioxidant enzyme heme oxygenase-1 expression. J Biol Chem 286: 2308-2319.
- 6. Truman LA, Ford CA, Pasikowska M, Pound JD, Wilkinson SJ, et al. (2008) Cx3cl1/fractalkine is released from apoptotic lymphocytes to stimulate macrophage chemotaxis. Blood 112: 5026-5036.
- 7. Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, Littman DR (2000) Analysis of fractalkine receptor cx(3)cr1 function by targeted deletion and green fluorescent protein reporter gene insertion. Mol Cell Biol 20: 4106-4114.
- Perry VH, Nicoll JA, Holmes C (2010) Microglia in neurodegenerative disease. Nat Rev Neurol 6: 193-201.

- 9. Prockop DJ, Oh JY (2012) Mesenchymal stem/stromal cells (mscs): Role as guardians of inflammation. Mol Ther 20: 14-20.
- Bedi SS, Hetz R, Thomas C, Smith P, Olsen AB, et al. (2013) Intravenous multipotent adult progenitor cell therapy attenuates activated microglial/macrophage response and improves spatial learning after traumatic brain injury. Stem Cells Transl Med 2: 953-960.
- 11. Dooley D, Vidal P, Hendrix S (2014) Immunopharmacological intervention for successful neural stem cell therapy: New perspectives in cns neurogenesis and repair. Pharmacol Ther 141, 21-31.
- 12. London A, Cohen M, Schwartz M (2013) Microglia and monocyte-derived macrophages: Functionally distinct populations that act in concert in cns plasticity and repair. Front Cell Neurosci 7: 34.
- Ekdahl CT, Claasen JH, Bonde S, Kokaia Z, Lindvall O (2003) Inflammation is detrimental for neurogenesis in adult brain. Proc Natl Acad Sci USA 100: 13632-13637.
- 14. Kyritsis N, Kizil C, Brand M (2014) Neuroinflammation and central nervous system regeneration in vertebrates. Trends Cell Biol 24: 128-135.
- 15. Monje ML, Toda H, Palmer TD (2003) Inflammatory blockade restores adult hippocampal neurogenesis. Sci 302: 1760-1765.
- 16. Hoehn BD, Palmer TD, Steinberg GK (2005) Neurogenesis in rats after focal cerebral ischemia is enhanced by indomethacin. Stroke 36: 2718-2724.
- 17. Liu Z, Fan Y, Won SJ, Neumann M, Hu D, et al. (2007) Chronic treatment with minocycline preserves adult new neurons and reduces functional impairment after focal cerebral ischemia. Stroke 38: 146-152.
- Guadagno J, Xu X, Karajgikar M, Brown A, Cregan SP (2013) Microglia-derived tnfα induces apoptosis in neural precursor cells via transcriptional activation of the bcl-2 family member puma. Cell Death Dis 4: e538.
- 19. Hamzei Taj S, Kho W, Aswendt M, Collmann FM, Green C, et al. (2016) Dynamic modulation of microglia/macrophage polarization by mir-124 after focal cerebral ischemia. J Neuroimmune Pharmacol 11: 733-748.
- 20. Hamzei Taj S, Kho W, Riou A, Wiedermann D, Hoehn M (2016) Mirna-124 induces

neuroprotection and functional improvement after focal cerebral ischemia. Biomaterials 91: 151-165.

- 21. Balseanu AT, Buga AM, Catalin B, Wagner DC, Boltze J, et al. (2014) Multimodal approaches for regenerative stroke therapies: Combination of granulocyte colony-stimulating factor with bone marrow mesenchymal stem cells is not superior to g-csf alone. Frontiers Aging Neurosci 6: 130.
- 22. Buga AM, Scheibe J, Moller K, Ciobanu O, Posel C, et al. (2015) Granulocyte colony-stimulating factor and bone marrow mononuclear cells for stroke treatment in the aged brain. Curr Neurovascul Res 12: 155-162.
- Popa-Wagner A, Buga AM, Doeppner TR, Hermann DM (2014) Stem cell therapies in preclinical models of stroke associated with aging. Front Cell Neurosci 8: 347.
- 24. Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, et al. (2005) Atp mediates rapid microglial response to local brain injury in vivo. Nat Neurosci 8: 752-758.
- 25. Nimmerjahn A, Kirchhoff F, Helmchen F (2005) Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. Sci 308: 1314-1318.
- 26. Catalin B, Cupido A, Iancau M, Albu CV, Kirchhoff F (2013) Microglia: First responders in the central nervous system. Romanian journal of morphology and embryology = Revue roumaine de morphologie et embryologie 54: 467-472.
- 27. Catalin B, Stopper L, Balseanu TA, Scheller A (2017) The in situ morphology of microglia is highly sensitive to the mode of tissue fixation. J Chem Neuroanat 86: 59-66.
- 28. Hoehn M, Küstermann E, Blunk J, Wiedermann D, Trapp T, et al. (2002) Monitoring of implanted stem cell migration in vivo: A highly resolved in vivo magnetic resonance imaging investigation of experimental stroke in rat. Proceedings of the National Academy of Sciences of the USA 99: 16267-16272.
- 29. Aswendt M, Adamczak J, Couillard-Despres S, Hoehn M (2013) Boosting bioluminescence neuroimaging: An optimized protocol for brain studies. PLoS One 8: e55662.
- Mezzanotte L, Aswendt M, Tennstädt A, Hoeben R, Hoehn M, et al. (2013) Quantitative evaluation of luciferases in vivo bioluminescence imaging of neural stem cells. Mol Imaging Biol 8, 505-513.

- 31. Boehm-Sturm P, Aswendt M, Minassian A, Michalk S, Mengler L, et al. (2014) A multimodality platform to image stem cell graft survival in the naive and stroke-damaged mouse brain. Biomaterials 35: 2218-2226.
- 32. Tennstaedt A, Aswendt M, Adamczak J, Collienne U, Selt M, et al. (2015) Human neural stem cells intracerebral grafts show spontaneous early neuronal differentiation after several weeks. Biomaterials 44: 143-154.
- 33. Tennstaedt A, Mastropietro A, Nelles M, Beyrau A, Hoehn M (2015) In vivo fate imaging of intracerebral stem cell grafts in mouse brain. PLoS ONE 10: e0144262.
- 34. Vogel S, Aswendt M, Nelles M, Henn N, Schneider G, Hoehn M (2017) Initial graft size and not the innate immune response limit survival of engrafted neural stem cells. Tissue Eng Reg Med 2017.
- 35. Doerr J, Schwarz M, Wiedermann D, Leinhaas A, Jakobs A, et al. (2017) Whole-brain 3d mapping of human neural transplant innervation. Nature Communications 8: 14162.
- 36. van Meer MP, Otte WM, van der Marel K, Nijboer CH, Kavelaars A, et al. (2012) Extent of bilateral neuronal network reorganization and functional recovery in relation to stroke severity. J Neurosci 32: 4495-4507.
- 37. van Meer MP, van der Marel K, Wang K, Otte WM, el Bouazati S, et al. (2010) Recovery of sensorimotor function after experimental stroke correlates with restoration of resting-satte interhemispheric functional connectivity. J Neurosci 30: 3964-3972.