

Pushing HSI to the Limit for the Quantification of Peanut Traces in Bulk Powder Foods

Pilar Barreiro^{1*}, Belén Diezma¹, Teresa R. Cuadrado Dominguez¹, Inés María López-Calleja², Lourdes Lleó¹, Ana Herrero-Langreo³, Puneet Mishra¹, Satyabrata Ghosh¹, Eva Cristina Correa¹, Teresa García², Nathalie Gorretta³, SitiNurHidayah Mohamad¹, Pablo Delgado Sánchez⁵, Facundo Ruiz⁶, Christophe B.Y. Cordell⁶, Douglas N. Rutled⁶, Jean Michel Roger³, Rosario Martínde Santos² and Margarita Ruiz-Altisent¹

¹LPF_TAGRALIA Departamento de Ingeniería Agroforestal, UPM CEI Moncloa, Madrid, España

²Departamento de Nutrición, Bromatología y Tecnología de los Alimentos, Facultad de Veterinaria, Universidad Complutense de Madrid, 28040 Madrid, Spain

³Irstea, UMR ITAP, 361 Rue J. F. Breton, 34196 Montpellier Cedex 5, France

⁴AgroParisTech UMR1145 Genial, Analytical Chemistry Laboratory, 16 rue Claude Bernard, F-75005 Paris, France 5 Facultad de Agronomía y Veterinaria, UASLP, San Luis Potosí, México

⁵Facultad de Agronomía y Veterinaria, UASLP, San Luis Potosí, México

⁶Facultad de Ciencias, UASLP, San Luis Potosí, México

Received March 1, 2018; Accepted March 26, 2018; Published June 26, 2018

REVIEW

CEI Moncloa as the Origin of this Interdisciplinary Research

The CEI_Moncloa was a proposal coordinated by the Complutense University of Madrid (UCM) and the Technical University of Madrid (UPM), approved in 2011, in which 13 institutions have been integrated. It houses around 10,000 researchers, 10% of the national scientific production and approximately 80,000 students (75% of the

UCM and 25% of the UPM), including the Agri-Food and Health Cluster. It is situated in the so-called Agroalimentary Corridor, which groups together the activities that all these groups carry out in the production of agricultural and livestock products, and their subsequent processing for the creation of safe, healthy and nutritious food and feed. In this context we have been committed to finding synergistic and stable relations among research groups and as a consequence of a think tank set up to do this, there has been a steady scientific production since the first published works in 2014 (Figure 1).

Corresponding Author: Pilar Barreiro, LPF_TAGRALIA Departamento de Ingeniería Agroforestal, UPM CEI Moncloa, Madrid, España, Email: pilar.barreiro@upm.es

Citation: Barreiro P, Diezma B, Dominguez TRC, Calleja IML, Lleó L, et al., (2018) Pushing HSI to the Limit for the Quantification of Peanut Traces in Bulk Powder Foods. Food Nutr Current Res, 1(2): 56-60.

Copyright: ©2018 Barreiro P, Diezma B, Dominguez TRC, Calleja IML, Lleó L, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. Time line of the research at CEI_Moncloa(UCM_UPM), Agri-Food and Health Cluster.

Detection Limit of the Reference Measurement

The main outcome of the think tank was the realisation that there was a need (both in biomedicine and the Agrofood industry) for low-cost real-time non-destructive methods for detecting peanut traces in powdered food ingredients such as flour, milk and cocoa. According to IUPAC¹, "trace" means any element having an average concentration of less than about 100 parts per million atoms (ppm) or less than 100 ug g⁻¹. One of the relevant issues for the development of a new procedure is the need of a reliable reference procedure which should be at least one order of magnitude more sensitive than the proposed one. In this research the reference procedure was developed (prior to CEI_Moncloa) by the UCM research group (TRADETBIO) and consisted of both PCR and real-time PCR [1,2] for the quantification of peanut traces with 0.1 ppm resolution.

On the other hand, the spatial resolution of the Hyper Spectral images (HSI) images proposed by UPM (LPF_Tagrafia) was identified as the cornerstone for the detection of peanut traces, therefore at the very initial steps the optimal configuration of HIS was set as to characterize particles of 70um size.

Selection of Wavelength Range

VIS and NIR spectroscopy and HIS have been evaluated in consecutive trials [3-9] using IRMMⁱ-481e originating from Jumbo Runners, USA to guarantee the highest variability

¹ IUPAC, International Union of Pure and Applied Chemistry
IRMM International Reference Material
<https://ec.europa.eu/jrc/en/reference-materials>

among peanut origins and treatments. The main conclusion being that VIS is sensitive to peanut variability, and does not provide sufficient specificity regarding milk and flour, based on results validated with commercial peanut samples (Figure 2). On the other hand, NIR spectroscopic imaging provides characteristic bands for peanut detection: 950 nm arising from O-H bond, 1212 nm arising from 2nd overtone of the C-H stretch of CH₂group and 1450 nm arising from the 1st overtone of the O-H stretch, while HIS measurements at 70 µm resolution has shown the feasibility to detect up to 100 ppm (0.01 %) of peanut traces in any of the proposed powder foods (Figure 3).

SPECIFICITY ANALYSIS

Detecting peanut traces is relevant when it is sufficiently specific. Therefore, it is mandatory to avoid both false positives and false negatives. To this end, 49 commercial samples of cereals, legumes, oilseeds and nuts were obtained from local market of Madrid, Spain. All samples were ground with precautions to avoid cross-contamination. After grinding, all powdered food samples were sifted by passing through a sieve of 212 µm, since diffuse reflectance properties are dependent on the size of particles, non-uniformity in size can cause scatter effects in the spectra. In this case the wavelength range proposed included all NIR data, except for the bands with highest sensitivity to peanuts. The result (see procedure in Figure 5a), attained for calibration and external validation [10,11] showed that the greatest challenge was in isolating peanuts from pine nuts. Figure 4b provides an indication or the most relevant wavelength bands for specificity.




Figure 2. Identification of peanuts (IRMM 481e) and bulk powdered food ingredients (flour, and milk) with VIS-HIS.




Figure 3. Process in the identification of peanut traces by NIR HSI.




Figure 4. Segregation of bulk food materials from nuts (left), and wavelength used for specificity analysis (right)

Calibration Transfer, Transcontinental Approach

Finally, it was necessary to determine whether our results were reproducible irrespective of the instrumental platform, a hypothesis which was dismissed after initial trials at the transcontinental level [12]. Therefore the development of

calibration transfer techniques is required, and this step is still ongoing. **Figure 5b** summarizes some pros and cons of NIR spectroscopic techniques for the specific identification and quantification of peanut traces in powdered foods.




Figure 5. Summary of Chemometric procedure for the specific identification of peanut traces (left), and the pros and cons of NIR Spectroscopy for the identification of peanut traces in bulk food materials.

Prospective of Use for the Agro-Food Industry

The main conclusion of nearly a decade of research is that low-cost non-destructive NIR multispectral equipment associated with corresponding chemometric approaches should be commercially available in the near future, as derived from our AECOC² award in 2016. This technology can be complemented with real-time PCR when required with a highly significant reduction in cost, since it will only be used for contamination levels below 100ppm (0.01 %).

ACKNOWLEDGEMENTS

This research was financially supported by IDEAS <http://www.idt.mdh.se/ideas/> and INDIA4EU2 project of Erasmus Mundus <http://www.india4eu.eu/> and PICATA of the Moncloa Campus of International Excellence <http://www.campusmoncloa.es/en/calls/picata.php> (UCM-UPM, 2012).

REFERENCES

1. Maria Lopez-Calleja I, de la Cruz S, Gonzalez I, Garcia T, Martin R (2014) Survey of undeclared allergenic pistachio (*Pistacia vera*) in commercial

foods by hydrolysis probe real-time PCR. Food Control 39: 49-55.

2. Maria Lopez-Calleja I, de la Cruz S, Pegels N, Gonzalez I, Garcia T, et al. (2013) High resolution TaqMan real-time PCR approach to detect hazelnut DNA encoding for ITS rDNA in foods. Food Chem 141: 1872-1880.
3. Mishra P, Barreiro Elorza P, Roger JM, Diezma Iglesias B, Herrero Langreo A, et al. (2014) Detection of peanut traces in wheat flour through NIR hyperspectral imaging spectroscopy. Hyperspectral imaging conference. Ricoh arena, Coventry, United Kingdom.
4. Mishra P, Roger J-M, Barreiro Elorza P, Diezma Iglesias B, Herrero Langreo A, Lleó García L, et al. (2014) Application of Improved Direct Calibration for Hyperspectral image processing: Detecting peanut traces in wheat flour. Food Analysis Congress, Barcelona, Spain.
5. Mishra P, Herrero-Langreo A, Barreiro P, Roger JM, Diezma B, et al. (2015) Detection and quantification of peanut traces in wheat flour by near infrared hyperspectral imaging spectroscopy using principal-component analysis. J Infrared Spectroscopy 23: 15-22.

² AECOC Asociación Española de Codificación Comercial

6. [Mishra P, Herrero Langreo A, Barreiro Elorza P, Roger J-M, Diezma Iglesias B, et al. (2015) Hyperspectral to multispectral imaging for detection of tree nuts and peanut traces in wheat flour. *J Spectral Imaging* 4: 1-9.
7. Mohamad SNH, Ghosh S, Mishra P, Barreiro Elorza P, Diezma Iglesias B. (2015) Star dust memories, or how to detect allergen traces in powder foods. VII Congreso de Estudiantes Universitarios de Ciencia, Tecnología e Ingeniería Agronómica. Madrid, Spain.
8. Ghosh S, Domínguez TRC, Diezma B, Lleó L, Barreiro P, et al. (2015) VIS/NIR spectral signature for the identification of peanut contamination of powder foods. *Agric Eng Int J* 17: 310-329.
9. Cuadrado Dominguez TR, Ghosh S, Barreiro P, Lleó García L, Diezma B, et al. (2014) VIS-NIR spectroscopy for the detection of peanuts traces in powder food. In: Tellez FA, Rodriguez AM, Sancho IM, Robinson MV, RuizAltisent M, Ballesteros FR, Hernando ECC, eds. VII Congreso Ibérico de Agroingeniería y Ciencias Hortícolas: Innovar y Producir para el Futuro Innovating and Producing for the Future. Madrid, Spain.
10. Mishra P, Cordella CBY, Rutledge DN, Barreiro P, Roger JM, et al. (2016) Application of independent components analysis with the JADE algorithm and NIR hyperspectral imaging for revealing food adulteration. *J Food Eng* 168: 7-15.
11. Ghosh S, Mishra P, Mohamad SNH, Martin de Santos R, Diezma Iglesias B, et al. (2016) Discrimination of peanuts from bulk cereals and nuts by near infrared reflectance spectroscopy. *Biosystems Eng* 151: 178-186.
12. Cuadrado Dominguez TR, Barreiro Elorza P, Lleó García L, Ruiz F, et al. (2016) Prospective of NIR Calibration transfer need for detecting peanut traces in powder foods: a transcontinental approach. 4th CIGR International Conference of Agricultural Engineering (CIGR-AgEng2016). Aarhus University, Denmark.