

Micronucleus Assay in Different Grades of Oral Squamous Cell Carcinoma-A Comparative Study

Dave Gunjan^{1*} and Parikh Neelampari²

^{*1} MDS Oral Pathology, PhD research fellow, Gujarat University, Ahmedabad, Gujarat, India

² Department of Oral and Maxillofacial Pathology, Faculty of Dental Sciences, Dharamsinh Desai University, Nadiad, Gujarat, India.

Received April 05, 2023; Revised May 01, 2023; Accepted May 04, 2023

ABSTRACT

Introduction: Oral squamous cell carcinoma (OSCC) is a multistage process which on exposure of carcinogens (smoking and smokeless associated) cause complex karyotypes like chromosomal deletions, translocations, and structural abnormalities. Detection of these genetic changes resulting from errors in DNA metabolism, in buccal mucosal cells can be done by using biomarkers that help to identify individuals vulnerable to malignancy. Micronucleus (MN) is one such standard biomarker in the field of oral cancer. It is evident that the individual cancer risk can be predicted on the basis of increase in number of micronuclei in the oral epithelial cells. Hence this study was done to compare the level of micronucleus & micronucleated cells (MNC) in normal subjects and OSCC subjects. Also, MN & MNC were compared in different grades of OSCC.

Material and Method: The study consisted of 2 groups, Group A Thirty healthy controls and Group B, thirty histopathologically proven cases of OSCC. The exfoliated cells were collected from buccal mucosa/lesional area of both study groups and were fixed, and then stained with Feulgen stain and evaluated under microscope for MN and MNC. Statistical analysis was done using ANOVA test and correlation was calculated using Pearson's correlation using SPSS 18.0 software.

Results and Observations: Statistically highly significant difference of 0.0001 was obtained between group A and group B. Mean of MN and MNC and MN% increased more than 3 folds in OSCC cases compared to controls. Also, all three parameters increased from well to moderate to poorly differentiated carcinoma but statistically significant difference was obtained only between well and poorly differentiated carcinoma.

Conclusion: Micronucleus assay can be used as a standard to identify genotoxic damage occurring due to carcinogens and it increases in OSCC cases as compared to controls in buccal mucosa cells.

Keywords: Micronucleus, Oral squamous cell carcinoma, Feulgen stain

Abbreviations: ANOVA: Analysis of Variance; DNA: Deoxyribonucleic Acid; DPX: Di-n-butyl Phthalate in Xylene, H and E, H & E: Hematoxylin and Eosin Stain; ICMR: Indian Council of Medical Research; MGG: May-Grunwald-Giemsa; MN%: Micronucleus Percentage; MN: Micronucleus; MNC: Micronucleated Cell; Mini: Micronuclei; NS: Not Significant; OSCC: Oral Squamous Cell Carcinoma; PAP: Papanicolaou; S: Significant; SCC: Squamous Cell Carcinoma; WHO: World Health Organization

INTRODUCTION

The National Cancer Registry Programmed report of The Indian Council of Medical Research (ICMR) estimated the number of cancer cases in India to be 13.9 lakhs in 2020. These numbers are likely to increase to 15.7 lakhs by 2025 [1]. Another set of figures say that around 77,000 new cases and 52,000 deaths are reported annually in India which are related to oral cancer. This makes India a world leader of oral cancer cases, which is about 1/3rd of total oral cancer cases globally [2]. This presents a real frightening picture but the good news is that these cancers can be

Corresponding author: Dave Gunjan, PhD Research fellow, Gujarat University, Ahmedabad, Gujarat, India, Tel: +919725944249; E-mail: gunjan_dave29@rediffmail.com

Citation: Gunjan D & Neelampari P. (2023) Micronucleus Assay in Different Grades of Oral Squamous Cell Carcinoma-A Comparative Study. J Oral Health Dent, 6(3): 520-527.

Copyright: ©2023 Gunjan D & Neelampari P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

prevented by screening or by early detection and treated in early stage [3]. Squamous cell carcinoma (SCC) accounts for 84-97% of all oral cancers [2].

Oral squamous cell carcinoma (OSCC) is a multistage process which on exposure of carcinogens (smoking and smokeless associated) cause complex karyotypes that involve many chromosomal deletions, translocations, and structural abnormalities [4,5]. Detection of these genetic changes resulting from errors in DNA metabolism, in buccal mucosal cells can be done by using biomarkers that help in identifying individuals vulnerable to malignant transformation [6]. Micronucleus is one such standard biomarker in the field of oral cancer [7]. Biologically, micronuclei (Mini) are the chromosome fragments or whole chromosomes that lag behind at anaphase during nuclear division [8].

After telophase, the undamaged chromosomes, as well as the centric fragments, give rise to regular daughter nuclei. The lagging elements are included in the daughter cells, but a considerable proportion is transformed into one or several secondar nuclei, which are, as a rule, much smaller than the principal nucleus and are therefore called micronuclei. Bigger micronuclei result from exclusion of whole chromosome following damage to the spindle apparatus of the cell (A eugenic effect), whereas smaller micronuclei result from structural aberrations; causing chromosomal fragments (Clastogenic effect) [9].

Besides these two important mechanisms, MN may be formed due to broken anaphase bridges. This may be

because of dicentric chromatids, intermingled ring chromosomes, union of sister chromatids or from extrachromosomal elements called double minutes [10].

The assessment of micronuclei in exfoliated cells is a promising tool for the study of epithelial carcinogens [11]. Screening of individuals who are at high risk of malignant transformation is more pivotal in preventing and reducing the number of deaths. It is evident that the individual cancer risk can be predicted on the basis of increase in number of micronuclei in the oral epithelial cells [5]. Hence this study was done to compare the level of micronucleus & micronucleated cells in normal subjects and OSCC subjects. Also, number of micronucleus and micronucleated cells were compared in different grades of oral squamous cell carcinoma.

METHOD AND MATERIAL

The study was conducted in the Department of Oral and Maxillofacial Pathology and Department of Oral Medicine and Radiology of Karnavati School of Dentistry, Uvaras, Gandhinagar, the data collection consisted of randomly selected sixty cases and divided into following groups.

Group A: Thirty healthy controls with no oral lesions, no previous tobacco-related habits, and no previous medical history with age and sex matching with diseased cases
Group B: Thirty histopathologically proven cases of oral squamous cell carcinoma. **Table 1** shows gender wise distribution of cases.

Table 1. Gender Wise Distribution of Cases.

Group	Total No. of Patients	Males	Females
Group A [Control cases]	30 (100%)	26 (86.66%)	4 (13.34%)
Group B [OSCC cases]	30 (100%)	25 (83.33%)	5 (16.67%)
Total	60 (100%)	51 (85%)	9 (15%)

In each group cases were given a standard questionnaire interview and their history was recorded. A written consent was taken from cases of both Groups. Routine hematological investigations were carried out in study cases prior to biopsies.

Collection of Exfoliated cells: The participants of study were asked to rinse their mouth with water and a pre-moistened wooden spatula was used to sample cells from lesional area of Group B (Oral Squamous Cell Carcinoma) cases and from the buccal mucosa of Group A (Control) cases. The cells were immediately smeared on pre-cleaned microscopic slides.

The smears were fixed with fixative solution for minimum fifteen minutes (80% methyl alcohol + glacialacetic acid fixative mixed in ratio 3:1). The smears were stained with

Feulgen stain.

Biopsy procedure: Incisional/punch biopsy in cases with oral lesions was taken for histopathologic examination. The specimens were placed in 10% neutral buffered formalin for fixation for minimum 24 hours. The biopsies were processed in automated tissue processor, embedded in paraffin wax blocks. 4-5µm thick sections were prepared, stained with routine H and E stain, and mounted using DPX mount ant. Then they were analyzed for confirmed histopathological diagnosis using light microscope. Only histopathologically proven cases of OSCC were considered in the study.

H and E-stained sections of Group B cases (Oral Squamous Cell Carcinoma cases) were evaluated and graded using Broder's classification.

Broder's classification (1920): In each group cases were

given a standard questionnaire interview and their history was recorded. A written consent was taken from cases of both Groups. Routine hematological investigations were carried out in study cases prior to biopsies.

Collection of Exfoliated cells: The participants of study were asked to rinse their mouth with water and a pre-moistened wooden spatula was used to sample cells from lesional area of Group B (Oral Squamous Cell Carcinoma) cases and from the buccal mucosa of Group A (Control) cases. The cells were immediately smeared on pre-cleaned microscopic slides.

The smears were fixed with fixative solution for minimum fifteen minutes (80% methyl alcohol + glacialacetic acid fixative mixed in ratio 3:1). The smears were stained with Feulgen stain.

Biopsy procedure: Incisional/punch biopsy in cases with oral lesions was taken for histopathologic examination. The specimens were placed in 10% neutral buffered formalin for fixation for minimum 24 h.

The biopsies were processed in automated tissue processor, embedded in paraffin wax blocks. 4-5µm thick sections were prepared, stained with routine H and E stain, and mounted using DPX mountant. Then they were analyzed for confirmed histopathological diagnosis using light microscope.

Only histopathologically proven cases of OSCC were considered in the study.

H and E-stained sections of Group B cases (Oral Squamous Cell Carcinoma cases) were evaluated and graded using Broder's classification.

Broder's classification (1920) Cases of Oral squamous cell carcinoma was graded on the basis of degree of differentiation and keratinization of tumor cells into:

- Grade I: Well differentiated tumors - 75-100% of cells are differentiated
- Grade II: Moderately differentiated tumors - 50-75% of cells are differentiated
- Grade III: Poorly differentiated tumors - 25-50% of cells are differentiated.
- Grade IV: Anaplastic tumor - 0-25% of cells are differentiated [12].

Evaluation of micronucleus in cytology:

All the slides of cytology smears were stained with Feulgen stain and observed under low magnification (10X) for

screening and high magnification (40X) for counting of MN.

The zigzag method was used for screening slides and 500 cells with intact nuclei and cell boundaries were counted per patient.

Scoring criteria:

The scoring of micronucleus was done according to criteria given by Tolbert [13]

1. Rounded smooth perimeter.
2. Less than one third diameter of the associated nucleus, but large enough to discern shape and color.
3. Staining intensity similar to that of nucleus
4. Texture similar to that of nucleus
5. Same focal plane as nucleus
6. Absence of overlap, with or bridge to, the nucleus

The exclusion criteria considered when scoring of micronuclei was done were broken eggs, nuclear budding, binucleated cell, karyomeric, karyolytic [7].

Observations were expressed in form of No. of micronucleus (MN) in each patient, No. of micronucleated cells (MNC) per patient and Micronucleus percentage (MN%).

Micronucleus percentage was counted using this formula: $100 \times \frac{\text{total number of micronuclei found in a Group}}{\text{No. of cells counted per patient (500)} \times \text{total number of patients in a Group}}$ Statistical Analysis: - The data obtained was statistically analyzed using ANOVA test and correlation was calculated using Pearson's correlation using SPSS 18.0 software.

RESULTS AND OBSERVATIONS

The study comprised of 60 subjects 30 controls and 30 cases of OSCC. As shown in **Table 1** were 26 males and 4 females in control group and 25 males and 5 females in OSCC case group. **Table 2** shows mean of micronucleus, micronucleated cell & MN % in group A and group B, 0.66% and 2.46% respectively. On comparison using ANOVA test highly significant difference of 0.0001 is obtained between group A and group B (**Table 3**). **Table 4** shows age distribution of cases in the study with interval of 10 years and no of cases in each group. Micronucleus and MNC found in each corresponding age group is shown. In present study highest no. of cases and highest no of MN & MNC were found in 36-45 age group.

Table 2. Comparison of Mean of MN and MNC in Group A (Control Cases), GroupB (OSCC cases).

Group	Mean of MN $\pm$ Standard deviation	Mean of MNC $\pm$ Standard deviation	MN%
Group A (Control Cases)	3.30 $\pm$ 1.022	3.30 $\pm$ 1.022	0.66%
Group B (OSCCcases)	10.10 $\pm$ 1.668	9.77 $\pm$ 1.524	2.15%

N=60

Table 3. Frequency of total number of MN and total number of MNC according to age distribution in Group A (Control cases) and Group B (OSCC cases).

Age (in years)	Group A (Control Cases)			Group B (OSCC cases)		
	No. of cases (30)	MN (No.)	MNC(No.)	No. of cases (30)	MN (No.)	MNC(No.)
16-25	1	3	3	1	9	9
26-35	5	16	16	8	78	77
36-45	12	38	38	11	118	113
46-55	10	32	32	5	47	46
56-65	1	3	3	4	40	39
66-75	1	5	5	1	12	12
Total	30	97	97	30	304	296

Table 4. Comparison of Occurrence of MN between Group A (Control cases), and Group B (OSCC cases) (ANOVA test).

Groups	p - value	Significance
Group A (Control cases) Vs Group B (OSCC cases)	<0.0001	Highly Significant

N=60

Amongst 30 cases of OSCC 17 were of Well differentiated OSCC, 10 were of Moderately differentiated OSCC and 3 were Poorly differentiated OSCC. The mean of MN and MNC increased from well to moderate to poorly

differentiated OSCC as shown in **Table 4** and **Graph 1** but statistically significant difference was only seen between well and poorly differentiated OSCC (**Tables 5 & 6**).

Table 5. Mean of MN & MNC in different grades of OSCC (Group B cases) (Well Differentiated, Moderately Differentiated and Poorly Differentiated) (Broder's Classification).


Grades of OSCC	No. of Cases (out of 30) (%)	Mean of MN $\pm$ Standard Deviation	Mean of MNC $\pm$ Standard Deviation
Well differentiated OSCC	17 (56.67%)	9.35 $\pm$ 1.320	9.18 $\pm$ 1.185
Moderately differentiated OSCC	10 (33.33%)	10.70 $\pm$ 1.494	10.30 $\pm$ 1.418
Poorly differentiated OSCC	3 (10%)	12.33 $\pm$ 1.528	11.33 $\pm$ 2.309
Total	30 (100%)	10.10 $\pm$ 1.668	9.77 $\pm$ 1.524

**No case of grade IV OSCC was found in present study.*


Table 6. Comparison of occurrence of MN between different grades of OSCC(Group B cases) (ANOVA test).

Grades of Cancer	p-value	Significance
Well differentiated OSCC vs Poorly differentiated OSCC	0.006	S
Well differentiated OSCC vs Moderately differentiated OSCC	0.057	NS
Moderately differentiated OSCC vs Poorly differentiated OSCC	0.196	NS

S = significant, NS = Not significant

**Graph 1.** Mean of MN and MNC in different Grades of OSCC (Group B cases).

MN% was calculated and it increased from 1.87% to 2.14% respectively (Graph 2). from well to moderate to poorly differentiated carcinoma

**Graph 2.** MN% in in different Grades of OSCC (Group B cases).

DISCUSSION

William Howell, an American, and Justin Jolly, a Frenchman were first to identify Micronuclei (MN), also known as Howell-Jolly bodies, in red cell precursors. Initially, Howell-Jolly bodies were described as remnants of nuclei of red blood cells circulating in organs with pathology. So later on, the micronucleus technique was proposed as a reliable method for measuring chromosomal damages caused by cytotoxic agents “*it vivo*” by Fenech

and Morley in 1985 [14]. Burkhardt [15] suggested that the MN test applied to exfoliated cells could provide valuable information and improve the evaluation of carcinomas by exfoliative cytology [15].

It has been documented by Gupta [16] that the habit of chewing gutkha had gained considerable popularity among the younger men in India [16]. The rapidly increasing prevalence of this habit can be judged from the reports that the Indian market for pan masala and gutkha is worth

43,410.2 crores in 2022 [17].

The carcinogenic agents in tobacco act by inducing changes at both genetic level and locally by providing a conducive local environment for hyperplastic transformation of the buccal cells [18]. Also, micronuclei in exfoliated cells emerge during mitosis of the basal layers of the epithelium and their absolute quantities could reflect the real situation in target cells [6]. In a study, levels in exfoliated buccal mucosa cells of patients with breast, lung, cervix uteri cancer, and patients with Hodgkin's disease were studied and levels of MN were found to be elevated in buccal mucosal cells. This reflected level of instability in somatic cell due to cancerous lesions even when there is cancer in distant body organs [19]. The frequency of micronucleated cells do reflect the capacity of target tissues to activate procarcinogens into reactive species or to inactivate or trap ultimate carcinogens [20].

Hence, this study was performed with primary aim of evaluating frequency of micronuclei in exfoliated cells of buccal mucosa from healthy control subjects and squamous cell carcinoma patients. Also, micronucleus & micronucleated cell frequencies in different grades of oral squamous cell carcinoma were compared.

The oral squamous cell carcinoma cases in our study comprised of 83.33% males and 16.67% females. These figures were found to be similar to the studies conducted by Dindigire [21] who had 85% males in malignant group and Casartelli [22] who had 85.71 % of males in their respective studies [22]. This gender distribution show that habits being prevalent more in males and that occurrence of cancerous lesions are seen more in them as compared to controls.

In present study maximum number of cancer cases was recorded in age range of 26-45 i.e., 2nd to 4th decade (as shown in **Table 3**). But contrary to our study the age range of cancer subjects in other studies [5,21-24] were seen more after 5th decade. The reason for prevalence of cancer in early decade of life in our study could be beginning of habits at early age during middle school and high school period and hence higher incidence of cancerous lesions in early half of life span [25].

In their study Grover [26] concluded that because of the possibility of mis-interpretation of nuclear anomalies (e.g., karyorrhexis, karyolytic, condensed chromatin binucleates and keratin granules) as MNi with nonspecific stains, a higher count was observed with PAP and H and E stain [26]. Contrary to this Kamboj [20] in their study concluded that fluorescent stain was more sensitive than Feulgen stain. Jyoti [27] also concluded that chances of errors during scoring MN are less when cells are stained using fluorescent stain such as acridine orange. Fluorescent staining technique involves use of acridine orange stain which is costly and also involve use of fluorescent microscope which is more technique sensitive and adds up to the cost.

So, in our study we used Feulgen staining technique and binocular microscope for analysis of micronuclei which was simple and economical. Feulgen staining technique is also known for high DNA specificity and a clear transparent appearance of the cytoplasm which enables easy identification of MN [26]. Also, it is most common stain used for scoring of MN.

In present study, MN count per 500 cells in controls was 97 and that in cancerous lesion group was 304. This shows that MN count increases more than 3 times in cancer cases as compared to that in controls.

Dindigire [21] in their study inferred that MN count per 500 cells was 28 (20 subjects) in tobacco users without any lesion and that in malignant conditions was 166 among 20 subjects. This study used PAP and MGG (May Grunwald Giemsa) stain [21]. Also, in their study to determine MN index as early diagnostic criteria of carcinoma Sivasankari PN et al (2008) found total 77 MN in control and total 145 MN in Malignant cases using MGG stain [24].

All these studies, including ours, have shown that there is almost double to triple fold increase in MN and MNC count as severity of disease increases.

Mean of MN and MN% increase from control to OSCC cases in present study. In relation to mean of MN, in study done by Casartelli [22] showed increase from 1.5 in controls to 4.4 in cancerous conditions. Similar were findings of other studies [20,21,24]. In present study mean of MN increased from 3.30 in controls to 10.10 in cancer cases. When viewed in relation to MN % apart from other studies (5,23,28) our study showed increase in MN% from control to OSCC cases. In our study MN% increased from 0.66% in controls to 2.15% in OSCC cases. Our study inferred that micronucleus frequencies and frequency of micronucleated cells increase from controls to cancerous lesions subjects. Also, the occurrence MN frequencies in controls and cancerous conditions showed statistically significant difference in our study ($p < 0.0001$) and these findings are similar to above mentioned studies.

These variations in micronucleus, micronucleated cells and MN% may be attributed to the ingredients in the quid, the number of quids per day and to the different lifestyles and food habits of people where study is conducted [20]. Also, variability may be due to scoring criteria and staining procedures used [7].

The levels of MN% in our study in SCC groups (from well differentiated OSCC to moderate differentiated OSCC to poorly differentiated OSCC) was between 1.87% to 2.46% whereas that in study done by Palve [9] was between 1.1% to 3% and that between Kumar V et al (2000) was between 1.4% - 9.15 %. The reason for this difference was different technique used for assessing exfoliated cells. Palve D et al (2008) used rapid papanicolaou technique whereas Kumar V et al (2000) used fluorescent dyes for staining purpose

whereas present study utilized feulgen staining technique. Also, Kumar V et al (2000) in their study obtained cells by mincing biopsy tissue and prepared cell suspensions from it, whereas in our study simple non-expensive method of exfoliative cytology was used. As different techniques are used to obtain samples and different staining techniques used there are variations in number of micronucleated cells and micronucleus counts in various studies [9].

The present study, studies of Palve [9], Saran [28] showed increase in micronucleus from low grade carcinoma to poor grade carcinoma. The reason for this increase in micronucleus count with increase in grades of carcinoma can be attributed to increased DNA damage in relation to tumor grade as there is gradual progression of carcinogenic process [28].

Hence this study shows that with increase in severity of disease the MN frequencies increase and also that with increase in MN% individual cancer risk increases. Present study has used border's classification for histopathological grading and Feulgen stain for identifying micronuclei. Future studies are suggested to be with larger sample size and performed longitudinally to evaluate the lesions and also addressing sources of variability like strict adoption of optimal scoring criteria using standard histopathological grading system and other DNA specific stains in the assay to confirm the findings. Comparison of MN with other biomarkers and use of MN assay as a biomarker of genotoxicity in predicting the effects of cancer intervention studies may also be considered.

CONCLUSION

- 1) Occurrence of micronucleus and micronucleated cell increases from Controls to OSCC cases and that there is statistically significant difference (p value < 0.001) for occurrence of MN between controls and OSCC cases ($p < 0.0001$).
- 2) Amongst three stages of cancer MN shows statistically significant difference between well and poorly differentiated carcinoma.

Micronucleus assay is a simple and rapid screening test applied for an early detection of cancer. Thus, this test can be used on exfoliated cells to identify the genotoxic damage in human tissues, which are targets for carcinogens and from which carcinomas develop.

Conflict of interest: None

ACKNOWLEDGMENT

Staff of Department of Oral Pathology and Department of Oral Medicine, Karnavati School of Dentistry, Uvarsad, Gandhinagar. Dr. Girish Mishra (MS ENT, Prof. and Guide, Department of Otolaryngology and Head and Neck surgery, Pramukhswami Medical College, Karamsad. Dr. Daxa

Mishra MPT, Ph.D., K.M. Patel Institute of Physiotherapy, Karamsad.

REFERENCES

1. Sharma LK (2020) ICMR-NCDIR National Cancer Registry Programme estimates 12% increase in cancer cases in the country by 2025: New Delhi. Indian Council of Medical Research. Accessed on: February 10, 2022. Available online at: https://ncdirindia.org/AllReports/Report_2020/PB/Press_release.pdf
2. Borse V, Konwar AN, Buragohain P (2020) Oral cancer diagnosis and perspectives in India. *Sens Int* 1: 100046.
3. Mathur P, Sathishkumar K, Chaturvedi M, Das P, Sudarshan KL, et al. (2020) Cancer Statistics, 2020: Report from National Cancer Registry Programme, India. *JCO Glob Oncol* 6: 1063-1075.
4. Priya B, Spadigam A, Dhupar A, Syed S (2020) Frequency of micronuclei in tobacco habitués and non-Habitués with oral lichen planus. *CHRISMED J Health Res* 7(3): 151.
5. Devi P, Vb T, Mehrotra V, Arora P. (2011) Micronucleus Assay for Evaluation of Genotoxicity in Potentially Malignant and Malignant Disorders. Kailasam S, editor. *J Indian Acad Oral Med Radiol* 23: 97-100.
6. Mahimkar MB, Samant TA, Kannan S, Patil T (2010) Influence of genetic polymorphisms on frequency of micronucleated buccal epithelial cells in leukoplakia patients. *Oral Oncol* 46(10): 761-766.
7. Jadhav K, Gupta N, Ahmed Mujib B (2011) Micronuclei: An essential biomarker in oral exfoliated cells for grading of oral squamous cell carcinoma. *J Cytol* 28 (1): 7.
8. Samanta S, Dey P (2012) Micronucleus and its applications. Baloch Z, editor. *Diagn Cytopathol* 40(1): 84-90.
9. Palve D, Tupkari J (2008) Clinico-pathological correlation of micronuclei in oral squamous cell carcinoma by exfoliative cytology. *J Oral Maxillofac Pathol* 12(1): 2.
10. Shimizu N (2011) Molecular mechanisms of the origin of micronuclei from extrachromosomal elements. *Mutagenesis* 26(1): 119-123.
11. Jois HS, Kale AD. (2010) Micronucleus as Potential Biomarker of Oral Carcinogenesis. *Indian J Dent Adv* 2(2):197-202.
12. Anneroth G, Hansen LS (1984) A methodologic study of histologic classification and grading of malignancy in oral squamous cell carcinoma. *Eur J Oral Sci* 92(5):

- 448-468.
13. Tolbert PE, Shy CM, Allen JW (1992) Micronuclei and other nuclear anomalies in buccal smears: methods development. *Mutat Res Mutagen Relat Subj* 271(1): 69-77.
 14. Luzhna L, Kathiria P, Kovalchuk O (2013) Micronuclei in genotoxicity assessment: from genetics to epigenetics and beyond. *Front Genet* [Internet]. 4. Available online at: <http://journal.frontiersin.org/article/10.3389/fgene.2013.00131/abstract>
 15. Burkhardt A (1985) Advanced methods in the evaluation of premalignant lesions and carcinomas of the oral mucosa. *J Oral Pathol Med* 10: 751-778.
 16. Gupta MK, Mhaske S, Ragavendra R (2008) Oral submucous fibrosis - Current Concepts in Etiopathogenesis. *People J Sci Res* 1: 39-44.
 17. India Pan Masala Market Size, Share and Trends 2023-2028 [Internet]. [cited 2023 Mar 30]. Available from: <https://www.imarcgroup.com/prefeasibility-report-pan-masala-processing-plant>
 18. Khan Z, Khan S, Christianson L, Rehman S, Ekwunife O, et al. (2016) Smokeless tobacco and oral potentially malignant disorders in South Asia: A protocol for a systematic review. *Syst Rev* 5(1): 142.
 19. Nersesyan A, Vardazaryan N, Gevorgyan A, Arutyunyan R (2022) Micronucleus level in exfoliated buccal mucosa cells of cancer patients. *Arch Oncol* 10(1): 35-36.
 20. Kamboj M, Mahajan S (2007) Micronucleus-an upcoming marker of genotoxic damage. *Clin Oral Investing* 2: 121-126.
 21. Dindigre SL, Gosavi S, Kumawat RM, Ganvir S, Hazarey V (2012) Comparative Study of Exfoliated Oral Mucosal Cell Micronucleus Frequency in Potentially Malignant and Malignant Lesions. *Int J Oral Maxillofac Pathol* 3(2): 15-20.
 22. Casartelli G, Bonatti S, De Ferrari M, Scala M, Mereu P, et al. (2000) Micronucleus frequencies in exfoliated buccal cells in normal mucosa, precancerous lesions and squamous cell carcinoma. *Anal Quant Cytol Histol* 6: 486-492.
 23. Halder A, Chakraborty T, Mandal K, Gure PK, Das S, et al. (2004) Comparative Study of Exfoliated Oral Mucosal Cell Micronuclei Frequency in Normal, Precancerous and Malignant Epithelium. *Int J Hum Genet* 4: 257-260.
 24. Reddy KS (2008) Micronucleus Index: An Early Diagnosis in Oral Carcinoma. *J Anat Soc India* 57: 8-13.
 25. Chitroda PK, Shah TJ, Katti G, Ghali S (2011) A Correlative Study of Smokeless Tobacco-induced Lesion and Smoke-induced Leukoplakia in Various Aspects. Kailasam S, editor. *J Indian Acad Oral Med Radiol* 23(2): 86-91.
 26. Grover S, Mujib A, Jahagirdar A, Telagi N, Kulkarni P (2012) A comparative study for selectivity of micronuclei in oral exfoliated epithelial cells. *J Cytol* 4: 230.
 27. Jyoti S, Khan S, Afzal M, Naz F, Siddique YH (2013) Evaluation of micronucleus frequency by acridine orange fluorescent staining in buccal epithelial cells of oral submucous fibrosis (OSMF) patients. *Egypt J Med Hum Genet* 2:189-193.
 28. Saran R, Tiwari RK, Reddy PP, Ahuja YR (2008) Risk assessment of oral cancer in patients with pre-cancerous states of the oral cavity using micronucleus test and challenge assay. *Oral Oncol* 44(4): 354-360.