Journal of Infectious Diseases and Research

JIDR, 6(S4): 26 www.scitcentral.com



Abstract: Open Access

Investigating Antimicrobial Resistance Patterns in *Gardnerella Vaginalis* Isolates

Thasmika Durga^{2*} and Nathlee S Abbai¹

¹School of Laboratory Medicine & Medical Sciences, Department of Medical Microbiology, College of Health Sciences, University of KwaZulu-Natal, Durban, KwaZulu-Natal, South Africa

²School of Clinical Medicine Research Laboratory, College of Health Sciences, University of KwaZulu-Natal, Durban, KwaZulu-Natal, South Africa.

Published November 28, 2023

ABSTRACT

Background and Aim: Antimicrobial resistance has increased in *G. vaginalis*. The aim was to determine the susceptibility patterns of isolates of *G. vaginalis* to antibiotics and investigated the genetic diversity of the isolates as well as linked the genetic data to patterns of antimicrobial resistance.

Methods: A cross sectional study where a total of n=150 pregnant women were enrolled. Vaginal swabs were used for cultivation of *G. vaginalis* and diagnosis of BV via Nugent scoring. Genetic diversity assessments were based on the genetic differences in the *tuf* gene using clade specific primers on a qPCR platform. The antimicrobial susceptibility profiles were generated using the SensititreTM Anaerobe MIC Plate (Thermo Fischer Scientific, United States).

Results: 17 isolates of *G. vaginalis* was isolated out of the 150 vaginal swabs. 5 were BV positive, 2 were BV negative and 9 were BV intermediate. All isolates were oxidase and catalase negative. The *16S rRNA* gene was amplified in all isolates. 0% belonged to clade 3. The frequencies for clades were as follows; 100% for clade 1, 37.5% for clade 2 and 43.75% for clade 4. Multiple clades were found in 56.25% of the isolates. 15 isolates were successfully cultured and tested. 60% were susceptible to metronidazole and 40% were resistant to metronidazole. BV positive women (5), 20% showed metronidazole resistance. BV negative women (2), 50% showed metronidazole resistance was in women with BV intermediate status. All women that were resistant to metronidazole experienced abnormal discharge. 100% of isolates with metronidazole resistance fell into clade 1. 16.66% of metronidazole resistant isolates fell into clades 1 and 4.

Conclusion: This study has shown a link between clade 1, BV intermediate women and metronidazole resistance. More research needs to focus on genotyping to understand the resistance patterns.

Keywords: Antimicrobial resistance, G. vaginalis, Sensititre™ Anaerobe MIC Plate, Metronidazole resistance

Abbreviations: BV: *Bacterial vaginosis; G. vaginalis: Gardnerella Vaginalis;* MIC: Minimum Inhibitory Concentration; qPCR: Quantitative Polymerase Chain Reaction; RNA: Ribonucleic Acid

Citation: Durga T & Abbai NS. (2023) Investigating Antimicrobial Resistance Patterns in Gardnerella Vaginalis Isolates. J Infect Dis Res, 6(S4): 26.

Copyright: ©2023 Durga T & Abbai NS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Corresponding author: Thasmika Durga, School of Clinical Medicine Research Laboratory, College of Health Sciences, University of KwaZulu-Natal, 719 Umbilo Road, Durban, 4001, South Africa, E-mail: thasmikadurga2@gmail.com