

Anesthetic Concerns of Ebstein's Anomaly Patient Undergoing Lower Limb Orthopedic Surgery

Lahiri Swati*, Adrija Chakraborty, Moumi Dey, Mukulanda Panja, Somnath Dey and Satyajit Bera

*Department of Anesthesiology and Critical Care, Medical College, Kolkata, West Bengal, India.

Received June 24, 2023; Revised June 27, 2023; Accepted June 30, 2023

ABSTRACT

Ebstein's Anomaly is a rare congenital disorder (<1%) in which distorted and displaced tricuspid valve leaflets cause atrialization of the right ventricle leading to tricuspid regurgitation of varying severity. The main aim of anesthetic management is avoidance of tachyarrhythmias and maintenance of preload and afterload by careful monitoring, fluid management and prevention of hypotension. Here we are presenting perioperative anesthetic consideration of a patient with Ebstein's anomaly who manifested three episodes of tachyarrhythmias intraoperatively however no episode of hypotension was observed under general anesthesia.

Keywords: Anesthesia general, Ebstein's anomaly, Congenital, Distal sciatic nerve block

Abbreviations: RV: Right Ventricle; TV: Tricuspid Valve; RA: Right Atrium; LV: Left Ventricle; MV: Mitral Valve; LA: Left Atrium

INTRODUCTION

Patients with Ebstein's Anomaly have apical displacement of tricuspid valve which mainly affects septal and posterior leaflets, apical displacement of tricuspid annulus and anteroapical displacement of tricuspid orifice [1]. Atrialization of Right Ventricle (RV) splits RV into 2 portions: atrialized RV and functional RV. The atrialized RV becomes dilated along with the right atrium from tricuspid regurgitation. Functional RV becomes very small and depending on severity of leaflet displacement [2]. Frequently the anterior cusp is sail like in structure with multiple fenestrations resulting in tricuspid insufficiency. Patient with Ebstein anomaly can be either cyanotic or acyanotic. Secundum atrial septal defect or a patent foramen ovale is also very common in Ebstein's Anomaly. Due to compression of AV node by septal leaflet conduction system anomalies and accessory pathways can also be associated with Ebstein's Anomaly. About one third of patients have more than one accessory pathway [3]. 5-25% patients with Ebstein's Anomaly present with Wolff-Parkinson-White (WPW) syndrome. From congestive cardiac failure in childhood to incidental diagnosis in adulthood - clinical manifestation varies in a wide range.

CASE REPORT

Here we are presenting a case of a 28 years old man (Body mass index 23.89 Kg/m²) diagnosed with Ebstein's

Anomaly incidentally during pre-anesthetic checkup. He fell from a tree and got a fracture of both tibia and fibula (left leg) with eye injury scheduled for closed reduction and internal fixation with intramedullary interlocking tibia nail. He was completely asymptomatic since childhood. He had no complaints of exertional dyspnea, palpitations, tachyarrhythmias or chest pain. There were no signs of congestive cardiac failure on physical examination. On auscultation pan systolic murmur was heard at lower left sternal border suggestive of tricuspid regurgitation. His laboratory investigations were normal except short PR interval, wide QRS, delta wave found in electrocardiogram and cardiomegaly in digital x-ray of chest (posterior - anterior view). These abnormalities lead to further detailed investigation and 2D Echo revealed a moderate size 18 mm ostium secundum Atrial septal defect with left to right shunt, elongated, sail like anterior leaflet of tricuspid valve, 3 mm

Corresponding author: Lahiri Swati, Assistant Professor, Department of Anesthesiology, Medical College, Kolkata, India, Tel: +91 9836979351; E-mail: drslahiril1@gmail.com

Citation: Swati L, Chakraborty A, Dey M, Panja M, Dey S, et al. (2023) Anesthetic Concerns of Ebstein's Anomaly Patient Undergoing Lower Limb Orthopedic Surgery. *Int J Surg Invasive Procedures*, 7(1): 215-218.

Copyright: ©2023 Swati L, Chakraborty A, Dey M, Panja M, Dey S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

apical displacement of septal leaflet of tricuspid valve (diagnostic of Ebstein's anomaly), no evidence of Pulmonary artery hypertension at rest, Left ventricular ejection fraction - 58% and jerky septal wall motion. A 24 h Holter recording was started soon after diagnosis and

ventricular ectopic beats with a total burden of 1.7%, which were of no significance were found. Patient was scheduled for surgery after getting fitness from a cardiological point of view. He was premedicated with oral alprazolam 0.25 mg on the night before and on the morning of surgery (**Figure 1**).




Figure 1. A: 12 lead ECG with long lead II showing sinus rhythm, short PR interval, delta wave, widened QRS; B: Echocardiography showing atrialization of right ventricle, elongated, anterior leaflet of tricuspid valve, apical displacement of septal leaflet of tricuspid valve, atrial septal defect.

Monitoring on the operation table was done with electrocardiography (ECG), pulse oximetry, noninvasive blood pressure and urine output. One 18G intravenous line was established. Cardiologist was appointed for any adverse situation and emergency medications alongside antiarrhythmic medications (injection amiodarone) were kept ready. Prophylactic treatment for infective endocarditis with Injection (Inj) ceftriaxone 1g intravenous (IV) and antiemetic treatment with inj ondansetron 8 mg IV were given slowly. Precautions were taken to prevent air bubbles and paradoxical air embolism. For better analgesia distal sciatic nerve block in popliteal fossa and femoral nerve block in femoral triangle was performed under ultrasound guidance (E cube 8 ultra sound: probe L3-12T straight) with 100 mm 22g USG compatible nerve block needle with 0.2 % ropivacaine of 20 ml for distal sciatic nerve and 10 mL for femoral nerve. Supine position was maintained during surgery.

General anesthesia was given with inj Etomidate 24 mg IV after giving inj Fentanyl 150 mcg IV, inj midazolam 2 mg IV as premedication. Inj loxicard was administered to prevent laryngeal surge thereby reducing any chances of sympathetic stimulation. Inj Rocuronium 60 mg IV used as muscle relaxant induced agents. Direct laryngoscopy was done and intubated with an endotracheal tube with internal diameter of 8.5 mm and fixed at 22 cm. Patient was then put on a ventilator of following settings - Volume Control mode, Tidal volume 450 ml, I: E 1:2; No positive end expiratory pressure was used. End Tidal CO₂ (EtCO₂) maintained between 25 and 35 mmHg General anesthesia was

maintained in closed circuit with inhalational agent sevoflurane and muscle relaxation maintained with inj vecuronium 1 mg at 30 minutes intervals. For analgesia infusion Paracetamol 1g was given through intravenous route before skin incision. The operation lasted for 150 minutes and in total 1.5 liters of ringer's lactate solution was transfused during the surgery after proper assessment of fluid status through inferior vena cava diameter which was maintained around 1.7 cm. Intraoperatively ECG showed short PR interval, delta wave, widened QRS and two episodes of tachycardia (HR 110-120 beats per minute) which reverted back to normal sinus rhythm without any intervention and was maintained till the end of the surgery. The patient remained hemodynamically stable with systolic blood pressure in the range of (110-130) mm of Hg and diastolic blood pressure (80-90) mm of Hg, with well-maintained saturation in the intraoperative period. Total urine output and blood loss during the intraoperative period was 500 mL and 600 mL respectively (**Figure 2**).

Patient was extubated with a stable arterial pressure, heart rate and a good urine output. Inj Loxicard and inj Neostigmine methyl sulfate 2.5 mg and inj glycopyrrolate 0.5 mg were administered during extubation. One episode of tachycardia was noted however no hypotension was observed even shortly after removal of the endotracheal tube. After the surgery the patient was shifted to the Post Anesthesia Care Unit (PACU) for 24 h for hemodynamic observation. Pain control was also assessed using Visual analogue scale (VAS) score in PACU and it was around 0-4 during the first 24 h of the postoperative period. No

antiarrhythmic medication was used postoperatively however inj Tramadol 100 mg intravenous for three times was scheduled for better pain management. Patient was

shifted from PACU to general ward after 24 h of close monitoring with stable hemodynamic condition.



Figure 2. Intraoperative vitals of the patient.

DISCUSSION

The goal of anesthesia in this patient is to maintain sinus rhythm and both preload and afterload. The major problems associated with Anesthesia in Ebstein's anomaly are the risk of arrhythmias and the possibility of decreasing the arterial oxygen tension further by an increase of the right to left shunt. Subarachnoid blocks and general anesthesia can cause sudden decrease in systemic vascular resistance which can potentially worsen the right to left cardiac shunt. We avoid Spinal anesthesia as it can lead into sharp fall in systemic vascular resistance which causes sudden hypotension. Sudden hypotension has a major effect on cardiac output as well as contractility of the right atrium and RV. General anesthesia is also associated with increased catecholamine levels and intrathoracic pressure. These can increase intracardiac shunt [4]. However, precautions before surgery, reduced intubation time, inhibition of sympathetic stimulation like laryngeal surge, watchful monitoring, taking good care of fluid status, maintaining deep plane of anesthesia and adequate analgesia make general anesthesia effective for lower limb surgery in a case of Ebstein's Anomaly. Here in this case, we use ultrasound guided direct visualization of the needle, nerve, and any nearby vascular structures which minimizes the chance of nerve injury, local

anesthetic toxicity, and accidental puncture of a blood vessel. By avoiding epidural analgesia, we minimize the chances of hypotension, nerve damage, epidural hematoma, spinal cord infection and accidental dural puncture.

Hannam [5] found that etomidate as an induction agent didn't cause reduction in mean arterial pressure (MAP) integral from baseline after induction of anesthesia and less frequent use of vasopressors [5]. Use of vasopressors with strong beta-adrenergic stimulant action is better to avoid in this context as they have direct effect on myocardium. Wierda, Schuringa and van den Broek [6] found that Rocuronium because of its weak vagolytic effect in its intubating dose apart from hemodynamic effects related to muscle paralysis, did not lead to increase in heart rate without change in MAP. We avoid Succinylcholine as muscle relaxant which further decreases any possibility of hyperkalemia induced arrhythmia. In this case central venous catheterization was not performed as it has higher propensity for tachyarrhythmias and paradoxical embolism [7]. Every effort was made to prevent any excess fluid administration which can lead to an increase in right to left shunt and hypoxia. Hypercapnea and hypercapnea induced hypotension inhibited by maintaining EtCO₂ in the range of 25-35 mm of Hg.

Therefore, we conclude that anesthetic challenges in patients with Ebstein's anomaly depend on physical status of the patients, type of surgery, mode of analgesia and monitoring techniques [8].

DECLARATION OF PATIENT CONSENT

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient has given his consent for his images and other clinical information to be reported in the journal. The patient understands that his name and initials will not be published and due efforts will be made to conceal his identity, but anonymity cannot be guaranteed.

REFERENCES

1. Fuchs MM, Connolly HM (2020) Ebstein Anomaly in the Adult Patient. *Cardiol Clin* 38(3): 353-363.
2. Holst KA, Connolly HM, Dearani JA (2019) Ebstein's Anomaly. *Methodist Debaquey Cardiovasc J* 15(2): 138.
3. Stephens EH, Dearani JA, Qureshi MY, Ammash N, Maleszewski JJ (2020) The Congenital Tricuspid Valve Spectrum: From Ebstein to Dysplasia. *World J Pediatr Congenit Heart Surg* 11(6): 783-791.
4. Lobato EB, Florete OG, Paige GB, Morey TE (1998) Cross-sectional area and intravascular pressure of the right internal jugular vein during anesthesia: Effects of Trendelenburg position, positive intrathoracic pressure, and hepatic compression. *J Clin Anesth* 10(1): 1-5.
5. Hannam J, Mitchell S, Cumin D, Frampton C, Merry A, et al. (2019) Haemodynamic profiles of etomidate vs propofol for induction of anesthesia: A randomized controlled trial in patients undergoing cardiac surgery. *Br J Anaesth* 122(2): 198-205.
6. Wierda JM, Schuringa M, van den Broek L (1997) Cardiovascular effects of an intubating dose of rocuronium 0.6 mg kg⁻¹ in anaesthetized patients, paralyzed with vecuronium. *Br J Anaesth* 78(5): 586-587.
7. Horishita T, Minami K, Koga K, Ogata J, Sata T (2005) Anesthetic management using echocardiography for surgery of lower extremity in a patient with Ebstein's malformation. *Anesth Analg* 101: 606-615.
8. Sinha P, Kumar B, Varma P (2010) Anesthetic management for surgical repair of Ebstein's anomaly along with coexistent Wolff-Parkinson-White syndrome in a patient with severe mitral stenosis. *Ann Card Anaesth* 13(2): 154.