

## Is ABO Blood Groups Relevant to COVID-19 Infection?

Francis Ajeneye\*

*\*Department of Blood Transfusion, Tunbridge Wells Hospital, NHS Trust TN2 4QJ, United Kingdom.*

*Received August 27, 2020; Revised September 24, 2020; Accepted January 28, 2021*

### ABSTRACT

Evaluation of peripheral blood morphology is an important screening tool for many diseases. When abnormalities are detected by the automated hematology analyzer, manual microscopic review of the blood smear is necessary to determine the next course of action. Thus, the International Consensus Group for Hematology Review prepare criteria to review peripheral blood slide after analyzing by hematology analyzers. For better advantage of this criterion laboratories must optimize it with their setting.

**Keywords:** COVID-19, ABO blood group, Hematology

### INTRODUCTION

The association of Covid-19 infection and blood groups remains elusive and must be approach with vigilance. The research designs and methodologies identified in recent literatures are not robust and convincing. A sound research design that identifies dependent and independent variables and adjust for confounding factors could be a way forward. However, Covid-19 infection could be the results of complex interactions of factors that could vary from genetics, behavioural, metabolic, psychological, social status and environmental risk factors.

Conversely, as suggested by Guillon and colleagues [1] that variable susceptibility to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections could be linked to circulating anti-A antibodies, and could interfere or even reduce the virus-cell adhesion process. This could be considering as a working hypothesis.

A genome-wide association study (GWAS) [2,3] found that blood type O carriers have increased IL-6 levels than non-O blood types, suggesting the advantages of blood type O over the other types in maintaining the dominant role of Angiotensin-converting enzyme-2 (ACE2) in the RAS and thus a reduced risk of developing hypertension. The GATC haplotype of the four polymorphisms of the ABO gene (rs8176746, rs8176740, rs495828, rs12683493), which is common among non-O blood type patients, was positively associated with ACE activity [2].

In addition to inflammatory response associated with Covid-19 infection [3], we cannot overlook mechanism purported by earlier studies that suggested ABO blood group profound influence on the haemostasis, as it is a major determinant of

plasma levels of Von Willebrand Factor (VWF) [4,5]. Literatures also defined blood group O as a risk factor for increased severe bleeding while blood group non-O is a risk factor for thromboembolic events. The risk of VTE is probably related to the level of VWF and factor VIII in non-group O subjects. A, B, and H blood group antigens are expressed on N-glycans of vWF and influence the half-life of the protein (10 hours for group O and 25 hours for non-O subjects), duchies explain for the greater levels in non-O patients. These observations raise the prospect that a greater tendency for blood clot formation in non-O patients [6].

The association between antigens A and B and arterial thrombosis, such as coronary heart disease (CHD), cerebrovascular disease or peripheral vascular disease, is still unclear. Canadian researchers evaluated the association between blood groups and thrombotic events in a cohort of blood donors from the province of Quebec, Canada. When the analysis was controlled to older women ( $\geq 40$  years), those with blood group A had a statistically significant 40% higher risk of CHD events, compared to those with group O (HR 1.40 [95% CI: 1.01-1.92]) [7].

Thromboembolic events (blood group O vs. others) occurred

**Corresponding author:** Francis Ajeneye, Associate Lecturer in Biomedical Sciences, Department Blood Transfusion, Senior Biomedical Scientist, Pathology, Maidstone and Tunbridge Well Hospitals, NHS Trust, TN2 4QJ, UK, Tel: 01892636063; E-mail: f.ajeneye1@nhs.net

**Citation:** Ajeneye F. (2021) Is ABO Blood Groups Relevant to COVID-19 Infection? J Blood Transfusions Dis, 4(1): 178-179.

**Copyright:** ©2021 Ajeneye F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

in 2.8 (O) vs. 3.3 (others) %,  $P=0.619$ . The Canadian retrospective study on 1281 multiple injured patients, there found no relevant influence of ABO blood group on haemorrhage, thromboembolic events and mortality.

They found Blood group AB was associated with a higher risk of hospitalisation or death because of thrombotic events such as coronary, cerebrovascular or peripheral diseases.

However, in another review, one justification for the association of ABO blood group and CHD was directed towards elevated levels of Von Willebrand Factor (VWF) and consequently, of factor VIII in the plasma, as a risk factor for CHD and also towards variants at ABO loci associated with increased levels of plasma lipid and inflammatory markers. Some studies have found conflicting results [2,8-10].

Other mechanisms have been proposed to explain the association between blood group and CHD, but a unifying theory remains elusive as discussed by Zhou et al. in their recent review [11].

An international database, encompass with a robust research design could unlock the biology of Covid-19 and its association with blood groups which eventually will improve our understanding of human susceptibility to the new emerging infection [12].

## REFERENCES

- Guillon P, Clement M, Sebille V, Rivain J-G, Chou C-F, et al. (2008) Inhibition of the interaction between the SARS-CoV Spike protein and its cellular receptor by anti-histo-blood group antibodies. *Glycobiol* 18: 1085-1093.
- Luo JQ, He FZ, Luo ZY, Wen JG, Wang LY, et al. (2014) Rs495828 polymorphism of the ABO gene is a predictor of enalapril-induced cough in Chinese patients with essential hypertension. *Pharmacogenet Genom* 24: 306-313.
- Naitza S, Porcu E, Steri M, Taub DD, Mulas A, et al. (2012) A genome-wide association scan on the levels of markers of inflammation in Sardinians reveals associations that underpin its complex regulation. *PLoS Genet* 8: e1002480.
- Li J, Wang X, Chen J, Cai Y, Deng A, et al. (2020) Association between ABO blood groups and risk of SARS-CoV-2 pneumonia. *Br J Haematol* 2020 [Epub ahead of print].
- Gill JC, Endres-Brooks J, Bauer PJ, Marks WJ, Montgomery RR (1987) The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood* 69: 1691-1695.
- Dentali F, Sironi AP, Ageno W, Bonfanti C, Crestani S, et al. (2013) Relationship between ABO blood group and hemorrhage: A systematic literature review and meta-analysis. *Semin Thromb Hemost* 39: 72-82.
- Blais C, Germain M, Delage G, Grégoire Y (2016) The association between blood group and the risk of vascular disease in Quebec blood, *Blood Transfusion* 14: 455-459.
- Jenkins PV, O'Donnell JS (2006) ABO blood group determines plasma von Willebrand factor levels: A biologic function after all? *Transfusion*. 46: 1836-1844.
- Dentali F, Sironi AP, Ageno W, Turato S, Bonfanti C, et al. (2012) Non-O blood type is the commonest genetic risk factor for VTE: Results from a meta-analysis of the literature. *Semin Thromb Hemost* 38: 535-548.
- Zhou S, Welsby I (2014) Is ABO blood group truly a risk factor for thrombosis and adverse outcomes? *World J Cardio* 16: 985-992.
- Zhao J, Yang Y, Huang H, Li D, Gu D, et al. (2020) Relationship between the ABO Blood Group and the COVID-19 Susceptibility. *MedRxiv* [Epub ahead of print].
- Franchini M, Capra F, Targher G, Montagnana M, Lippi G, et al. (2007) Relationship between ABO blood group and von Willebrand factor levels: from biology to clinical implications. *Thromb J* 5:14.